

Variation in the yield and essential oil of four chamomile varieties grown in Finland in 1985—1988

B. GALAMBOSI¹, ZS. SZEBENI-GALAMBOSI¹, M. REPCAČ² and P. CERNAJ²

¹ Agricultural Research Centre, South Savo Research Station, Karila, SF-50600 Finland

² Dept. of Special Biology, P.I. Safarik University of Kosice, Manesova 23, 04167 Kosice, Czechoslovakia

Abstract. Four chamomile (*Matricaria recutita* L.) varieties were grown at Puumala, Finland (61°40' N, 28°15' E) in 1985—1989 from spring sowings on stony till soil, pH 5.8. The weather conditions in the experimental years were very different. For instance between 1987 and 1988 there was a difference of 5.3°C in the mean temperature of the growing periods (May–September). The four-year average of the fresh flower yield was 0.4 (0.1–0.7) kg/m². The yields varied greatly depending on the number of harvests, which depended on the weather conditions. One harvest could be obtained in 1986 and 1987, two in 1985 and three in 1988.

In spite of the different weather conditions there were no differences in the content and composition of the essential oil distilled from the dry flowers.

Index words: chamomile, varieties, flower yield, essential oil

Introduction

Chamomile (*Matricaria recutita* L.) is a native plant of the Finnish flora, but it occurs quite rarely due to the widespread use of herbicides in agriculture and on roadsides. All chamomile used in the pharmaceutical industry and sold in health shops a total of 8.7 t to a value of FIM 521.000 was imported in 1982 (HÄLVÄ 1985).

Field experiments for production of chamomile were started in 1984 within the herb cultivation project of the University of Helsinki,

Finland, organised by the Department of Horticulture and the Pharmacognosy Division (GALAMBOSI 1989). In addition to the agrotechnical experiment, observations on different cultivars and chemotypes have been started in order to study the effects of the climatic conditions on the herb yield and essential oil. So far there are no long time observations on chamomile cultivated in the Finnish climate. The very few experiments with chamomile in Finland are concerned with the accumulation of essential oil and the chamazulene content during ontogenesis (VON SCHANTZ and SALO-

NEN 1966), or with the effects of various climatic factors on chamomile in comparative pot trials in Finland (Helsinki), central Europe (Münich), and Asia Minor (Ismir, Turkey) reported by FRANZ *et al.* (1986). The latest observations were carried out in 1984 during a one year experiment in which the morphological characters, yield and essential oil content and composition of seven chamomile species grown in parallel in Hungary and South-Finland were studied (GALAMBOSI *et al.* 1988).

The purpose of this work was to study the variation of flower yield and essential oil of four chamomile varieties during a longer period under normal field conditions in Finland.

Material and methods

The experiments were carried out at Puumala (61°40' N, 28°15' E) in a stony till soil, pH 5.8. The agricultural methods were the same each year and similar to the general Hungarian methods (SVÁB 1978). The chamomile was sown in spring using 5 kg/ha seed (5 g/10 m²) with diffuse surface sowing. The plot size was 10 m² in two replications. The plots were fertilized with 50—80—85 kg/ha nitrogen, phosphor and potassium at the time of soil preparation.

Harvesting was carried out with a hand-harvester, the flower and herb yields (stems with flowers) being separated by a 18 mm screen. During the four growing seasons one harvest was gathered in 1986 and 1987, two in 1985 and three in 1988. The flower and herb samples were dried at 30°C for 12 h and stored in paper bags at room temperature. The essential oil analyses were carried out in the laboratory of the Department of Special Biology, University of Kosice, Czechoslovakia, in 1986—1988 from the first harvests, in 1988 from the second harvest as well. The essential oil was obtained by 2 h distillation of 2 g dry samples of the ground drug in three replications. The content of essential oil was determined gravimetrically (HÖLZ and DEMUTH 1975). The main components of the essential

oil were determined by means of gas chromatography (REPCAK *et al.* 1980).

The chamomile varieties used in the experiment were Budakalasz-2 (tetraploid, Hungary), Degumille (diploid, FRG), Csömöri (diploid, Hungary) and Bona (diploid, CSSR). Some morphological characters of the varieties, the sowing times and length of the growing periods, are presented in Tables 1—2.

The data on the essential oil content and composition was studied by analyses of variance (SVÁB 1981).

The meteorological conditions during the experimental periods as measured by the Finnish Meteorological Institute, are presented in Figure 1. The climatic conditions were quite different in each year. In 1986 and 1988, the average temperature of the growing season was higher than 30 year average. In May-June-July the average temperature was 2 to 4°C higher than the long term average. The effective temperature sum (calculated from the sum of the mean daily temperatures above +5°C between May 15th and August 31th) was 1072°C in 1986 and 1168°C in 1988.

In 1987, on the other hand, the growing season was colder and rainier than the 30 year average, the temperature being 2 to 3°C lower and the precipitation 130 mm higher than the long-term average. The effective temperature sum was only 750°C. The temperature in 1985 did not differ too much from the average, the effective temperature sum being 928°C, but the second half of the growing period was very rainy. Thus the chamomile

Table 1. Length of the vegetation periods of chamomile during 1985—1988 at Puumala.

Year	Sowing date	Length of periods in days from sowing to		
		First harvest	Second harvest	Third harvest
1985	17.5.	77	98	—
1986	15.5.	64	—	—
1987	28.5.	84	—	—
1988	16.5.	59	78	115
	mean:	71	88	—




Fig. 1. Mean temperature (A) and precipitation (B) at Puumala during the experimental years and long-term averages (1930-1960).

Table 2. Height and morphological characters of flowers of chamomile varieties.

Variety	Plant height at the first harvest time (cm)				Torus diameter of flower head (cm) (1985)	100 flower mass (g)	
	1985	1986	1987	Mean		fresh (1985)	dry (1985)
Budakalasz-2	72.6	54.8	63.7	63.7	0.955	28.1	5.27
Degumille	60.8	47.6	49.5	52.6	0.825	15.0	2.81
Csömöri	57.0	55.2	47.5	53.2	0.820	16.0	3.00
Bona	52.8	54.9	46.2	51.3	0.865	15.1	2.81

(figures are means of 20 plants, 50 flowerheads and fresh and dry mass of 10 × 100 flowerheads)

varieties were grown under very different weather conditions.

Results and discussion

Length of the growing periods

The average length of the period from sowing to the first flower harvest was 71 days, but in 1986 and 1988 it was shorter due to the warm and dry weather (Table 1). In 1987, as a consequence of the cold and rainy summer, the first harvest was collected 84 days after sowing, and a second flowering did not appear. The spring sowings can generally be made in the middle of May after melting of the snow, when the soil surface is rather dry.

The length of the growing periods of chamomile depends on the weather conditions: if the spring sowing is successful and there is enough precipitation for second and third flowerings it can be 100–115 days long.

Flower and herb yield

The fresh flower and herb yields are presented in Tables 3–4 and in Figure 2. The quantities were determined by the number of harvests and by the weather conditions.

The four year average yield of fresh flower was 0.4 kg/m². From only one harvest the yields were generally low (0.17 kg/m²), especially in 1986 and 1987. If second and third harvests were available, the yields reached 0.6–0.7 kg/m². The yields from spring sowings are at the same low level as reported by SVÁB (1978) and ZALECKI (1978). There was no connection between the flower or herb yield and the temperature sum of the growing season. The highest yields were achieved in 1985 and 1988 with low and high temperature sums (928 and 1168°C, respectively), and the lowest yields were obtained in 1986 and 1987, also with high and low values (1072 and 750°C). The critical factor affecting the yield is the precipitation in early spring. If there is enough precipitation in the period of the vegetative growth of the roots and rosettes,

the yield will be good as in 1985. In 1988, the warm summer caused continuous flowering and gave three harvests, but the yield remained lower due to lack of moisture in the soil in May.

The fresh herb yield obtained after separation of the flowers and which is suitable for distillation or extraction varied from 0.03 to 0.60 kg/m², being on average 0.20 kg/m². The quantitative herb yield of the second harvest was affected by the high precipitation in July, 1985. The response of the different varieties to the varying weather conditions was different. The highest flower and herb yields were produced by the varieties Degumille and Budakalasz-2. The yield of Bona was lowest due to the low flower mass. The average proportion of flowers in the total phytomass varied from 30 to 38 %, which is a little higher than the results of GASIC *et al.* (1989).

Content and composition of the oil

The oil contents of Budakalasz-2 and Degumille were stable 1 %, and no changes were observed from year or between the first and second harvests in 1988 (Table 5). The average essential oil content of Csömöri and Bona was 0.9 %. In 1986 and 1987, the oil content of Csömöri was 0.8 %, the oil content of Bona in 1986 and in the second harvest of 1988 was 0.7 % and 0.8 %, respectively ($p < 0.1$).

The variation of the main components of the essential oil is shown in Table 5. There were clear differences between the varieties: Degumille, Csömöri and Bona seemed to be α -bisabolol types (35–60 %), while Budakalasz-2 is a bisabololoxide type cultivar. There were no big differences in the content of the main components caused by the very different weather conditions during the growing periods. The chamazulene content of the oils varied between 11 and 21 %. Some variation can be observed in the content of *trans*- β -farnesene, the content being highest in the warmest year 1988 and especially in the second harvest. This observation is in accordance




Fig. 2. Fresh flower yield of the chamomile varieties at Puumala 1985—1988. Roman numerals in the columns indicate first, second and third harvest.

with results of GASIC *et al.* (1989), who reported the highest farnesene content of 16 chamomile varieties in the warmest year stud-

ied. GALAMBOSI *et al.* (1988) stated the same tendency in a comparative study of 6 chamomile varieties grown in Hungary and Finland.

Table 3. Fresh flower yield (g/m^2).

Variety	1985			1986	1987	1988				Mean		
	I	II	I+II	I	I	I	II	III	I+III	All together	From I harvests only	From I-II-III harvests
Budakalasz-2	341	453	794	107	146	314	149	96	558	401	227	676
Degumille	200	1048	1248	100	91	154	329	306	789	557	136	1018
Csömöri	112	640	752	103	183	332	192	57	581	404	182	666
Bona	136	180	326	124	127	261	250	41	552	282	162	439
Mean	197	580	780	108	137	265	230	125	620	411	177	699

Table 4. Fresh herb yield after separation of the flowers (g/m^2).

Variety	1985			1986	1987	1988				Mean		
	I	II	I+II	I	I	I	II	III	I+III	All together	From I harvests only	From I-II-III harvests
Budakalasz-2	176	356	432	96	56	184	76	40	300	221	128	366
Degumille	137	466	603	70	39	90	169	130	389	275	84	496
Csömöri	92	259	351	110	97	194	98	24	316	218	123	333
Bona	41	56	97	101	35	153	128	17	298	132	82	197,5
Mean	111	284	370	95	56	155	118	53	326	211	104	348

Table 5. Content and main compounds of the essential oil of different chamomile varieties 1986—1988.

Year	Variety	Oil	<i>trans</i> - β -farnesene	Chamazulene	α -bisabolol	Bisabololoxide		Bisabolonoxide
		%	%	%	%	A %	B %	%
1986	Budakalasz-2	1.06	17.40	15.73	3.33	39.00	5.13	3.66
	Degumille	1.12	15.53	17.90	40.93	6.16	7.50	+
	Csömöri	0.82	11.63	16.70	57.56	1.20	2.40	+
	Bona	0.69	14.00	11.56	60.10	1.40	2.16	+
	X	0.92	14.64	15.47				
1987	Budakalasz-2	1.08	29.60	12.00	1.30	37.80	3.94	0.40
	Degumille	1.04	30.70	17.80	34.60	5.90	2.90	+
	Csömöri	0.87	23.80	15.40	54.80	—	0.20	+
	Bona	1.01	19.90	17.90	50.10	0.50	—	+
	X	1.00	26.00	15.77				
1988 (a)	Budakalasz-2	1.06	37.30	17.60	1.25	33.40	5.10	1.85
	Degumille	1.16	27.60	20.00	37.40	2.45	2.45	+
	Csömöri	1.03	43.75	10.75	38.60	0.75	0.45	+
	Bona	1.09	33.55	16.90	44.35	0.39	0.65	+
	X	1.08	35.67	16.31				
1988 (b)	Budakalasz-2	1.07	40.65	16.95	1.55	35.30	2.35	0.95
	Degumille	1.16	34.75	20.15	37.90	2.20	2.05	+
	Csömöri	1.00	42.75	18.25	32.40	1.65	0.15	+
	Bona	0.84	37.75	21.75	33.35	0.65	0.10	+
	X	1.02	38.97	19.27				
	p <	0.1 % 5 % (1987)	0.1 % 5 % (1986)	0.1 %	0.1 %	0.1 %	0.1 %	n.c.

+ = traces
n.c. = not calculated
a = first harvest
b = second harvest

In Hungary the β -farnesene content was higher when the summer was warm.

The four-year experiment on the cultivation of chamomile in Finland carried out under very different weather conditions confirmed the results reported earlier by SVÁB *et al.* (1967), HONCARIV *et al.* (1979) and FRANZ *et al.* (1986). The production of phytomass and the flower yield are influenced by climatic factors rather than by the genetically determined chemical characters of the different varieties or chemotypes.

FRANZ *et al.* (1986) stated that the oil formation is related to the flower formation.

Since the summer conditions in Finland are suitable for the flower formation of chamomile, there were no substantial changes in the content and composition of the essential oil during the four years of cultivation. The flower yield depends on the field conditions and suitable cultivation techniques. Growers must use the appropriate growing technique for chamomile, especially in the spring period, elaboration of such methods requires more detailed research.

Acknowledgement. The authors wish to thank Y. Holm (Ph.D.) for valuable advice during the preparation of this paper.

References

- FRANZ, Ch., HÖLZ, J. & VÖMEL, A. 1978. Preliminary morphological and chemical characterization of some populations and varieties of *Matricaria chamomilla* L. Acta Hort. 73: 109—114.
- FRANZ, Ch., MÜLLER, E., PELZMANN, H., HÄRDH, K., HÄLVÄ, S. & CEYLAN, A. 1986. Influence of ecological factors on yield and essential oil of chamomile (*Chamomilla recutita* (L.) Rauschert syn. *Matricaria chamomilla* L.). Acta Hort. 188, Medicinal and Aromatics V. 157—162.
- GALAMBOSI, B., MARCZAL, G., LITKEY, K., SVAB, J. & PETRI, G. 1988. Comparative examination of chamomile varieties grown in Finland and Hungary. Herba Hung. 27: 45—55.
- GALAMBOSI, B. 1989. Phytomass production of Medicinal Plants in Finland. Acta Agron. Hung. 38: 89—97.
- GASIC, O., LUKIC, V., ADAMOVIĆ, R. & DURKOVIC, R. 1989. Variability of content and composition of essential oil in various chamomile cultivars (*Matricaria chamomilla* L.) Herba Hung. 28: 21—28.
- HONCARIV, R. & REPCAK, M. 1979. Chemotypes of *Matricaria chamomilla* L. Herba Polonica. XXV: 261—267.
- HÖLZ, J. & DEMUTH, G. 1975. Einfluss ökologischer Faktoren auf die Bildung des ätherischen Öls der Flavone verschiedenen Kamillenherkunfte. Kritischer Vergleich der quantitativen Bestimmungsmethoden. Planta Medica 27: 37—45.
- HÄLVÄ, S. 1985. Consumption and production of herbs in Finland. J. of Agric. Sci. in Finland. 57: 231—237.
- REPCAK, M., HALASOVA, J., HONCARIV, R. & PODHRADSKY, D. 1980. The content and composition of essential oil in the course of anthodium development in wild chamomile (*Matricaria chamomilla* L.). Biol. Plant. (Praha) 22(3): 183—191.
- SCHANTZ, M. & SALONEN, R. 1966. Untersuchungen über den Ölgehalt und die Gesamtazuleenmenge während der Entwicklung der Blütenkörbchen von in Finnland wildwachsender Kamille (*Matricaria chamomilla* L.). Sci. Phar. 34: 177—185.
- SVÁB, J., C. EL-DIN AWAAD & FAHMY, T. 1967. The influence of highly different ecological effects on the volatile oil content and composition in the chamomile. Herba Hung. 6: 177—199.
- SVÁB, J. 1978. Kamilla (*Matricaria chamomilla* L.) in: ed. Hornok, L.: Gyógynövények termesztése és feldolgozása. p. 248—254. Budapest.
- SVÁB, J. 1981. Biometrical methods in research work. Budapest.
- ZALECKI, R. 1978. Organizacja uprawy rumianku pospolitego na surowiec. Wiadomosci Zielarskie. 7: 2—2.

Ms received June 14, 1990

SELOSTUS

Neljän kamomillalajikkeen kukkasato ja haihtuvien öljyjen pitoisuudet Suomessa v. 1985—1988

B. Galambosi¹, Zs. Szebeni-Galambosi¹, M. Repcak² & P. Cernaj²

¹ Maatalouden tutkimuskeskus, Etelä-Savon tutkimusasema Karila, SF-50600 Mikkeli, Finland.

² P.J. Safarik -yliopiston Erikoisbiologinen laitos, Manesova 23, CS-04167 Kosice, Czechoslovakia

Vaikka kamomilla (*Matricaria recutita* L.) kasvaa Suomessa luonnonvaraisena rohdoskasvina käyttävät lääketeollisuus ja luontaistuotekaupat ulkomailta tuotua kamomillaa. Hälvän (1985) selvityksen mukaan maamme tuottiin esim. v. 1982 8700 kg kamomillan kukkaa, jonka arvo oli 521 000 mk. Lääketeollisuus vaatii kamomillalle tiettyä kemiallista koostumusta, joten eri

kemotyyppien tutkimus on välttämätöntä pohjoisissa ilmasto-olosuhteissa.

Kamomillan viljelytutkimus kuului Helsingin yliopiston v. 1984 aloittamaan Puumalan mausteprojektiin. Siinä tutkittiin Suomen olosuhteisiin sopivia viljelymenetelmiä ja niiden koneellistamista sekä eri lajikkeiden viihtyvyyttä.

Pirttimäen koetilalla tutkittiin v. 1985—1988 neljän

lajikkeen satoisuutta ja viihtyvyyttä. Lajikkeet olivat unkarilainen tetraploidi Budakalasz-2 sekä diploidit länssisaksalainen Degumille, unkarilainen Csömöri ja tšekkoslovakialainen Bona. Kamomillan siemeniä kylvettiin keväällä 5 kg/ha hajakylvönä maan pinnalle muokattuun maahan 10 m²:n koeruutuihin. Kokeessa oli kaksi kerannetta, maalajina hiekkamoreeni ja maan pH-arvo 5,8. Koealueelle levitettiin typpeä 50 kg/ha, fosforia 80 kg/ha ja kaliumia 85 kg/ha. Kukkasato korjattiin käsipuimurilla riipien ja lajiteltiin 18 mm:n seulalla. Kaikki kukat eivät irtoa puinnin ja lajittelun aikana. Osa sadosta muodostuu varsista, joissa kukat ovat yhä jäljellä. Tämä ns. kamomillan heinäsato on sopivaa kemialliseen jatkojalostukseen uttamis- ja tislaamisen menetelmien avulla.

Viljelyvuosien sääolosuhteet vaihtelivat paljon. Vuosina 1986 ja 1988 kasvukausi oli 2—4°C keskimääräistä lämpimämpi ja sademäärä vastaavasti keskimääräistä pienempi. Vuonna 1987 kasvukausi oli 3°C keskimääräistä kylmempi ja myös sateinen. Kasvukauden sää vaikutti merkittävästi kamomillan kukka- ja heinäsatoon. Vuon-

na 1985 saatiin korjattua kaksi satoa, mutta kahtena seuraavana vuonna vain yksi onnistunut sato. Paras tuotos saavutettiin v. 1988, jolloin satoa korjattiin kolme kertaa.

Lajikkeiden tuore kukkasato oli neljän vuoden keskiarvona 0.4 kg/m² ja tuore heinäsato 0.2 kg/m². Tämä vastaa keskieurooppalaista kevätkylvöstä saatua satoa, jota saadaan yleensä vähemmän kuin syyskylvöstä.

Kuivattujen kukkien kemiallinen analyysi osoittaa, että tetraploidi lajike on bisabololoksidi-A -tyyppiä ja diploidit lajikkeet ovat alfa-bisabololi -tyyppiä. Lajikkeiden haihtuvan öljyn pitoisuuksissa ja koostumuksessa ei tapahtunut merkittäviä muutoksia, vaikka viljelyvuosien sääolosuhteet vaihtelivatkin paljon.

Tämä viljelytutkimus antoi näytön siitä, että eteläisessä Suomessa on mahdollista saada hyvälaatuinen kamomillan kukkasato. Tulevaisuudessa onkin tarpeellista tutkia myös kamomillan syyskylvöä ja sen vaikutusta sadon määrään.