

Bryology in Serbia: from floristics to chemosystematics

Review Article

Milan Veljić

University of Belgrade, Faculty of Biology, Institute of Botany and Botanical Garden, Takovska 43, 11000 Belgrade, Serbia
veljicm@bio.bg.ac.rs (corresponding author)

Abstract:

Bryological research in Serbia can be divided into several periods and research fields, the first of which being floristic research. An important exploration of bryophytes in Serbia started at the end of the 19th century, and with a few discontinuances it continued up today. During the last decade of the 20th century couple of researchers started their work on the exploration of bryophytes. They focused on bryologically important habitats like sources, riverbanks, peat bogs and mountains. A significant number of new species was reported for the country, but the revision of the old records has been done as well, hence today flora of Serbia numbers 831 bryophyte taxa in total. Progress of the botanical research has brought interdisciplinarity into bryology. As a result, bryology in Serbia nowadays spread in different types of bryophyte research. For example, bryophytes are studied to investigate the biological, antimicrobial, antifungal and cytotoxic activity, biotransformation of molecules, phytochemical composition and as model organisms in stress physiology. Special emphasis is also given on active conservation of the species. This represents only some directions of what is currently studying on bryophytes in Serbia.

Key words:

bryophytes, Serbia, history, floristics, phytochemistry, biological activity

Apstract:

Kolekcija mahovina BEOU - zanemareno nacionalno istraživačko blago

Istoriju briologije u Srbiji možemo podeliti na nekoliko perioda i pravaca. Prvi je pravac florističke (inventarizacione) briologije. Naime značajnija briološka istraživanja u Srbiji započeta su krajem devetnaestog veka i sa znatnim prekidima traju do danas. U poslednjoj dekadi dvadesetog veka pojavilo se nekoliko istraživača koji su se prevashodno posvetili istraživanju mahovina. Istraživanja su usmerena na briološki zanimljiva staništa kao što su vrela, reke, tresetišta, planine. Otkriven je značajan broj novih taksona ali izvršena je i revizija nekih ranijih podataka, tako da trenutno brioflora Srbije broji 831 takson. Kako je opšti način botaničkih istraživanja napredovao, i u briologiji se prešlo na multidisciplinarnost. Grupa mlađih istraživača bavi se istraživanjima biološke aktivnosti mahovina, antimikrobne, antifungalne i citotoksične, fitohemijom mahovina, biotransformacijom molekula izolovanih iz mahovina ali i fiziologijom stresa. Poseban akcenat je stavljen na aktivnu zaštitu i reintrodukciju. Ovo je samo deo onoga što se danas iz Briologije radi u Srbiji.

Ključne reči:

briofite, Srbija, istorija, floristika, fitohemija, biološka aktivnost

Introduction

Bryological research in Serbia could be divided into three periods of investigation, and several research fields. First period coincide to the onset of the botany in Serbia, or with the work of Josif Pančić and his students. This period is characterized mainly by floristic or inventory studies, with relatively little bryophyte records made. After a brief break in investigation, research slightly intensified during the period

from 1920s to 1960s with the work of group of phytocenologists. Third, or the modern period, began in the end of 20th century, and is the most fruitful period so far (Pantović & Sabovljević, 2017).

During this period, in addition to extensive floristic research, ecological investigations, the biological activity of moss extracts, their phytochemical analysis, as well as the transformation of molecules isolated from the extracts, were also initiated and successfully realized. At present, a lot of research is



conducting in the field of stress physiology in Serbia, as well. A group of researchers is intensively working on the conservation of bryophytes, with an emphasis on active protection and reintroduction of the rare and threatened species. The Institute of Botany and Botanical Gardens “Jevremovac” has valuable collection of over 260 species of living bryophytes from around the world, of which about 60% are regionally or globally endangered.

Floristic-ecological research of the bryophytes of Serbia

As a part of general botanical research in the territory of Serbia, research of the bryophyte flora had started in the middle of the 19th century. The first bryophyte records were published in the works of the researchers of that time (Grisebach, 1843; Pančić, 1859, 1863). More extensive research of bryophyte flora has began at the end of the century when Svetozar Obradović, Đura Ilić and Lujo Adamović started researching bryophytes; they were sending most of the collected material for determination and verification by more experienced botanists (Wettstein, 1890; Simić, 1892; Matouschek, 1899, 1901). Simić (1897, 1898) published part of his research on the Vranje surrounding in the reports of the Vranje Gymnasium. Schiffner (1897) identified one species from the Bornmuller’s herbarium. Research results obtained during this period were summarized by Jurišić (1900) and Simić (1900). In these papers, they included part of their unpublished results as well - Jurišić cites 209 and Simić 145 species. For the next ten years, bryological research was done mostly by Katić (1900, 1903, 1906, 1907a, 1907b i 1909) and Košanin (1909a, 1909b, 1910).

After this period, bryological studies were done by Podpera (1922), Černjavski (1929, 1932,

1937, 1938), Pichler (1931, 1939, 1940), Grebenschikov (1943, 1949, 1950), Soška (1949), Knapp (1944), Rudski (1949a, 1949b) and Rajevski (1951). Pavletić (1955) stated in his work “Prodromus of the bryophyte flora of Yugoslavia” 374 taxa (57 Marchantiophyta, 317 Bryophyta) to be present in Serbia.

Since the 1950s, only sporadic bryophyte records were published within the general floristic research (Gigov & Nikolić, 1954; Gigov, 1956; Slavnić, 1956; Pavletić, 1956, 1968; Janković & Janković 1962; Blečić & Tatić, 1962; Čolić et al., 1963; Pavlović, 1951, 1964; Mišić & Popović, 1960; Popović, 1966; Filipović, 1966; Guelmino, 1970, 1972; Erdeši, 1971; Babić, 1972; Tešić et al., 1979; Martinčić, 1980; Janković & Mišić 1980; Jovanović 1980; Janković & Stevanović, 1981; Mišić, 1982; Grgić, 1983; Gajić, 1983, 1986, 1988, 1989; Gajić & Karadžić, 1991; Petković et al., 1988, 1990, 1991; Mišić & Panić, 1989). Based on these studies, the list of bryophyte flora has been updated by Gajić et al. (1991), who cited 415 bryophyte species in Serbia (60 Marchantiophyta, 355 Bryophyta).

The third period (or modern period) of bryophyte research began in the last decade of the 20th century when several researchers devoted their research to the bryophytes: Randelović, 1994; Stevanović et al., 1995; Veljić et al., 1996, 2001a, 2001b, 2004, 2006, 2008, 2013, 2016a, 2016b; Veljić & Marin, 1997; Pavić et al., 1998; Sabovljević & Stevanović, 1999; Petković et al., 2000; Sabovljević, 1998, 1999, 2000, 2003a, 2003b, 2006; Papp & Sabovljević, 2001, 2002, 2010; Grdović & Blaženčić, 2001; Sabovljević & Cvetić, 2003; Pócs et al., 2004; Cvetić & Sabovljević, 2005; Papp & Erzberger, 2005; Grdović, 2005; Grdović & Stavretović, 2004, 2006; Sabovljević & Grdović, 2009; Papp & Erzberger, 2005, 2007a, 2007b, 2009; Erzberger & Papp, 2007; Papp et al., 2004, 2006, 2012a, 2012b, 2013, 2014a, 2014b, 2014c, 2016a, 2016b, 2016c; Randelović, Zlatković, 2010; Pantović & Sabovljević, 2013, 2017; Pantović et al., 2014; Ilić et al., 2015). Research was focused on biologically interesting and unexplored habitats such as springs, rivers, peatlands, mountains. This phase of intensive research of the flora and ecology of bryophytes is ongoing, and more recently two Ph.D. dissertations focusing on bryophytes in Serbia were done: “Biogeographic and ecological study of the bryophyte flora of Serbia” (Pantović, 2018) and “Diversity, distribution, differentiation of micro-habitats and the structure of moss communities of Fruška gora mountain” (Ilić, 2019). In this short period of time,



Fig. 1. Collection of bryophytes in the BEOU Herbarium. Shown are digitized specimens included in the Bryo database (photo J. Pantović)



Fig. 2. Reintroduction of *Henediella heimii* (marked red) and *Entostodon hungaricus* (marked yellow), propagated axenically *in vitro*, acclimatized and finally released to nature on potentially suitable site in Banat, Serbia! (photo M. Sabovljević)

a significant number of new taxa has been reported for the country, but also some previous data has been revised, so Serbia currently numbers 830 bryophyte taxa (1 Anthocerotophyta, 142 Marchantiophyta, 687 Bryophyta).

Historical data on the distribution and ecology of bryophytes has been collected and digitized into a unique database of bryophytes of Serbia, the Bryo database (**Fig. 1**). Updated lists of liverworts and mosses of Serbia have been prepared and are currently in press.

Protection and conservation of bryophytes

Bryophytes, like all other living organisms, are exposed to various negative environmental factors. They are threatened similarly as vascular plants, but the attention and concern for their conservation is much lower. Among bryophytes, there are taxa that are extinct, in different categories of threat, or rare, which has been confirmed by the increasing number of the regional Red Lists (Sabovljević et al., 2004; Marka et al., 2012; Hodgetts et al., 2019). Recognizing this global problem, a group of associates gathered around Marko and Aneta Sabovljević (BBGB - Bryophyte Biology Group Belgrade) has developed into the widely recognized group for bryophyte conservation (Sabovljević et al., 2009; Sabovljević et al., 2014; Vujičić et al., 2012). The Institute of Botany and Botanical Garden “Jevremovac” has an *in vitro* living collection of over 260 species of bryophytes from around the world, of which about 60% are regionally or globally endangered. Nearly 52%

of the species in the collection are on the European and regional red lists.

Special attention is given to the active protection, like reintroduction and population strengthening of the rare and endangered taxa. *In vitro* propagated and grown plants are being prepared for reintroduction to the potentially suitable habitats. Several species such as *Henediella heimii* and *Entostodon hungaricus* (**Fig. 2**) have been propagated this way and reintroduced to habitats in Banat, Serbia (Sabovljević et al., 2012a, 2012b, 2016, 2018).

Ecophysiological research of bryophytes in Serbia

With progress of the botanical research, a multidisciplinary approach has been adopted in bryology as well. A group of young researchers has started working on bryophyte stress physiology (Rajčić et al., 2015; Vujičić, 2016; Vujičić et al., 2016; Čosić et al., 2018). As a part of these studies, a doctoral dissertation entitled “Biochemical and ecophysiological properties of selected moss species during salt stress in the axenical conditions” (Vujičić, 2016) has been done. Monitoring the parameters of morphogenesis revealed that bryophytes and vascular plants respond differently to the short-term and long-term salt stress, indicating different survival strategies with the increased salinity of the microenvironment. The effect of salt stress is also observed by the change in the concentration of tocopherols in mosses (*Physcomitrella patens*, *Entostodon hungaricus* and *Henediella heimii*) (**Fig. 3**). Namely, tocopherol as a non-enzymatic component plays an important role in the moss response to the stress caused by increased NaCl concentration. The results of the study show that under long-term stress, the concentration of tocopherols increases on moderately saline media (50-100 mM NaCl) in *H. heimii*. The salt stress affects concentration of proline and the ratio of proline and free amino acids in mosses (*P. patens*, *E. hungaricus* and *H. heimii*). Proline as an essential amino acid plays a very important role in protecting the cell from osmotic stress, and it is involved in osmotic adaptation. It was discovered that proline concentration increases with the increase in salt concentration in all tested species during short-term stress, and that the ratio of proline and free amino acids increases with the increase in salt concentration in the medium. Long-term salt stress induces a decrease of proline concentration in plants grown on a medium with the high concentration of



Fig. 3. The appearance of the investigated species cultivated for three weeks in culture in vitro on a nutrient medium with the addition of various salt concentrations (photo M. Vujičić)

salt. During long-term stress, the ratio of proline to free amino acids also increases with the increase of salt concentration in the medium. In addition to the non-enzymatic components, as a response to the salt

stress enzyme activity (peroxidase, catalase, and superoxide dismutase) was monitored as well.

Phytochemical research of the bryophytes

Together with the ecophysiological studies, phytochemical studies of bryophytes have also been done. Using the UHPLC/-HESI-MS/MS methods, phenolic compounds were analyzed in *Physcomitrella patens* (Vujičić, 2016). In plants exposed to the short-term stress compared to plants exposed to the long-term salt stress, a significantly greater variety of compounds was observed. The predominantly detected compound was *p*-coumarin acid, which was detected in all treatments except in the control group. Qualitative analysis of phenolic compounds in the plants exposed to the long-term stress showed that the number of detected phenolic

compound decreases with the increase in salt concentration. In all tested species dominant phenolic compound was *p*-coumaric acid.

The chemical composition has been investigated on selected moss species: *Rodobrium ontariense*, *Atrichum undulatum*, *Hypnum andoi* (Pejin et al., 2012a, 2012c). However, there are more studies of the chemical composition of liverworts. Several liverwort taxa have been the subject of a doctoral dissertation: “Chemical composition and antimicrobial activity of various liverwort extracts of *Scapania aspera* M. & H. Bernet, *S. nemorea* (L.) Grolle, *Porella cordaeana* (Hub.) Moore and *P. arboris-vitae* (With.) Grolle)” (Bukvički, 2014). Analysis of the volatile compounds of different extracts identified 105 compounds in *Scapania aspera* (Bukvički et al., 2013), 79 in *Scapania nemorea* (Bukvički et al., 2014), 84 in *Porella cordaeana* (Bukvički et al., 2012a), and 66 in *P. arboris-vitae* (Tyagi et al., 2013). Chemical analysis of the volatile compounds of the extracts revealed the presence of large amounts of sesquiterpene and monoterpenes, and a much lower percentage of hydrocarbons, alcohols, aldehydes and ketones. GC-MS analysis of the hexane solution of methanolic extract of *Lophocolea bidentata* showed two major compounds - two sesquiterpene lactones. Diplophyllolide (eudesmanic-type of sesquiterpene lactone), an important chemotaxonomic marker for Marchantiophyta, and 5-hydroxydiplophyllolide have also been identified. Bis-bibenzyls are a significant group of chemical compounds in liverworts. Methanolic extract of *Marchantia polymorpha* subsp. *ruderalis* contains bisbibenzyl marchantin A as a




Fig. 4. Structures of seven new bisbibenzyls isolated from the methanol extract of the liverwort *Lunularia cruciata* (Novaković et al., 2019).




Fig. 5. Transmission electron micrographs of untreated and treated *Salmonella enteritidis* cells. (a) Untreated cells, (b) Methanolic *P. arboris-vitae* extract, and (c) Ethanolic *P. arboris-vitae* extract (Tyagi, 2013)

major component (Sabovljevic et al., 2017). Seven new bis-bibenzyls were isolated from the methanol extract of the liverwort *Lunularia cruciata* (Fig. 4) along with one previously known bibenzyl and five known bisbibenzyls (Novaković et al., 2019). These are acyclic bisbibenzyls, perrotetins and cyclic analogues, ricardines.

Recently, microbial transformation of molecules isolated from mosses has been the subject of the investigation. Biotransformation represents an important strategy in the structural modification of natural compounds by microorganisms.

Biological activity of bryophytes

Secondary metabolites of bryophytes have important antibacterial and antifungal potential. This has been confirmed by a series of studies on both mosses and liverworts. Among mosses, following species were studied: *Bryum argenteum* (Sabovljevic et al., 2006), *Pleurozium schreberi*, *Palustriella commutata*, *Homalothecium philippeanum*, *Anomodon attenuatus*, *Rhytidium rugosum*, *Hylocomium splendens*, *Dicranum scoparium*, *Leucobryum glaucum* (Veljić et al., 2008), *Fontinalis antipyretica*, *Hypnum cupressiforme*, *Ctenidium molluscum* (Veljić et al., 2009), *Abietinella abietina*, *Neckera crispa*, *Platyhypnidium riparoides*, *Cratoneuron filicinum*, *Campylium protensum* (Bukvički et al., 2012b) and *Rhodobryum ontariense* (Pejin et al., 2012b). In general, all extracts showed stronger antifungal than antibacterial potential. However, significantly higher antimicrobial potential has been shown by secondary metabolites in liverworts. In addition to the species from *Scapania* and *Porella* genera that were already mentioned (Bukvički, 2014), *Ptilidium pulcherrimum* was investigated as well (Veljić et al., 2010).

TEM analysis was performed to confirm antimicrobial activity and the mode of action. Transmission electron micrographs of *Salmonella enteritidis* (Fig. 5) show the effect of the extracts: (a) control

cells; (b) bacterial cell after the treatment with methanolic extract and (c) bacterial cell after the treatment of ethanol extract of the liverwort *P. arboris-vitae* (Fig. 5).

More recently, the cytotoxic potential of various molecules (like terpenes, flavonoids, sterols, coumarins, and interesting macrocyclic compounds - bibenzyls and bis-bibenzyls) present mainly in Marchantiophyta, have also been investigated. The cytotoxic activity of the newly isolated compounds, as well as the already known ricardin G was investigated. The isolated compounds showed cytotoxic activity against A549 lung cancer cells with IC₅₀ values of 5.0, 5.0 and 2.5 mM.

Pinguisane-type of compounds isolated from *Porella cordaeana* showed immunomodulatory activity (Radulović et al., 2016).

This chronological overview gives a partial review of bryological researches in Serbia, but also shows endless directions and possibilities of bryology. Bryology has large potential and perspective, considering that every new initiative in bryophyte research has given many internationally significant results. At this point, perhaps main problem in the further development of bryology is the lack of interested researchers.

Acknowledgements. This research was supported by the grant from the Ministry of Education, Science and Development of Serbia (Project No. 173029).

References

- Babić, N.** 1972: Močvarna i livadska vegetacija Koviljskog rita. *Matica srpska, Zbornik za prirodne nauke*, 41: 19–87.
- Blečić, V., Tatić, B.** 1962. Prilog poznavanju smrčevih šuma Golije planine. *Glasnik Prirodnjačkog muzeja u Beogradu*, B18: 39–47.
- Bukvički, D.** 2014: *Hemijski sastav i antimikrobna aktivnost različitih ekstrakata jetrenjača Scapania*

aspera M. & H. Bernet, *S. nemorea* (L.) Grolle, *Porella cordaeana* (Hub.) Moore i *P. arboris-vitae* (With.) Grolle". PhD thesis. Univerzitet u Beogradu, Biološki fakultet.

Bukvički, D., Gottardi, D., Veljić, M., Marin, P. D., Vannini, L., Guerzoni, M. E. 2012a: Identification of Volatile Components of Liverwort (*Porella cordaeana*) Extracts Using GC/MS-SPME and Their Antimicrobial Activity. *Molecules*, 17: 6982-6995.

Bukvički, D., Gottardi, D., Tyagi, A.K., Veljić, M., Marin, P.D., Vujisić, Lj., Guerzoni M.E. Vannini, L. 2014: *Scapania nemorea* liverwort extracts: Investigation on volatile compounds, *in vitro* antimicrobial activity and control of *Saccharomyces cerevisiae* in fruit juice, *LWT – Food Science and Technology*, 55: 452-458.

Bukvički, D., Tyagi, A., Gottardi, D., Veljić, M., Janković, S., Guerzoni, M.E., Marin, P. D. 2013: Assessment of the Chemical Composition and *in vitro* Antimicrobial Potential of Extracts of the Liverwort *Scapania aspera*. *Natural Product Communications*, 8 (9): 1313-1316.

Bukvički, D., Veljić, M., Soković, M., Grujić, S., Marin, P. D. 2012b: Antimicrobial activity of metanol extracts of *Abietinella abietina*, *Neckera crispa*, *Platyhypnidium riparoides*, *Cratoneuron filicinum* and *Campylium protensum* mosses. *Archive of Biological Sciences*, 64 (3): 911-916.

Ćosić, M., Vujičić, M., Sabovljević, M., Sabovljević, A. 2018. What do we know about salt stress in bryophytes? *Plant Biosystems*, 153 (3): 478-489.

Černjavski, P. 1929: Ein betrag zur kenntnis der geh-lzflora der kalktuffes (travertins) von Plevlje und Prijepolje. *Glasnik botaničkog zavoda i bašte Univerziteta u Beogradu*, 1(2): 205-208.

Černjavski, P. 1932: Beitrag zur postglazialen Geschichte des Blace- „Sees“ in Serbien. *Bulletin de l'Institut et du Jardin botaniques de l'Universite de Belgrade*, 2: 80-90.

Černjavski, P. 1937: Tablice za određivanje biljaka iz okoline Beograda. „Lito-Štampa“, Beograd.

Černjavski, P. 1938: Postglacijalna istorija vlasinskih šuma. Izdavačko i književno preduzeće „Geca Kon“, Beograd.

Čolić, D., Mišić, V., Popović, M. 1963: Fitocenološka analiza visokoplaninske zajednice šleske vrbe i planinske jove (*Saliceto-Alnetum viridis* ass. nova) na Staroj Planini, *Knjiga 6* (5): 1-43.

Cvetić, T., Sabovljević, M. 2005: A contribution

to the bryophyte flora of Fruška Gora (Vojvodina, Serbia). *Phytologia balcanica*, 11(1): 35-43.

Erdeši, J. 1971: *Fitocenoze šuma jugozapadnog Srema*. Doktorska disertacija. Šumsko gazdinstvo Sremska Mitrovica, Sremska Mitrovica.

Erzberger, P., Papp, B. 2007: New and noteworthy bryophyte records from Montenegro and Serbia. *Willdenowia*, 37: 339-351.

Filipović, D. 1966: Limnološka karakteristika izvorskog regiona Lisinskog potoka na Kopaoniku. *Arhiv Bioloških nauka*, 18 (3-4): 325-337.

Gajić, M. 1983: *Flora Deliblatske pešcare*. Prirodno-matematički fakultet OOUR Institut za biologiju Novi Sad; Šumsko industrijski kombinat „Pančevo“ OOUR specijalni prirodni rezervat „Deliblatski pesak“, 475 p.

Gajić, M. 1986: *Flora i vegetacija Subotičko-Horgoške pešcare*. Šumarski fakultet Beograd, Šumsko gazdinstvo Subotica, 495 p.

Gajić, M. 1988: *Flora Nacionalnog parka Tara*. Šumarski fakultet, Beograd; Nacionalni park Tara, Bajina Bašta, 288 p.

Gajić, M. 1989: *Flora i vegetacija Golije i Javora*. Šumarski fakultet, Beograd; OOUR Šumarstvo „Golija“, Ivanjica, 592 p.

Gajić, M., Karadžić, D. 1991: Flora ravnog Srema sa posebnim osvrtom na Obedsku baru. *Šumarski fakultet Beograd, Šumsko gazdinstvo Sremska Mitrovica*, 67-70.

Gajić, M., Korać, M., Obratov, D. 1991: Pregled mahovina u Srbiji, *Zbornik radova sa simpozijuma „Nedeljko Košanin i botaničke nauke“*, 401-407.

Gigov, A. 1956: Analiza polena na nekim tresavama Stare planine. *Arhiv bioloških nauka*, 1: 45-57.

Gigov, A., Nikolić, V. 1954: Rezultati analize polena na tresavama planine Ostrozub. *Arhiv bioloških nauka*, 6(1-2): 99-116.

Grdović, S. 2005: Mahovine šireg područja Beograda i njihov bioindikatorski značaj, *Zadužbina Andrejević*, Beograd, 130 str.

Grdović, S., Blaženčić, Ž. 2001: Prilog poznavanju mahovina rezervata „Zasavica“, *Monografija „Zasavica“*, 35-41.

Grdović, S., Stavretović, N. 2004: Mahovine kao korov u travnim površinama. *Acta Herbologica*, 13(1): 235-242.

Grdović, S., Stevanović, V. 2006: The moss flora in the central urban area of Belgrade. *Archives of Biological Sciences*, 58(1): 55-59.

- Grebenščikov, O.** 1943: Prilog poznavanju vegetacije planine Koprivnik kod Peći. Srpska kraljevska akademija, Posebna izdanja 136, Prirodnjački i matematički spisi 35, *Ohridski zbornik*, 2: 241–267 (1–29).
- Grebenščikov, O.** 1949: Pregled radova na polju proučavanja mahovina u Srbiji, *Glasnik Prirodnjačkog muzeja srpske zemlje*, B, 1-2: 315–321.
- Grebenščikov, O.** 1950: O vegetaciji centralnog dijela Stare planine. *Zbornika radova Instituta za ekologiju i biogeografiju*, 1: 1–36.
- Grgić, P.** 1983: Prilog poznavanju mahovina u ekosistemima sa pančičevom omorikom. *Godišnjak Biološkog instituta*, 36: 69–72.
- Grisebach, A.** 1843: *Spicilegium Florae rumelicae et bitynicae exhibens synopsis plantarum, Brunsvigae.*
- Guelmino, J.** 1970: Prva nalazišta mahovine *Funaria hungarica* Boros u Jugoslaviji, *Zbornik za prirodne nauke, Matica Srpska*, 37: 176–178.
- Guelmino, J.** 1972: Nova nalazišta dve malo poznate mahovine kod nas (*Funaria hungarica* Boros i *Tortula Velenovsky* Schiffner). *Zbornik za prirodne nauke, Matica srpska*, 42(1): 106–109.
- Hodgetts, N., Cáliz, M., Englefield, E., Fettes, N., García Criado, M. et al.** 2019: A miniature world in decline: European Red List of Mosses, Liverworts and Hornworts. Brussels, Belgium: IUCN.
- Ilić, M.** 2019: *Diverzitet, distribucija, diferencijacija mikrostaništa i struktura zajednica mahovina Fruške gore*. Phd thesis. Univerzitet u Novom Sadu, Prirodno-matematički fakultet, Departman za biologiju.
- Ilić, M., Vukov, D., Rućando, M., Ćuk, M., Igić, R.** 2015: Contribution to the bryophyte flora in beech forests of Vidlič Mountain (Serbia). *Zbornik Matice Srpske za Prirodne Nauke*, 174: 67–27.
- Janković, M., Janković, M. M.** 1962: Ekološki uslovi vegetacije u vrelu Mlave kod Žagubice, sa posebnim osvrtom na biljku *Callitriche verna*. *Arhiv bioloških nauka*, 14 (3-4), 157-168.
- Janković, M., Mišić, V.** 1980: *Šumska vegetacija i fitocenoze Fruške gore*. Monografije Fruške gore. Matica srpska, Novi Sad, 191 p.
- Janković, M., Stevanović, V.** 1981: Prilog poznavanju fitocenoza sa srpskom ramondijom (*Ramonda serbica* Panč.) u klisurama severnih ogranaka Šarplanine. *Ekologija*, 16(1): 1–34.
- Jovanović B., Vukićević, E., Avdalović, V.** 1983: Neke planinske zajednice crne i bele jove u okolini Sjenice. *Zaštita prirode*, 36: 49–68.
- Jovanović, B.** 1980: Šumske fitocenoze i staništa Suve planine. *Glasnik Šumarskog fakulteta*, serija A - Šumarstvo, posebno izdanje 55.
- Jurišić, J. Ž.** 1900: Prilog poznavanju mahovina u Srbiji. *Spomenik Srpske Akademije Nauka*, 45: 47–60.
- Katić, D.** 1900: Drugi prilog flori okoline Kragujevca. *Izveštaj Gimnazije kneza Miloša u Kragujevcu za 1899/1900. godinu*, 43–45.
- Katić, D.** 1903: Treći floristički prilog (iz Kragujevačke okoline). *Izveštaj Gimnazije kneza Miloša Velikog u Kragujevcu za 1902/1903*, 3–7.
- Katić, D.** 1906: Beitrag zur Mossflora von Serbien. *Hedwigia*, 45: 92–99.
- Katić, D.** 1907a: Nekoliko mahovinskih prinova flori Srbije. *Prosvetni glasnik*, 12: 884–888.
- Katić, D.** 1907b: Sitniji prilozi flori Srbije. *Nastavnik, list profesorskoga društva*, 18: 184–191.
- Katić, D.** 1909: Priložak mahovinskoj flori Srbije. *Nastavnik*, 20 (7-8): 285–286.
- Knapp, R.** 1944: Vegetationsstudien in Serbien. Halle.
- Košanin, N.** 1909a: Das Vorkommen von *Polytrichum alpinum* L. auf einem Hochmoor in Serbien. *Hedwigia*, 48: 205–206.
- Košanin, N.** 1909b: Moose aus dem Gebiete des Golia-Gebirges in Südwest-Serbien. *Hedwigia*, 48: 207–209.
- Košanin, N.** 1910: Elementi Vlasinske flore (Algae, Bryophyta, Pteridophyta et Phanerogamae). *Prosvetni Glasnik Beograd*, 31 (8-9): 684–700.
- Lakušić, D.** 1996: Pregled flore Kopaonika (JZ Srbija, Jugoslavija). *Ekologija*, 31(2): 1–35.
- Lazarević, P., Pantović, J., Szurdoki, E., Papp, B., Sabovljević, M.** 2016: Distribution, ecology and threat status evaluation of *Sphagnum* species in Serbia. *Wulfenia*, 23: 37–51.
- Marinković, P., Gajić, M.** 1956: O jednom nalazištu sfagnumske tresave u Srbiji. *Šumarstvo*, 4–5: 258–262.
- Marka, J., Papp, B., Erzberger, P., Colacino, C., Sabovljević, M.** 2012: Towards a red list of the Albanian bryophytes. *Studia Botanica Hungarica*, 43: 13-25.
- Martinčić, A.** 1980: Prispjevek k poznavanju mahovne flore Jugoslavije II - Šar planina. *Biološki vestnik*, 28 (2): 87–102.

- Matouschek, F.** 1899: Beitrag zur Mooskenntnis von Südserbien. *Verhandlungen der Zoologisch-Botanischen Gesellschaft in Wien*, 49: 386–390.
- Matouschek, F.** 1901: Bryologisch floristisches aus Serbien. *Allgemeine botanische Zeitschrift für Systematik, Floristik, Pflanzengeographie etc.*, 7 (2): 21–22.
- Mišić, V.** 1982: Reliktne polidominantne šumske zajednice Srbije. *Matica Srpska*, 178 p.
- Mišić, V., Panić, I.** 1989: Šumska vegetacija doline Studenice. *Zaštita prirode*, 41-42: 33–52.
- Mišić, V., Popović, M.** 1960: Fitocenološka analiza smrčevih šuma Kopaonika. *Zbornik radova Biološkog instituta N. R. Srbije*, 3(5): 1–26.
- Novaković, M., Bukvički, D., Andjelković, B., Ilić Tomić, T., Veljić, M., Tešević, V., Asakawa, Y.** 2019: Cytotoxic Activity of Riccardin and Perrottetin Derivatives from the Liverwort *Lunularia cruciata*. *Journal of Natural Products*, DOI:10.1021/acs.jnatprod.8b00390, in press.
- Pančić, J.** 1859: Die Flora der Serpentinberge in Mittel-Serbien. *Verhandlungen der Zoologisch-Botanischen Gesellschaft in Wien*, 9: 139–150.
- Pančić, J.** 1863: Živi pesak i bilje što raste na njemu. *Glasnik društva srbske slovesnosti*, 16: 197–233.
- Pantović, J.** 2018: *Biogeografska i ekološka studija flore briofita Srbije*. Phd thesis. Univerzitet u Beogradu, Biološki fakultet.
- Pantović, J., Grdović, S., Sabovljević, A., Sabovljević, M.** 2014: New and interesting bryophyte records for the flora of Serbia. *Archives of Biological Sciences*, 66 (2): 701–704.
- Pantović, J., Sabovljević, M.** 2013: Contribution to the bryophyte flora of Mt Boranja (West Serbia). *Phytologia Balcanica*, 19 (1): 23–27.
- Pantović, J., Sabovljević, M.** 2017: Bryophytes of Kosovo. *Phytotaxa*, 306 (2): 101–123.
- Pantović, J., Sabovljević, M. S.** 2017: Overview of bryophyte flora research in Serbia with presentation of the Serbian BRYO database. *Botanica Serbica*, 41(2): 153-162.
- Papp, B., Alegro, A., Erzberger, P., Szurdoki, E., Šegota, V., Sabovljević, M.** 2016a: Bryophytes of saline areas in the Pannonian region of Serbia and Croatia. *Studia Botanica Hungarica*, 47 (1): 141–150.
- Papp, B., Erzberger, P.** 2005: The bryophyte flora of Golija - Studenica biosphere reserve and some adjacent sites (SW Serbia - Montenegro). *Studia Botanica Hungarica*, 36: 101–116.
- Papp, B., Erzberger, P.** 2007a: Contributions to the bryophyte flora od western Stara planina Mts (E Serbia). *Studia Botanica Hungarica*, 38: 95–123
- Papp, B., Erzberger, P.** 2007b: New and noteworthy bryophyte records from Montenegro and Serbia. *Willendowia*, 37: 339–351.
- Papp, B., Erzberger, P.** 2009: Contributions to the bryophyte flora of southeastern Serbia: Suva planina Mts and its surroundings. *Studia Botanica Hungarica*, 40: 125–142.
- Papp, B., Erzberger, P., Sabovljević, M.** 2004: Contributions to the bryophyte flora of Kopaonik Mts (Serbia, Serbia-Montenegro). *Studia Botanica Hungarica*, 35: 67–79.
- Papp, B., Erzberger, P., Sabovljević, M.** 2006: Contribution to the bryophyte flora of the Djerdap National Park (E Serbia). *Studia Botanica Hungarica*, 37: 131–144.
- Papp, B., Natcheva, R., Erzberger, P., Sabovljević, M.** 2012a: *Didymodon sicculus*, new to Bulgaria and Serbia and notes on its ecology. *Nova Hedwigia*, 95 (1-2): 221–226.
- Papp, B., Pantović, J., Sabovljević, M., Szurdoki, E.** 2014a: *Myurella sibirica*, a moss species new to Montenegro and Serbia: its range extension towards south-eastern Europe. *Cryptogamie, Bryologie*. 35(3): 1–6.
- Papp, B., Pantović, J., Szurdoki, E., Sabovljević, M.** 2014b: Interesting and new species for the bryophyte flora of Serbia. *Herzogia*, 27(1): 221–225.
- Papp, B., Sabovljević, M.** 2001: Contribution to the knowledge of the bryoflora of the region of Petnica (W Serbia, Yugoslavia). *Studia Botanica Hungarica*, 32: 107–120
- Papp, B., Sabovljević, M.** 2002: The bryophyte flora of Tara National Park (W Serbia, Yugoslavia). *Studia Botanica Hungarica*, 33: 25–39.
- Papp, B., Sabovljević, M.** 2010: Contribution to the Bryophyte flora of the Vršacke Planine Mts., Serbia. *Botanica serbica*, 34(2): 107–110.
- Papp, B., Szurdoki, E., Pantović, J., Sabovljević, M.** 2013: *Physcomitrium euryostomum* and *Pohlia prolifera*, new mosses in the bryophyte flora of Serbia. *Archives of Biological Sciences*, 65 (2): 703–706.
- Papp, B., Szurdoki, E., Pantović, J., Sabovljević, M.** 2014c: Contributions to the bryophyte flora of the Pešter plateau, SW Serbia. *Studia Botanica Hungarica*, 45: 33–47.

- Papp, B., Szurdoki, E., Pantović, J., Sabovljević, M.** 2016b: An insight into the bryophyte flora of the Ibar gorge and its surroundings (Central and SW Serbia). *Acta Botanica Hungarica*, 58: 411–423.
- Papp, B., Szurdoki, E., Pantović, J., Sabovljević, M.** 2016c: New records of Mediterranean-Atlantic mosses in the flora of Serbia. *Herzogia*, 29:185–189.
- Papp, B., Szurdoki, E., Sabovljević, M.** 2012b: Bryophyte flora of lake Vlasina and its surroundings (SE Serbia). *Studia botanica hungarica*, 43: 27–45.
- Pavić, S., Sabovljević, M., Stevanović, V.** 1998: Diversity and threat status of the Yugoslav bryoflora. *Lindbergia*, 23: 38–44.
- Pavletić, Z.** 1955: *Prodromus flore briofita Jugoslavije*. Jugoslavenska Akademija znanosti i umetnosti, 578 p.
- Pavletić, Z.** 1956: Prilog poznavanju briofitskog endemizma u flori Jugoslavije. *Acta Musei Macedonici Scientiarum naturalium*, 4: 2–33.
- Pavletić, Z.** 1968: Flora mahovina Jugoslavije. Institut za botaniku Sveučilišta u Zagrebu, 431 p.
- Pavlović, Z.** 1951: Vegetacija planine Zlatibor. *Srpska akademija nauka, Zbornik radova II, Institut za ekologiju i biogeografiju*, 2: 115–182.
- Pavlović, Z.** 1964: Borove šume na serpentinama u Srbiji. *Glasnik Prirodnačkog muzeja u Beogradu*, B 19: 25–65.
- Pejin, B., Sabovljević, A., Soković, M., Glamočlija, J., Ćirić, A., Vujičić, M., Sabovljević, M.** 2012a: Antimicrobial activity of *Rhodobryum ontariense*. *Hemijska Industrija*, 66 (3): 381–384.
- Pejin, B., Bianco, A., Newmaster, S., Sabovljević, M., Vujisić, Lj., Tešević, V., Vajs, V., De Rosa, S.** 2012b: Fatty acids of *Rhodobryum ontariense* (Bryaceae). *Natural Product Research*, 26 (8): 696–702.
- Pejin, B., Vujisić, Lj., Sabovljević, M., Tešević, V., Vajs, V.** 2012c: Fatty acid chemistry of *Atrichum undulatum* and *Hypnum andoi*. *Hemijska Industrija*, 66 (2): 207–209.
- Petković, B., Tatić, B., Ilijin–Jug, M.** 1988: Dve nove zajednice srpske Ramondije (*Ramonda serbica* Panč.) u gornjem toku sliva reke Ibra. *Glasnik Instituta za botaniku i botaničke bašte Univerziteta u Beogradu*, 22: 107–116.
- Petković, B., Tatić, B., Marin, D. P., Dimić, J.** 1990: Prilog poznavanju zajednica sa *Edraianthus jugoslavicus* Lakušić sa Mokre Gore (jugozapadna Srbija). *Bilten Društva ekologa Bosne i Hercegovine*, B5: 131–135.
- Petković, B., Tatić, B., Marin, P., Ilijin–Jug, M.** 1991: *Musco-Seslerio-Edraianthetum jugoslavici*, ass. nov. nova zajednica klisura i kanjona tutinskog područja. *Drugi simpozijum o flori jugoistočne Srbije i mogućnostima njenog racionalnog korišćenja*, Univerzitet u Nišu, Tehnološki fakultet u Leskovcu, Leskovac. Zbornik radova 1(7-8/89-90): 133–138.
- Petković, B., Tatić, B., Marin, P., Veljić, M.** 2000: Vegetacija spomenika prirode „Đavolja Varoš”. 6. *Simpozijum o flori jugoistočne Srbije i susednih područja, Sokobanja*, Zbornik radova: 323–347.
- Pichler, A.** 1931: Prilog poznavanju mahova tresetara Jugoslavije. *Acta botanica croatica*, 6: 47–55.
- Pichler, A.** 1939: Die Lebermossflora auf morschen Baumstämmen, vermoderndem Holz und auf faulenden Baumstümpfen der Wälder Jugoslaviens. *Glasnik Skopskog naučnog društva*, 20: 141–153.
- Pichler, A.** 1940: Hepatike na kori drveća i grmlja naših šuma. *Šumski list*, 7: 334–354.
- Pócs, T., Sabovljević, M., Puche, F., Segarra Moragues, J. G., Gimeno, C., Kürschner, H.** 2004: *Crossidium laxefilamentosum* Frey & Kürschner (Bryopsida: Pottiaceae), new to Europe and to North Africa. *Journal of Bryology*, 26: 113–124.
- Podpera, J.** 1922: Ad Bryophytorum Haemi peninsulae cognitionem additamentum. *Acta Botanica Bohemica*, 1: 1–18.
- Popović, M.** 1966: Prilog poznavanju mahovina u rezervatima i zaštićenim područjima u Srbiji. *Zaštita prirode*, 33: 219–228.
- Radulović, N.S., Filipović, S.I., Zlatković, D.B., Đorđević, M.R., Stojanović, N.M., Randjelović, P.J., Mitić, K.V., Jevtpivčić-Stoimenov, T.M., Randelović, V.N.** 2016: Immunomodulatory pinguicane-type sesquiterpenes from the liverwort *Porella cordaeana* (Porellaceae): the “new old” furanopinguicane and its oxidation product exert mutually different effects on rat splenocytes. *RSC Advances*, 6(48), 41847–41860
- Rajčić, M., Nerić, V., Vujičić, M., Sabovljević, A., Sabovljević, M.** 2015: The effect of short- and long-termed salt stress on the *Atrichum undulatum* (Hedw.) P. Beauv. Sixth Balkan Botanical Congress, 14–18 September 2015, Rijeka, Croatia. Book of Abstracts: 102.
- Rajevski, L.** 1951: Borove šume u predelima od Mokre Gore do reke Uvac. *Srpska akademija nauka, Zbornik radova 11, Institut za ekologiju i biogeografiju*, 2: 183–192.
- Randelović, V.** 1994: *Geobotanička studija vlasinske*

tresave. Biološki fakultet, Univerziteta u Beogradu.

Randelović, V., Zlatković, B. 2010: *Flora i vegetacija Vlasinske visoravni*. Prirodno-matematički fakultet, Univerzitet u Nišu.

Rowntree, J. K., Pressel, S., Ramsay, M.M., Sabovljević, A., Sabovljević, M. 2011: In vitro conservation of European bryophytes. *In Vitro Cell Dev Biol – Plant*, 47: 55–64.

Rudski, I. 1949a: Ekскурzija na Žljeb: Moku planinu. *Prirodnjački muzej srpske zemlje*, 23. Beograd.

Rudski, I. 1949b: Tipovi lišćarskih šuma jugoistočnog dela Šumadije. *Prirodnjački muzej srpske zemlje*, 25: 1–67.

Sabovljević, A., Soković, M., Sabovljević, M., Grubišić, D. 2006: Antimicrobial activity of *Bryum argenteum*. *Fitoterapia*, 77 (2): 144–145.

Sabovljević, A., Sabovljević, M., Jocković, N. 2009: In vitro culture and secondary metabolite isolation in bryophytes. In: Mohan JS, Saxena PK, editors. Protocols for In vitro cultures and secondary metabolite analysis of aromatic and medicinal plants. Methods in Molecular Biology. New York, NY: Humana Press, 117–128.

Sabovljević, M. 1998: The rare Bryophytes of Šara Mountain, Yugoslavia, *Planta Europa Proceedings of the Second European Conference on the Conservation of Wild Plants*, 2: 159–161.

Sabovljević, M. 1999: *Anastrophyllum minutum* (Schreb.) Schust., new to Serbia (FR Yugoslavia) and its distribution in the Balkans. *Phytologia Balcanica*, 5(2-3): 93–96.

Sabovljević, M. 2000: Checklist of hepatics of the Federal Republic of Yugoslavia. *Lindbergia* 25: 37–42.

Sabovljević, M. 2003a: Bryophyte Flora of South Banat (Vojvodina, Yugoslavia). *Cryptogamie, Bryologie*, 24 (3): 241–252.

Sabovljević, M. 2003b: Données sur la présence et la chorologie des taxons du genre *Schistidium* (Grimmiaceae) dans la République Fédérale de Yougoslavie (Serbie et Monténégro). *Bocconeia*, 16 (2): 991–999.

Sabovljević, M. 2006: Contribution to the bryophyte flora of the Djerdap National Park (E Serbia). *Phytologia Balcanica*, 12 (1): 51–54.

Sabovljević, M. 2015: *Flora Briofita Srbije I. Tresetnice (Sphagnophyta)*. Srpska Akademija Nauka i Umetnosti, 110 p.

Sabovljević, M., Cvetić, T. 2001: *Rhodobryum ontariense* (Kindb.) Kindb. new to Yugoslavia and some notes on the genus *Rhodobryum* (Schimp.) Limpr. in the Federal Republic of Yugoslavia. *Ekologija*, 36 (2): 145–153.

Sabovljević, M., Cvetić, T. 2003: Bryophyte flora of Avala Mt. (C. Serbia, Yugoslavia). *Lindbergia*, 28: 90–96.

Sabovljević, M., Cvetić, T., Stevanović, V. 2004: Bryophyte red list of Serbia and Montenegro. *Biodiversity and conservation*, 13 (9): 1781–1789.

Sabovljević, M., Grdović, S. 2009: Bryophyte Diversity Within Urban Areas: Case Study of the City of Belgrade (Serbia). *International Journal of Botany*, 5 (1): 85–92.

Sabovljević, M., Marka, J. 2009: The biological evidence of climate changes: a case study of liverwort *Lunularia cruciata* (L.) Dum. ex Lindb. in Serbia. *Botanica serbica*, 33(2): 185–187.

Sabovljević, M., Sérgio, C. 2002: Contribution to the bryoflora of Serbia: a bryophyte collection from 1996. *Portugaliae Acta Bioogica*, 20: 65–74.

Sabovljević, M., Stevanović, V. 1999: Moss conspectus of Federal Republic of Yugoslavia. *Flora Mediterranea*, 9: 65–95.

Sabovljević, M., Stevanović, V. 2000: *Sphagnum denticulatum* Brid., novedad para la brioflora de Serbia (Yugoslavia). *Botanica Complutensis*, 24: 61–63.

Sabovljević, M., Stevanović, V. 2006: Contribution to knowledge of the bryophyte flora of Bačka (Vojvodina, Serbia). *Archives of Biological Sciences*, 58 (2): 135–138.

Sabovljević, M., Papp, B., Sabovljević, A., Vujičić, M., Szurdoki, E., Segarra-Moragues, J. G. 2012a: In vitro micropropagation of rare and endangered moss *Enthostodon hungaricus* (Funariaceae). *Bioscience Journal*, 28: 632–640.

Sabovljević, M. S., Segarra-Moragues, J. G., Puche, F., Vujičić, M., Cogoni, A., Sabovljević, A. 2016: Eco-physiological and biotechnological approach to conservation of the world-wide rare and endangered aquatic liverwort *Riella helicophylla* (Bory et Mont.) Mont. *Acta Botanica Croatica*, 75 (2): 194–198.

Sabovljević, M. S., Nikolić, N., Vujičić, M., Sinžar-Sekulić, J., Pantović, J., Papp, B., Sabovljević, A. 2018: Ecology, distribution, propagation in vitro, ex situ conservation and native population strengthening of rare and threatened halophyte moss *Enthostodon hungaricus* in Serbia. *Wulfenia*, 25: 117–130.

- Sabovljević, M., Vujičić, M., Pantović, J., Sabovljević, A. 2014: Bryophyte conservation biology: *In vitro* approach to the *ex situ* conservation of bryophytes from Europe. *Plant Biosystems*, 148 (4), 857-868.
- Sabovljević, M., Vujičić, M., Šinžar-Sekulić, J., Segarra-Moragues, J. G., Bapp, B., Skorić, M., Dragačević, L., Sabovljević, A. 2012b: Reviving, *in vitro* differentiation, development and micropropagation of the rare and endangered moss *Bruchia vogesiaca* (Bruchiaceae). *HortScience*, 47 (9): 1347-1350.
- Sabovljević, M. S., Vujičić, M., Wang, X., Garraffo, M., Bewley, C. A., Sabovljević, A. 2017: Production of the macrocyclic bis-bibenzyls in axenically farmed and wild liverwort *Marchantia polymorpha* L. subsp. *ruderalis* Bischl. et Boisselier. *Plant Biosystems*, 151 (3): 414-418.
- Schiffner, V. 1897: Musci Bornmüllerian, ein Beitrag zum kryptogramenflora des Orients. *Osterr Bot Zeitschr*, 125 p.
- Simić, M. 1892: Nekoliko srpskih mahovina, *Nastavnik, list profesorskoga društva*, III knjiga, 447-450.
- Simić, M. 1897: Nekoliko kriptogamskih biljaka u okolini vranjskoj. *Izveštaj vranjske gimnazije za školsku 1896/97. godinu*, 3-8.
- Simić, M. 1898: Kriptogamne biljke u okolini vranjskoj. *Izveštaj vranjske gimnazije za školsku 1897/98. godinu*, 13-16.
- Simić, M. 1900: Prilog flori mahovina u Srbiji, *Spomenik Srpske Akademije Nauka*, 35: 5-46.
- Slavnić, Ž. 1956: Vodena i barska vegetacija Vojvodine. *Zbornik Matice srpske, serija prirodnih nauka*, 10: 5-72.
- Soška, T. 1949: Pregled mahovina i lišajeva u okolini Beograda. *Glasnik 10. prirodnjackog muzeja srpske zemlje*, 1-2: 93-112.
- Stevanović, V., Pavić, S., Stevanović, B. 1995: Diverzitet flore mahovina (*Bryophyta*) Jugoslavije sa pregledom vrsta od međunarodnog značaja. – In: Stevanović, V., Vasić, V.(eds): *Biodiverzitet Jugoslavije sa pregledom vrsta od međunarodnog značaja*. Biološki fakultet i Ecolibri, Beograd.
- Tešić, Ž., Gigov, A., Bogdanović, M., Milić, Č. 1979: Tresave Srbije. *Zbornik radova Geografskog instituta Jovan Cvijić*, 31: 19-64.
- Tyagi, A.K., Bukvički, D., Gottardi, D., Veljić, M., Guerzoni, M.E., Vannini, L., Malik, A., Marin, P. D. 2013: Antimicrobial Potential and Chemical Characterization of Serbian Liverwort (*Porella arboris-vitae*): SEM and TEM Observations. *Evidence-Based Complementary and Alternative Medicine*, Article ID 382927,
- Veljić, M. 2013: Flora mahovina planine Kopaonik, Srbija. *Zaštita prirode*, 53 (1-2): 73-91.
- Veljić, M., Bukvički, D., Marin, P. D. 2016a: *Sphagnum fimbriatum*, a species new for the flora of Serbia. *Botanica Serbica*, 40 (1): 101-103.
- Veljić, M., Ljubić, B., Marin, P. D., Marin, M. 2008: Bryophyte flora on the northern slopes of Zlatar mountain (Southwest Serbia). *Archives of Biological Sciences*, 60 (1): 145-150.
- Veljić, M., Marin, P. D. 1997: New moss taxa for the flora of Serbia. *Flora Mediterranea*, 7: 133-138.
- Veljić, M., Marin, P. D., Boža, P., Petković, B. 1996: Flora mahovina vrela Grze i Crnog Timoka (Srbija). *Glasnik Instituta za botaniku i Botaničke bašte Univerziteta u Beogradu*, 30: 97-106.
- Veljić, M., Marin, P. D., Boža, P., Petković, B. 2001a: Bryoflora of some well-springs of the Dinaric Alps and Carpathian karst in Serbia. *Bocconea*, 13: 343-351.
- Veljić, M., Marin, P. D., Petković, B., Ljubić, B. 2001b: New species for the bryophyte flora of Yugoslavia. *Cryptogamie, Bryologie*, 22 (4): 275-277.
- Veljić, M., Marin, P. D., Petković, B., Ljubić, B. 2004: New records for the bryophyte flora of Serbia. *Archives of Biological Sciences*, 56 (1-2): 11-12.
- Veljić, M., Marin, P., Lakušić, D., Ljubić, B. 2006: Bryophyte flora of the Uvac river george (southwest Serbia). *Archives of Biological Sciences*, 58 (3): 187-194.
- Veljić, M., Tarbuk, M., Marin, P. D., Ćirić, A., Soković, M., Marin, M. 2008: Antimicrobial activity of Methanol Extracts of Some Genuine Mosses from Serbia, *Pharmaceutical Biology*, 46 (12), 871-875.
- Veljić, M., Ćirić, A., Soković, M., Janačković, P., Marin, P. D. 2010: Antibacterial and antifungal activity of the liverwort (*Ptilidium pulcherrimum*) metanol extract. *Archives of Biological Sciences*, 62 (2): 381-395.
- Veljić, M., Đurić, A., Soković, M., Ćirić, A., Glamočlija, J., Marin P. D. 2009: Antimicrobial activity of Methanol Extracts of *Fontinalis antipyretica*, *Hypnum cupressiforme* and *Ctenidium molluscum*. *Archives of Biological Sciences*, 61 (2): 225-229.
- Veljić, M., Rajčević, N., Bukvički, D. 2016b: A

revision of the moss collection of the University of Belgrade Herbarium (BEOU) from the Ostrožub Mountain in Serbia. *Biologica Nyssana*, 7 (1): 11-17.

Vujičić, M. 2016: *Biohemijski i ekofiziološki odgovori odabranih vrsta mahovina na kontrolisani stres izazvan solima u akseničnim uslovima*. Phd thesis. Univerzitet u Beogradu, Biološki fakultet.

Vujičić, M., Sabovljević, A., Milošević, S., Segarra-Moragues, J. G., Sabovljević, M. 2016: Effects of abscisic acid (ABA) on development of

selected bryophyte species. *Plant Biosystems*, 150: 1023-1029.

Vujičić, M., Sabovljević, A., Šinžar-Sekulić, J., Škorić, M., Sabovljević, M. 2012: *In vitro* development of the rare and endangered moss *Molendoa hornschurchiana* (Hook.) Lindb. ex Limpr. (Pottiaceae, Bryophyta). *HortScience*, 47: 84–87.

Wettstein, R. 1890: Ueber das Vorkommen von *Trochobryum Carniolicum* in Südserbien. *Oesterreichische botanische Zeitschrift*, 40: 170–171.