

**CONTROL OF AFLATOXIGENIC *Aspergillus flavus* IN PEANUTS USING  
NONAFLATOXIGENIC *A. flavus*, *A. niger* and *Trichoderma harzianum***

OKKY SETYAWATI DHARMAPUTRA

SEAMEO BIOTROP, P.O. Box 116, Bogor, Indonesia and  
Faculty of Mathematics and Natural Sciences, Bogor Agricultural University, Bogor,  
Indonesia

ASMARINA S.R. PUTRI, INA RETNOWATI and SANTIAMBARWATI

SEAMEO BIOTROP, P.O. Box 116, Bogor, Indonesia

**ABSTRACT**

The effects of nontoxigenic *Aspergillus flavus*, *A. niger* and *Trichoderma harzianum* inoculated into planting media on toxigenic *A. flavus* infection and its aflatoxin production in peanut kernels at harvest were investigated together with (1) the moisture content of planting media before peanut planting, at the time of inflorescence, and at harvest, (2) the population of aflatoxigenic and nonaflatoxigenic *A. flavus*, *A. niger* and *T. harzianum* in peanut planting media before peanut planting, at the time of inflorescence, and at harvest, (3) the moisture content of peanut kernels at harvest, and (4) toxigenic *A. flavus* invasion in peanut plant parts (roots, stems, petioles, leaves and flowers) at the time of inflorescence.

The fungal isolates were inoculated into planting media at the same time with the planting of peanut seeds. Peanut plants were grown under glasshouse conditions. Treated planting media were inoculated with the combined use of (1) toxigenic and nontoxigenic *A. flavus*, (2) toxigenic *A. flavus* and *A. niger*, and (3) toxigenic *A. flavus* and *T. harzianum*. Planting media inoculated only with each fungal isolate and uninoculated planting media were used as controls. Two watering treatments of peanut plants were carried out, i.e. watering until harvest and not watering for 15 days before harvest. The populations of the fungal isolates in the planting media and peanut kernels were determined using dilution method followed by pour plate method; the percentages of toxigenic *A. flavus* and test fungal colonizations in peanut plant parts were determined using plating method; the moisture content of planting media and peanut kernels were determined using oven method; the aflatoxin content of peanut kernels was determined using Thin Layer Chromatography method.

The results indicated that at the time of harvest the decrease in moisture contents of planting media not watered for 15 days before harvest was higher than those watered until harvest. The lowest population of toxigenic *A. flavus* was in planting media inoculated with the combined use of toxigenic and nontoxigenic *A. flavus* at the time of inflorescence and at the time of harvest. Toxigenic *A. flavus* could invade the roots, stems and flowers of peanut plants. The lowest percentage of invasion was on the plant parts which planting media were inoculated with the combined use of toxigenic and nontoxigenic *A. flavus*. The moisture content of peanut kernels originated from watered plants until harvest were higher than those not watered for 15 days before harvest. The population of toxigenic *A. flavus* in peanut kernels derived from the plants whose planting media were inoculated with the combined use of toxigenic *A. flavus* and each test fungi, was lower than those inoculated only with toxigenic *A. flavus*. It indicated that the test fungi inoculated into planting media could inhibit toxigenic *A. flavus* infection in peanut kernels. Aflatoxin was only detected in peanut kernels originated from one plant whose planting medium was inoculated only with toxigenic *A. flavus* and the plant was watered until the time of harvest. Toxigenic *A. flavus* infection and aflatoxin production were not influenced by planting media which were not watered for 15 days before harvest.

**Keywords:** Biocontrol / Aflatoxigenic / Nonaflatoxigenic / *Aspergillus flavus* / *Aspergillus niger* / *Trichoderma harzianum* / Peanuts

## INTRODUCTION

Aflatoxins are potential hepatotoxic and carcinogenic metabolites produced by the fungi *Aspergillus flavus*, *A. parasiticus*, and *A. nomius*. Contamination of aflatoxins occurs when peanut kernels become infected by the three fungal species under drought stress before harvest, during the drying phase in the field, or under unsuitable storage conditions.

Pitt and Hocking (1996) reported that 45, 33 and 22% of 215 peanut samples collected from farm storage, middlemen and retailers in Bogor and Yogyakarta contained aflatoxins of more than 50, 300 and 1000 ppb, respectively. Levels of 1000 ppb of aflatoxins will cause liver damage in man and animals. Lower levels of aflatoxins consumed from peanut products can cause liver cancer and premature death in humans, as well as reducing productivity of livestock. Based on the report of the 23<sup>rd</sup> Session of the Joint FAO/WHO Food Standards Programme, held in Rome, Italy, 28 June - 3 July 1999, Codex Alimentarius Commissions adopted 15 ppb as the maximum level of total aflatoxins to be contained in peanuts intended for further processing.

The entry of aflatoxin in the peanut plant could be from the roots, flowers, leaves damaged by insects, and from air or dust. The most important point of entry to the developing peanut, however, is directly from the soil surrounding it. So, the point of application of any biocontrol measure such as competitive exclusion should be in the soil of peanut - growing fields.

There are only few reports on the reduction of aflatoxins by competitive fungi. Ehrlich *et al.* (1985) reported that by growing *A. parasiticus* in mixed culture with *Penicillium oxalicum*, toxin productions of both species were reduced. To assess this effect under more practical conditions, Wicklow *et al.* (1988) inoculated corn kernels with toxigenic *A. flavus* at preharvest, with or without a range of other fungi present. They found that much lower levels of preharvest aflatoxin were produced in the presence of some other fungal species. Dorner *et al.* (1999) found that aflatoxin contamination of corn was reduced when the soil of corn plots were inoculated with nonaflatoxigenic strain of *A. flavus*. Dharmaputra *et al.* (2001) reported that non-aflatoxigenic *A. flavus* BIO 2127, *A. niger* BIO 2129 and *Trichoderma harzianum* BIO 19130 were antagonistic to aflatoxigenic *A. flavus* BIO 2128 *in vitro*. *Aspergillus niger* was the most promising fungal antagonist, because it caused the highest percent inhibition of aflatoxin production of *A. flavus* BIO 2128.

The objectives of this study were to investigate the effects of nontoxigenic *A. flavus*, *A. niger* and *T. harzianum* inoculated into the peanut planting media on toxigenic *A. flavus* infection and its aflatoxin production in peanut kernels at harvest. Investigations were also carried out on (1) the moisture contents of planting media before peanut planting, at the time of inflorescence, and at harvest, (2) the population of aflatoxigenic and nonaflatoxigenic *A. flavus*, *A. niger* and *T. harzianum* in peanut planting media before peanut planting, at the time of inflorescence and at harvest, (3) the moisture content of peanut kernels at the time of harvest, and (4) toxigenic *A. flavus* invasion in peanut plant parts (roots, stems, petioles, leaves and flowers) at the time of inflorescence.

## MATERIALS AND METHODS

### Planting media, peanut variety, isolates of toxigenic *A. flavus* and test fungi

A mixture of soil (latosol type), sand and casting (2:1:1) was used as planting medium, while peanut variety Komodo was obtained from the Research Institute for Legumes and Tuber Crops (RILET) in Malang. Four fungal isolates were used: toxigenic *A. flavus* BIO 2128, nontoxigenic *A. flavus* BIO 2127, *A. niger* BIO 2129, and *T. harzianum* BIO 19130. The latter three fungal isolates were used as test fungi.

### Preparation of *A. flavus*, *A. niger* and *T. harzianum* inocula

Mixtures of ground rice bran and tap water (4:3) were placed in transparent polyethylene bags (125 g/bag). They were then sterilized in an autoclave for 1 h, and incubated at room temperature for one night. For the time being, each fungal isolate was grown on Potato Dextrose Agar in Petri dishes (diam 9 cm) and incubated at room temperature for 6 days. Five ml of spore suspension ( $10^6$  spores/ml) of each fungal isolate was inoculated into the mixture of sterilized ground rice bran and tap water. They were then incubated at room temperature for 7 days to obtain the inocula of planting media.

### Inoculation of toxigenic *A. flavus* and test fungi into planting media, and seed planting

Pasteurized planting media were placed in black polyethylene bags (15 kg/bag). The inocula of planting media were then mixed with planting media at the time of germinated-seed planting (1 seed<sup>^</sup>ag).

Treated planting media were inoculated with the combined use of (1) toxigenic and nontoxigenic *A. flavus*, (2) toxigenic *A. flavus* and *A. niger*, (3) toxigenic *A. flavus* and *T. harzianum*. The comparison of toxigenic *A. flavus* and each test fungal isolate was 1:1 (one portion was equivalent with 30 g of inoculum). Planting media inoculated only with each fungal isolate and uninoculated planting media were used as controls. Six replicates (6 bags) were used for each treatment (including the controls). The bags with the peanut plants were placed under greenhouse conditions.

Two watering treatments of peanut plants were carried out, i.e. (1) watering until harvest and (2) not watering for 15 days before harvest. Sterilized tap water was used for watering peanut plants. The individual plants were watered with the same volumes of sterilized tap water.

### Sampling of planting media

Sampling of planting media were conducted :

Before peanut-seed planting, but after each treated planting medium as well as each control (except uninoculated planting media) have been mixed with fungal

inocula homogeneously. They were then placed in clean polyethylene bags (about 1 kg/bag/treatment or control).

- At the time of inflorescence and at the time of harvest. The peanut plants were pulled prior to sampling of planting media. Each treated planting medium as well as each control was mixed homogeneously; they were then placed in clean polyethylene bags (about 1 kg/bag/treatment or control).

Each sample of planting media (about 1 kg) was divided several times manually to obtain working samples for moisture content and fungal analysis, and for reserved sample.

**Determination of moisture contents of planting media; population of toxigenic *A. flavus* and test fungi in planting media before peanut seed planting, at the time of inflorescence and at the time of harvest**

Moisture content of planting media derived from each bag was determined using oven method at 105°C for 12 hours (Johnson *et al.* 1960). The population of each fungal isolate in planting media before peanut seed planting, at the time of inflorescence and at the time of harvest was determined using serial dilution method, followed by pour plate method (Johnson and Curl 1972) on PDA containing chloramphenicol (100 mg/L media).

**Sampling of peanut plants at the time of inflorescence**

At the time of inflorescence (25 - 30 days after planting), peanut plants grown on treated planting media as well as on the controls, were pulled. Their roots, stems, petioles, leaves and flowers were then cut into pieces (10 pieces/plant) about 1 cm long, surface sterilized using 0.5% Na-hypochloride for one minute, rinsed with sterile distilled water, dried on sterilized filter paper, and finally plated on Potato Dextrose Agar (PDA) containing chloramphenicol (100 mg/L media). The percentage of each fungal isolate colonization in various plant parts was determined using the following formula:

$$\begin{array}{l}
 \text{Number of plant part pieces} \\
 \text{Colonized by each fungal isolate} \\
 \text{plant part pieces Colonized} = \frac{\text{Number of plant part pieces}}{\text{Number of plant part pieces}} \times 100\% \\
 \text{by each fungal isolate} \qquad \qquad \qquad \text{plated on agar media}
 \end{array}$$

**Determination of moisture content of peanut kernels; population of toxigenic *A. flavus* and test fungi, and aflatoxin content in peanut kernels**

Peanuts were harvested at 90 days after planting. They were then shelled manually and aseptically. Peanut kernels derived from treated planting media and

the controls were divided using a box sample divider to obtain working samples for moisture content, fungal and aflatoxin analyses. The moisture content of peanut kernels was determined using the oven method (BSI 1995). The population of toxigenic *A. flavus* and test fungi was determined using serial dilution method, followed by pour plate method on PDA containing chloramphenicol (100 mg/L media). Aflatoxin content in peanut kernels was determined using Thin Layer Chromatography method (AO AC 1995).

### **Statistical analyses**

The experimental design used in this study was Completely Randomized Factorial Design with two factors, i.e. the kind of fungal inocula in planting media and watering of peanut plants. The kinds of fungal inocula consisted of (1) the combined use of toxigenic and nontoxigenic *A. flavus*, (2) toxigenic *A. flavus* and *A. niger*, (3) toxigenic *A. flavus* and *T. harzianum*, (4) toxigenic *A. flavus*, (5) nontoxigenic *A. flavus*, (6) *A. niger*, (7) *T. harzianum*, and (8) uninoculated planting media. Peanut plants (1) were watered until harvest and (2) not watered for 15 days before harvest.

## **RESULTS AND DISCUSSION**

### **Moisture content of planting media before peanut seed planting, at the time of inflorescence and harvest**

The fungal inocula did not give any significant difference on the moisture contents of planting media at the time of inflorescence. Besides, the moisture contents of planting media before peanut planting were not significantly different from the time of inflorescence. It indicated that up to the time of inflorescence, the moisture contents of planting media were always stable, although the media were inoculated with various fungal inocula.

The moisture content of planting media treated with various fungal inocula and the time of watering decreased from before peanut planting up to harvest. Fungal inocula and their interaction with time of watering were not significantly different with the decrease of moisture contents, while time of watering showed significant differences.

Table 1 shows that the decrease in moisture content of planting media not watered for 15 days before harvest (22.9%) was higher than that watered until harvest (11.1%).

Table 1. The decrease in moisture content of planting media before peanut planting until harvest time

Treatment	Decrease of moisture content (%)
Watered planting media	11.09 a
Not watered planting media	22.85 b

Numbers followed by the same letter do not differ significantly according to Duncan Multiple Range Test at 95% confidence level

### Population of toxigenic *A. flavus* and test fungi in planting media before peanut planting, at the time of inflorescence and at the time of harvest

In the planting media, populations of toxigenic *A. flavus* and test fungi increased at the time of inflorescence as well as at the time of harvest. At the time of inflorescence, fungal inocula gave significant differences on the increase of toxigenic *A. flavus* and test fungal populations. At the time of harvest, fungal inocula and time of watering gave significant differences on the increase of fungal populations.

The increase of toxigenic *A. flavus* population in planting media inoculated with the combined use of toxigenic and nontoxigenic *A. flavus* was lower compared with planting media inoculated (1) only with toxigenic *A. flavus*, (2) with the combined use of toxigenic *A. flavus* and *A. niger*, and (3) with the combined use of toxigenic *A. flavus* and *T. harzianum*, either at the time of inflorescence or at the time of harvest (Table 2).

The increase of each test fungal population in either the planting media inoculated with each test fungi or in the planting media inoculated with the combined use of toxigenic *A. flavus* and each test fungi were higher than the increase of toxigenic *A. flavus* population in the planting media inoculated only with toxigenic *A. flavus* and in the planting media inoculated with the combined use of toxigenic *A. flavus* and test fungi, either at the time of inflorescence or at the time of harvest (Table 2). It indicated that the growth of toxigenic *A. flavus* was more inhibited compared with the growth of test fungi, either in the planting media inoculated with each fungal isolate or in the planting media inoculated with the combined use of toxigenic *A. flavus* and test fungi.

At the time of harvest, the increase of fungal population in the watered planting media was higher than that not watered for 15 days before harvest (Table 3). It was related to the decrease of moisture content which was higher in the not watered planting media compared with the watered planting media (Table 1), consequently fungal growth was inhibited.

Table 2. The increase of toxigenic *Aspergillus flavus* and test fungal populations in the planting media inoculated with various inocula at the time of inflorescence and harvest

Fungal inocula in the planting media	Increase of fungal population (cfu/g dry weight)			
	Fungi	At the time of inflorescence	At the time of harvest	
			Transformed into log x	Not transformed
Toxigenic <i>Aspergillus flavus</i>	Toxigenic <i>A. flavus</i>	362 214 ab	4.6421 ab	53 655
Toxigenic <i>A. flavus</i> vs nontoxigenic <i>A. flavus</i>	Toxigenic <i>A. flavus</i>	282 333 a	4.4666 a	39 727
Toxigenic <i>A. flavus</i> vs <i>A. niger</i>	Toxigenic <i>A. flavus</i>	513 562 ab	4.8380 b	115 011
Toxigenic <i>A. flavus</i> vs <i>Trichoderma harzianum</i>	Toxigenic <i>A. flavus</i>	624 964 ab	4.6148 ab	51 481
Nontoxigenic <i>A. flavus</i>	Nontoxigenic <i>A. flavus</i>	1 533 069 b	5.7928 cd	694 390
Toxigenic <i>A. flavus</i> vs nontoxigenic <i>A. flavus</i>	Nontoxigenic <i>A. flavus</i>	3 838 215 c	5.6359 c	619 960
<i>A. niger</i>	<i>A. niger</i>	2 955 578 c	5.9010 cd	1 024 393
Toxigenic <i>A. flavus</i> vs <i>A. niger</i>	<i>A. niger</i>	3 809 040 c	6.0054 d	1 135 405
<i>T. harzianum</i>	<i>T. harzianum</i>	1 385 558 ab	5.8040 cd	807 209
Toxigenic <i>A. flavus</i> vs. <i>T. harzianum</i>	<i>T. harzianum</i>	1 294 316 ab	5.6993 cd	852 893

Numbers followed by the same letter in the same column do not differ significantly according to Duncan Multiple Range Test at 95% confidence level

Table 3. The increase of fungal population in watered and not watered planting media at harvest time

Treatment	Increase of fungal population (cfu/g dry weight)	
	Transformed into log x	Not transformed
Watered planting media	5.4753 a	697 710
Not watered planting media	5.2047 b	381 114

Numbers followed by the same letter do not differ significantly according to Duncan Multiple Range Test at 95% confidence level

At the time of harvest in the uninoculated planting media, three fungal isolates were found: nontoxigenic *A. flavus*, *Cladosporium cladosporioides* and *T. harzianum*. It was assumed that fungal contamination originated from the air or from peanut seeds.

### **Toxigenic *A. flavus* and test fungal invasion in peanut plants at the time of inflorescence**

Pitt *et al.* (1991) stated that the invasion of toxigenic *A. flavus* in peanut plants could originate from the soils or from peanut seeds grown under greenhouse conditions.

Toxigenic *A. flavus* invasion was observed in peanut plants where the planting media were inoculated with the fungus as well as its combination with the test fungi (Table 4). Its percentage of invasion occurring in plant parts adjacent to the soils (root and stem) was higher than the other parts (petioles and leaves). Pitt *et al.* (1991) reported that *A. flavus* could invade all plant parts, with the highest invasion occurring in plant parts adjacent to the soils.

The percentage of toxigenic *A. flavus* invasion in plants where the planting media were inoculated with the combined use of toxigenic and nontoxigenic *A. flavus*, toxigenic *A. flavus* and *A. niger* was lower than that only inoculated with toxigenic *A. flavus*. Nevertheless, the percentage of toxigenic *A. flavus* invasion was higher in plants where the planting media were inoculated with the combined use of the fungus and *T. harzianum* (Table 4). In the planting media inoculated with the combined use of toxigenic *A. flavus* and nontoxigenic *A. flavus*, also the combined use of toxigenic *A. flavus* and *A. niger*, the invasion of toxigenic *A. flavus* were only in the roots, while in the planting media inoculated only with toxigenic *A. flavus*, this fungus could invade roots, stems and flowers. In the planting media inoculated with toxigenic *A. flavus* and *T. harzianum*, toxigenic *A. flavus* could invade all the plant parts. The percentage of toxigenic *A. flavus* invasions was related to the increase of toxigenic *A. flavus* population in the planting media (Table 2). The lowest increase of toxigenic *A. flavus* population was in the planting media inoculated with the combined use of toxigenic and nontoxigenic *A. flavus*, while the highest was in the planting media inoculated with the combined use of toxigenic *A. flavus* and *T. harzianum*.

In the planting media inoculated only with toxigenic *A. flavus* as well as those inoculated with the combined use of toxigenic *A. flavus* and *T. harzianum*, toxigenic *A. flavus* could also invade flowers (Table 4). It is related to the position of peanut flowers, i.e. adjacent to the planting media. If the flowers come in contact with the planting media, they could be invaded by toxigenic *A. flavus*.

Toxigenic *A. flavus* derived from the planting media can systemically invade peanut plant parts adjacent to the soil up to the plant parts above the soil surface. The fungus can also invade young peanut plants.

Based on microscopic examination, little or no damage was observed in the cells of stems invaded by *A. flavus* (Pitt *et al.* 1991). Toxigenic *A. flavus* invasion did not cause any disease in peanut plants.



Table 4. The percentage of plant parts colonized by toxigenic *Aspergillus flavus* and test fungi at the time of inflorescence

Fungal inocula in planting media	Fungi	Plant parts colonized by fungi (%)					Remark
		Root	Stem	Petiole	Leaf	Flower	
Uninoculated	Toxigenic <i>A. flavus</i>	0	8.3	0	0	0	It was assumed that fungal contamination originated from air or seeds
	Nontoxigenic <i>A. flavus</i>	6.6	18.3	0	0	5	
	<i>A. niger</i>	10	0	10	0	8.7	
	<i>Fusarium</i> sp.	13.3	0	0	0	0	
	<i>Gilmaniella</i> sp.	0	15	0	0	0	
	<i>Nigrospora</i> sp.	0	0	0	6.6	0	
Toxigenic <i>A. flavus</i>	Toxigenic <i>A. flavus</i>	10*	18.3	0	0	5	* Contaminated by <i>Fusarium</i> sp. (31.6%)
Nontoxigenic <i>A. flavus</i>	Nontoxigenic <i>A. flavus</i>	26.6	43.3	0*	0*	16.6	* Contaminated by <i>Nigrospora</i> sp. (8.3%)
<i>A. niger</i>	<i>A. niger</i>	56.6	60	43.3	23.3	51.6	All plant parts were contaminated by <i>Fusarium</i> sp. (85%)
<i>Trichoderma harzianum</i>	<i>T. harzianum</i>	46.6	10	3.3	0	0	
Toxigenic <i>A. flavus</i> vs nontoxigenic <i>A. flavus</i>	Toxigenic <i>A. flavus</i>	3.3*	0	0	0	0	* Contaminated by <i>Fusarium solani</i> (33.3%)
	Nontoxigenic <i>A. flavus</i>	68.3	40*	6.6	0	22.5	* Contaminated by <i>Fusarium</i> sp. (8.3%)
Toxigenic <i>A. flavus</i> vs <i>A. niger</i>	Toxigenic <i>A. flavus</i>	1.6*	0	0	0	0	* Contaminated by <i>Sphaeropsis</i> sp. (20%)
	<i>A. niger</i>	21.6	16.6	8.3	10	6.8	
Toxigenic <i>A. flavus</i> vs <i>T. harzianum</i>	Toxigenic <i>A. flavus</i>	16.6	21.6	5	9.1	11.6*	* Contaminated by <i>Exosporium</i> sp. (16.6%) and <i>Tubercularia</i> sp. (8.3%)
	<i>T. harzianum</i>	20	13.3	5	3.3	0	

Test fungi could also colonize roots, stems, petioles, leaves and flowers (Table 4). Some plant parts, either on planting media inoculated with toxigenic *A. flavus*, test fungi, or uninoculated planting media, were contaminated with other fungi such as *Exosporium* sp., *Fusarium* spp., *Gilmaniella* spp., *Sphaeropsis* sp. and *Tuber-*

*cularia* sp. (Table 4). It was assumed that the fungal contamination originated from the air or peanut seeds.

### Moisture content of peanut kernels at the time of harvest

The fungal inocula and their interaction with time of watering did not give any significant difference on the moisture content of peanut kernels at the time of harvest, while only time of watering gave significant differences. The moisture content of peanut kernels derived from watered plants until harvest was higher than those of not watered plants for 15 days before harvest (Table 5).

Table 5. Moisture content of peanut kernels in watered and not watered planting media at harvest time

Treatment	Moisture content (%)
Watered planting media	38.38 a
Not watered planting media	33.58 b

Numbers followed by the same letter do not differ significantly according to Duncan Multiple Range Test at 95% confidence level

### The effects of nontoxigenic *A. flavus*, *A. niger* and *T. haniaum* on toxigenic *A. flavus* infection and aflatoxin production in peanut kernels

From six replications derived from the planting media inoculated with toxigenic *A. flavus* and its combination with test fungi, in general only one replication of peanut kernels was infected by the fungus (Table 6). It indicated that the capability of toxigenic *A. flavus* in infecting peanut kernels was relatively low.

Toxigenic *A. flavus* population in peanut kernels derived from the plants where the planting media were inoculated with the combination of toxigenic *A. flavus* and each test fungal isolate was lower than that of only inoculated with toxigenic *A. flavus* (Table 6). It indicated that the test fungal isolates inoculated into planting media could inhibit toxigenic *A. flavus* infection in peanut kernels.

Aflatoxin was only detected in peanut kernels derived from the plants where the planting media were only inoculated with toxigenic *A. flavus* and the plants were always watered until harvest time. Aflatoxin B<sub>1</sub> was produced with a concentration of 32 ppb (Table 7). It indicated that toxigenic *A. flavus* (21 cfu/g dry weight) in peanut kernels (Table 6) could produce aflatoxin.

Aflatoxin was not detected in peanut kernels derived from the plants where the planting media were inoculated with a combination of toxigenic *A. flavus* and test fungi, either in watered plants or unwatered plants for 15 days before harvest, because the test fungi could inhibit aflatoxin production of toxigenic *A. flavus*. Inhibition of aflatoxin production caused by the test fungi could be affected by the

dose of biocontrol agent inocula. According to Dorner *et al.* (1998), aflatoxin production in peanut kernels decreased with the increase of the dose of biocontrol agents inocula (nontoxigenic *A. flavus* and *A. parasiticus*) inoculated into the soils. Aflatoxin content in peanut kernels with the dose of biocontrol agents inocula 0, 2, 10 and 50 g/m were 337.6, 73.7, 34.8 and 33.3 ppb, respectively.

Table 6. Toxigenic *Aspergillus flavus* and test fungal populations in peanut kernels at harvest \*)

Fungal inocula and time of watering	Fungi	Fungal population (cfu/g dry weight)					
		Replication					
		1	2	3	4	5	6
Toxigenic <i>A. flavus</i> - watered	Toxigenic <i>A. flavus</i>	0	21	0	0	0	0
Toxigenic <i>A. flavus</i> -not watered	Toxigenic <i>A. flavus</i>	0	0	0	0	0	0
Nontoxigenic <i>A. flavus</i> - watered	Nontoxigenic <i>A. flavus</i>	0	0	0	0	0	0
Nontoxigenic <i>A. flavus</i> - not watered	Nontoxigenic <i>A. flavus</i>	0	0	0	0	0	0
<i>A. niger</i> - watered	<i>A. niger</i>	0	0	0	0	0	0
<i>A. niger</i> - not watered	<i>A. niger</i>	0	0	0	4	0	0
<i>T. harzianum</i> - watered	<i>T. harzianum</i>	26	0	0	30	0	0
<i>T. harzianum</i> - not watered	<i>T. harzianum</i>	0	0	0	0	0	0
Toxigenic <i>A. flavus</i> vs nontoxigenic <i>A. flavus</i> - watered	Toxigenic <i>A. flavus</i> Nontoxigenic <i>A. flavus</i>	10 0	0 0	0 0	0 0	0 0	0 0
Toxigenic <i>A. flavus</i> vs nontoxigenic <i>A. flavus</i> - not watered	Toxigenic <i>A. flavus</i> Nontoxigenic <i>A. flavus</i>	0 0	0 0	0 10	0 0	0 0	0 0
Toxigenic <i>A. flavus</i> vs <i>A. niger</i> - watered	Toxigenic <i>A. flavus</i> <i>A. niger</i>	0 0	0 0	0 0	0 0	0 0	0 6
Toxigenic <i>A. flavus</i> vs <i>A. niger</i> - not watered	Toxigenic <i>A. flavus</i> <i>A. niger</i>	10 72	0 0	0 0	0 53	0 0	0 0
Toxigenic <i>A. flavus</i> vs <i>T. harzianum</i> - watered	Toxigenic <i>A. flavus</i> <i>T. harzianum</i>	0 26	0 0	0 0	0 5	0 0	0 0
Toxigenic <i>A. flavus</i> vs <i>T. harzianum</i> - not watered	Toxigenic <i>A. flavus</i> <i>T. harzianum</i>	0 65	0 0	0 0	0 35	0 0	0 0

\*) Peanut seeds derived from the plants where the planting media were treated with various fungal inocula and time of watering

Table 7. Aflatoxin B<sub>1</sub> content of peanut kernels at harvest time \*)

Treatment	Aflatoxin B <sub>1</sub> content (ppb)
Toxigenic <i>A. flavus</i> - watered	32
Toxigenic <i>A. flavus</i> - not watered	Not detected
Toxigenic <i>A. flavus</i> vs nontoxigenic <i>A. flavus</i> - watered	Not detected
Toxigenic <i>A. flavus</i> vs nontoxigenic <i>A. flavus</i> - not watered	Not detected
Toxigenic <i>A. flavus</i> vs <i>A. niger</i> - watered	Not detected
Toxigenic <i>A. flavus</i> vs <i>A. niger</i> - not watered	Not detected
Toxigenic <i>A. flavus</i> vs <i>T. harzianum</i> - watered	Not detected
Toxigenic <i>A. flavus</i> vs <i>T. harzianum</i> - not watered	Not detected

\*) Peanut seeds derived from the plants where the planting media were treated with various fungal inocula and time of watering.

## CONCLUSIONS

At the time of harvest, decrease in the moisture content of peanuts in the planting media not watered for 15 days before harvest was higher than those watered until harvest.

The lowest population of toxigenic *A. flavus* was in the planting media inoculated with the combined use of toxigenic and nontoxigenic *A. flavus* at the time of inflorescence and at the time of harvest.

Toxigenic *A. flavus* could invade the roots, stems and flowers of peanut plants. The lowest percentage of invasion was on the plant parts where the planting media were inoculated with the combined use of toxigenic and nontoxigenic *A. flavus*.

The moisture content of peanut kernels originated from plants watered until harvest was higher than those not watered for 15 days before harvest.

The population of toxigenic *A. flavus* in peanut kernels derived from the plants where the planting media were inoculated with the combined use of toxigenic *A. flavus* and each test fungi, was lower than that only inoculated with toxigenic *A. flavus*. It indicated that the test fungi inoculated into planting media could inhibit toxigenic *A. flavus* infection in peanut kernels.

Aflatoxin B<sub>1</sub> was only detected in peanut kernels originated from one plant where the planting medium was only inoculated with toxigenic *A. flavus* and the plant watered until harvest.

Toxigenic *A. flavus* infection and aflatoxin B<sub>1</sub> production were not influenced by the planting media which were not watered for 15 days before harvest.

## ACKNOWLEDGEMENT

The authors gratefully acknowledge the financial support of the Government of Indonesia. Thanks are due to the Research Institute for Legumes and Tuber Crops (RILET), Malang, in providing peanut seeds var. Komodo. We also thank Ms. Shanti Saraswati, Department of Biology, Faculty of Mathematics and Natural Sciences, Bogor Agricultural University, and the technicians of the Laboratory of Pest and Disease Management, SEAMEO BIOTROP for their valuable assistance in conducting the experiment.

## REFERENCES

- AOAC. 1995. Natural toxins. *In* E. Scott (ed.). Official Methods of Analysis of Natural Poison. 6<sup>th</sup> ed. Association of Official Analytical Chemists, Arlington. Chap. 49. p. 1-5.
- BS1. 1995. Oilseeds-determination of Moisture and Volatile Matter Content. British Standard International.
- Dharmaputra, O.S., A.S.R. Putri, I. Retnowati and S. Ambarwati. 2001. Soil mycobiota of peanut fields at Wonogiri reGENCY, Central Java: their effect on the growth and aflatoxin production of *Aspergillus flavus* *in vitro*. BIOTROP No. 17: 3058
- Domer, J.W., R.J. Cole and P.D. Blankenship. 1998. Effect of inoculum rate of biological control agents on preharvest aflatoxin contamination of peanuts. *Biological Control* 12: 171-176
- Domer, J.W., R.J. Cole and D.T. Wicklow. 1999. Aflatoxin reduction in corn through field application of competitive fungi. *Journal of Food Protection* 62(6): 650-656
- Ehrlich, K., A. Ciegler, M.A. Klich and L. Lee. 1985. Fungal competition and mycotoxin production on corn. *Experimentia* 41: 691 - 693.
- Johnson, L.F., E.A. Curl, J.H. Bond and H.A. Fribourg. 1960. Methods for Studying Soil Microflora-Plant Disease Relationships. Burgess Publishing Company, Minneapolis.
- Johnson, L.F. and E.A. Curl. 1972. Methods for Research on the Ecology of Soil-borne Plant Pathogens. Burgess Publishing Company, Minneapolis.
- Pitt, J.I., S.K.. Dyer and S. McCammon. 1991. Systemic invasion of developing peanut plants by *Aspergillus flavus*. *Letters in Applied Microbiology* 13: 16-20.
- Pitt, J.I. and A.D. Hocking. 1996. Current knowledge of fungi and mycotoxins associated with food commodities in Southeast Asia. *In* Highley, E. and G.I. Johnson (eds). *Mycotoxin Contamination in Grains*. Pp. 5-10. ACIAR Technical Reports 37, Canberra.
- Wicklow, D.T., B.W. Horn, O.L. Shotwell, C.W. Hesseltine and R.W. Caldwell. 1988. Fungal interference with *Aspergillus flavus* infection and aflatoxin contamination of maize grown in a controlled environment. *Phytopathology* 78: 68 - 74.