

Correction of Abeditehrani, H., Dijk, C., Sahragard Toghchi, M., & Arntz, A. (2020). Integrating Cognitive Behavioral Group Therapy and Psychodrama for Social Anxiety Disorder: An Intervention Description and an Uncontrolled Pilot Trial

Clinical Psychology in Europe, 2021, Vol. 3(4), Article e7727, <https://doi.org/10.32872/cpe.7727>

Published (VoR): 2021-12-23

Correction note to:

Abeditehrani, H., Dijk, C., Sahragard Toghchi, M., & Arntz, A. (2021). Integrating cognitive behavioral group therapy and psychodrama for social anxiety disorder: An intervention description and an uncontrolled pilot trial. *Clinical Psychology in Europe*, 2(1), Article e2693. <https://doi.org/10.32872/cpe.v2i1.2693>

In the originally published version of the above mentioned article, there was a typographical error in Table 4. In the second row ("LSAS"), an incorrect value in the "p" column was provided (.19). Instead, the correct data for "LSAS" under the "p" column is: .019. This change has no effect on the conclusions drawn in the article as the authors had already explained correctly that there was a significant decrease ($p = .019$) in social anxiety symptoms assessed with the LSAS. The corrected table can be found below (see Table 4). The authors apologize for any inconveniences caused.

Table 4

Pretest and Posttest Comparison for the CBPT Intervention

Scale	Pre		Post		<i>t</i> (4)	<i>M</i> difference [CI 99%]			Cohen's <i>d</i>	Hedges' <i>g</i>
	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>		<i>LL</i>	<i>UL</i>	<i>p</i>		
BFNE	35.60	7.02	28.40	4.10	2.86	-4.39	18.79	.046	1.03	0.82
LSAS	99.40	16.99	58.40	24.81	3.82	-8.44	90.44	.019	2.41	1.93
SADS	14.40	5.64	11.80	7.73	1.31	-6.56	11.76	.261	0.46	0.37
PAS	133.20	13.88	131.60	17.21	0.20	-34.67	37.87	.849	-0.12	-0.09
OPQ	56.20	23.18	35.80	16.63	3.22	-8.74	49.54	.032	0.88	0.70



Scale	Pre		Post		<i>t</i> (4)	<i>M</i> difference [CI 99%]			Cohen's <i>d</i>	Hedges' <i>g</i>
	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>		<i>LL</i>	<i>UL</i>	<i>p</i>		
OCQ	64.80	23.47	47.00	26.67	5.95	4.03	31.57	.004	0.76	0.61
BDI	19.60	5.86	12.60	8.20	2.03	-8.82	22.82	.111	1.19	0.96
QOLI	29.40	21.31	36.00	25.17	-0.88	-41.13	27.93	.429	0.31	0.25

Note. Observed Means (*M*) and Standard Deviations (*SD*) for the Pre and Post assessment points; results of *t*-test analyses (*t*, *p*-value) and effect sizes Cohen's *d* and Hedges' *g*. BFNE = Brief Fear of Negative Evaluation; LSAS = Liebowitz Social Anxiety Scale; SADS = Social Avoidance and Distress Scale; PAS = Personal Attitude Scale-II; OPQ = Social cost and probability by the Outcome Probability Questionnaire; OCQ = Outcome Cost Questionnaire; BDI = Beck Depression Inventory; QOLI = Quality of Life Inventory. Cohen's *d* was estimated as $d = (\text{mean pre-post change}) / (\text{pretest } SD)$. Hedges' *g* was calculated as follows: $g = J * d$, with $d = \text{Cohen's } d$; $J = (1 - 3 / (4 * df - 1))$; $df = N - 1$. The sign of the effect size was chosen so that a positive effect size indicates improvement and negative effect size represents worsening.

EACLIPT

Clinical Psychology in Europe (CPE) is the official journal of the European Association of Clinical Psychology and Psychological Treatment (EACLIPT).



leibniz-psychology.org

PsychOpen GOLD is a publishing service by Leibniz Institute for Psychology (ZPID), Germany.