



UNIVERSITY OF TORONTO  
LIBRARIES

CPOJ

ISSN: 2561-987X

PROFESSIONAL OPINION

# Canadian Prosthetics & Orthotics Journal

All articles are permanently available online to the public without restrictions or subscription fees. All articles are free to be used, cited, and distributed, on condition that appropriate acknowledgment is included. Authors are the *copyright* holders of their original contributions and grant the Canadian Prosthetics & Orthotics Journal (CPOJ) a license to publish the article and identify itself as the original publisher. CPOJ articles are licensed under the Creative Commons Attribution 4.0 International License.



CPOJ Website: <https://jps.library.utoronto.ca/index.php/cpoj/index>

Editorial Office: [cpoj@online-publication.com](mailto:cpoj@online-publication.com)

ISSN 2561-987X

VOLUME 1, ISSUE 2

2 0 1 8

## PROFESSIONAL OPINION

## THE IMPORTANCE OF ADDITIONAL MID SWING TOE CLEARANCE FOR AMPUTEES

Lechler K\*, Kristjansson K

Össur Medical Office, R&amp;D Össur ehf, Reykjavik, Iceland.

## ABSTRACT

Increased prosthetic hip to toe distance and insufficient mid swing toe clearance of a prosthetic foot is a well-recognized inadequacy for lower limb prosthesis users with wide and possible grave consequences on their ambulation capabilities. Most important are increased risk of falls and abnormal compensatory gait patterns with secondary unwanted physical effect like generally decreased mobility, muscular-skeletal pain and joint degeneration, i.e. osteoarthritis, with possible significant health economic effect. Even though insufficient toe clearance can be device related and technically or even intentionally induced for attaining equal length of the lower extremities in a neutral standing position or the stance phase, it is important to be aware of available technical solutions that can counteract the problem, like swing phase dorsiflexing feet, vacuum suspension systems, polycentric axis knees rather than single-axis knees and adequate knee flexion in early swing and swing-flexion assistance in the case of bionic knees.

## CITATION

Lechler K, and Kristjansson K. The importance of additional mid swing toe clearance for amputees. Canadian Prosthetics & Orthotics Journal. Volume1, Issue2, No.1, 2018. DOI: <https://doi.org/10.33137/cpoj.v1i2.30813>

## KEYWORDS

Prosthetics, Toe clearance, Lower limb prosthesis, Mobility, Prosthetic foot, Polycentric axis knees, Amputee.

## INTRODUCTION

The shortest distance between the ground and the toe during level ground walking, referred to as the minimum toe clearance (MTC), occurs during mid swing phase (mid swing toe clearance: MSTC) and for abled bodies this averages to be around 15 mm.<sup>1</sup> This is the point in time during the gait phase where tripping is most likely to occur.<sup>2-4</sup> Trip-related falls have been identified as the leading cause of falls for community-dwelling, middle-aged and older adults.<sup>5-7</sup> Looking at elderly non-amputated women only, trips amount for up to 33% of the causes for their falling.<sup>8</sup> Whereas women are more likely to fall while walking men are more likely to fall from loss of support by an external object rather than tripping, as well as falling while sitting down or rising from sitting.<sup>9</sup> While most elderly amputees are men<sup>10</sup> the risk of amputees in general to fall over a one-year period is 50% or higher.<sup>11</sup>

Historically the first reported prosthetic leg offering additional toe clearance in swing dates back to 1805

(Fig. 1,2,3)<sup>12</sup>, but still today most conventional prosthetic feet do not provide features for additional MSTC. This is despite the fact that historically various stumble preventing mechanisms for both trans-tibial and trans-femoral community ambulators have been offered since the early days of the industrial revolution, underlining the longstanding awareness of the importance of trips and stumbles as a cause for falls by amputees.<sup>12</sup>

## WHY DO MANY PROSTHETICS SOLUTIONS NOT PROVIDE FOR MINIMUM TOE CLEARANCE?

The main reason for insufficient MSTC is the fact that conventional prosthetic feet do not dorsiflex during swing which places the MSTC on the prosthetic side below the MTC.<sup>13</sup> Another well recognized reason is the vertical displacement between the socket and the residual during swing phase (pistoning) resulting in functional elongation of the prosthetic leg.<sup>14-16</sup>

## \*CORRESPONDING AUTHOR

**Knut Lechler**, Medical Director Prosthetics, Össur ehf, Grjóthals 1-5, 110 Reykjavik, Iceland.


Email: [klechler@ossur.com](mailto:klechler@ossur.com)

Tel: +49 151 50459110.

**Disclosure:** Full time employees of a medical device manufacturer, Össur ehf.

**DOI:** <https://doi.org/10.33137/cpoj.v1i2.30813>

Equal leg length in the stance phase yields an increased functional leg length in the swing phase of at least 16mm for a non-amputee.<sup>17,18</sup> The functional prosthetic leg length with single axis knee is even further increased,<sup>18</sup> resulting in a 22 mm lower MSTC compared to polycentric knees.<sup>19</sup> In a 3D simulation with data from trans-femoral amputees' single-axis knees provided the least clearance (4.6 cm) during prosthetic swing phase, followed by four-bar knees (5.8 cm), and the two ankle mechanisms (6.4 and 8.7 cm).<sup>19</sup>



**Figure 1.** Anglesey TT and TF (1805) **Figure 2.** Erfurt TF (1880) **Figure 3.** Martin TT (1857)

### BIOMECHANICAL AND PHYSICAL EFFECT OF INSUFFICIENT TOE CLEARANCE

For the purpose of overcoming amputees' experience of vertical displacement between socket and residual and the associated functional elongation of the prosthetic leg in swing phase<sup>14-16</sup> the prosthesis is commonly aligned significantly shorter than the sound side. Amongst 113 trans-femoral and trans-tibial amputees 70 % were found to have a significant leg length discrepancy radiographically while balanced standing due to shorter alignment of the prosthesis compared to the sound side.<sup>20</sup> Gaunard et. al. showed that 66% of trans-femoral amputees had leg length discrepancies and for 57% the prosthetic side was shorter.<sup>21</sup>

Back pain is very common in lower limb amputees<sup>22</sup> and has been linked to poor prosthetic fit and alignment, postural changes, leg-length discrepancy, amputation level, and general deconditioning.<sup>23</sup> The significance of this approach of leg length misalignment for the purpose of trying to make up for functional leg length increase during swing and the associated decrease in the MSTC is that leg length discrepancy can lead to various

health problems like back pain<sup>24</sup> and studies have shown it to contribute to back pain in amputees.<sup>25,26</sup>

Not trying to adjust for the functional prosthetic leg length increase and to fit the prosthesis at an equal length to the sound side also carries with it a problem as pistoning in the socket and the changes in the trajectory of the prosthetic foot changes the point in time at which the physical maximum functional leg length occurs. For this additional functional swing phase leg length and time shift the amputees will try to compensate by developing their individual adaptation mechanisms. One is to elevate the contralateral side by early heel rise resulting in what is referred to as "Vaulting". Another is an arc-like movement of the prosthetic limb referred to as "Circumduction" and thirdly a lateral trunk inclination with definite raising of the hip on the prosthesis side. As a matter of fact, these three are the basic and main gait abnormalities characteristic for lower limb amputee in various degrees of combinations, even obvious to a lay observer.

Both stationary and functional leg length discrepancies cause abnormal forces on bones and joints likely to lead to degenerative conditions like osteoarthritis which has been found to be more prevalent in the contralateral limb than the residual limb of people with amputation.<sup>27,28</sup> The increased prevalence of osteoarthritis is of concern, especially for people who have lived with an amputation for a longer time.<sup>28</sup> This actually turns out to be the case for many traumatic amputees who have been shown to have a significantly increased risk of suffering from osteoarthritis.<sup>29</sup>

### THE HEALTH ECONOMIC RELEVANCE OF TOE CLEARANCE

The proven relationship between insufficient toe clearance to trips and hence falls bring this feature into the scope of possible significant health economic effects. Although there is scarce published literature on the economic costs of falls within the amputee population, the costs of falls among older adults have been well studied. The estimated average one-year cost attributed to an elderly adult falling and requiring subsequent medical attention is between \$3,408 and \$4,872. Furthermore, if the fall results in a hospitalization, the cost may increase up to \$35,144 on average. Within this population, it is estimated that up to 1 in every 9 falls will lead to hospitalization.<sup>30</sup>

A study on falls of amputees shows that up to 40% of their falls result in an injury and every other fall necessitates medical attention, which is higher than for the non-amputated elderly which has been estimated to be 30%.<sup>31</sup> One study on trans-femorals indicated an estimated cost of \$25,652 at 6 months, for falls resulting in hospitalization, similar to the costs within the elderly.<sup>32</sup> Direct medical costs of all falls in the USA was \$31.3 billion in 2015, up from \$30.3 billion in 2012.<sup>33</sup>

Possible secondary health economic effect through back- and musculoskeletal pain and degenerative joint disease is also of real significance with osteoarthritis already accounting for 2% of years lived with disabilities.<sup>34</sup> OA is a rising concern specifically to amputees because they have a higher risk to suffer from hip and knee osteoarthritis.<sup>35</sup> Osteoarthritis often coexist with an amputation as pain does<sup>28</sup> and even further limits amputees in their mobility.<sup>36</sup> Osteoarthritis in the contralateral limb knee is 17 times higher than in age-matched non-amputees<sup>35</sup> and knee pain in unilateral amputees is twice as common.<sup>35</sup> Imaging studies have confirmed the increased prevalence of degenerative changes in the contralateral knee.<sup>37,38</sup>

Since TTAs are known to load their intact limb to a greater extent than their prosthetic limb during gait,<sup>23</sup> the marked asymmetry in knee pain and degeneration suggests that mechanical loading factors are likely an important contributor to the increased incidence of OA. Not only asymmetries but also minor compensations can increase stress on the contralateral limb and possibly predispose the long-term prosthesis wearer to premature degenerative arthritis.<sup>39-40</sup>

## REMEDIES FOR INSUFFICIENT TOE CLEARANCE

The most straight forward and natural action is the provision of prosthetic feet that dorsiflex in the swing phase and thereby compensating for the elongation in swing phase at the same time allowing for leg length equality in the stance phase or when standing.


Motorized and actively dorsiflexing prosthetic ankles have shown to significantly increase MSTC<sup>41</sup> by about 70% and increase in MSTC of  $25.6 \pm 5.4$  mm has been shown to lead to a reduction in falls<sup>42</sup> over a one-year period for community dwelling amputees. Hydraulic ankles with dorsiflexion in swing have also been shown to increase the MSTC.<sup>43</sup>

A recent study has shown in addition to increased MSTC during level ground walking that this can also be provided for during incline and decline walking (Figure 4).<sup>44</sup> An anecdotal user report from this same trial states: "My work environment involves walking through sloped tunnels and up/down stairs. I usually trip 1-3 times per day with my prescribed foot, but I never tripped with the Proprio-Foot!". Several studies had previously looked at and compared the MSTC of conventional ESAR feet versus dorsiflexing ankles (Table 1).

**Table 1:** Comparison of prosthetic side toe clearances in mm.

	Toe Clearance on the Prosthetic Side		
	Level (mm)	Slope (mm)	Decline (mm)
Fallers, Rosenblatt 2017	12.3		
Non-Fallers, Rosenblatt 2017	25.6		
Hydraulic, Johnson 2014	20.7		
Motorized, Fradet 2010		16.0	12.0
Motorized, Rosenblatt 2014	33.6		
Motorized, Lamers 2018*	14.6*	18.4*	11.7*

\* compared and in addition to sound side



**Figure 4.** MTC with MPA(Proprio), MPA-Locked and conventional ESAR in level ground.<sup>44</sup>

Other technical solutions that can improve the MSTC are vacuum suspension systems for reduced pistoning and functional swing phase prosthetic leg length increase. A study by Rosenblatt where falling rate and pattern was tracked over a one-year period for 27 amputees (15 vacuum assisted socket suspension (VASS) users, 12 non-VASS users) showed 75% less risk of VASS trans-tibial amputee users having multiple falls compared to non-VASS users.

Even though less effective than providing dorsiflexion ankles, selecting polycentric knees over single axis knees also helps with the toe clearance as previously outlined.<sup>19</sup> Also providing a bionic knee like the Rheo Knee that has a rapid foot off velocity results in a higher MSTC compared to single axis knees.<sup>45</sup> Additionally, the rapid toe off velocity and lower hip torque qualities of the Rheo Knee results in earlier toe off at a slower than preferred walking speed and significantly less vaulting of the intact ankle, compared to for example a hydraulic knee ( $P = 0,028$ ).<sup>46</sup>

## SUMMARY

Providing sufficient toe clearance on the prosthetic side allows aligning the prosthesis more equally and favorable influence in short term on low back pain and in longer term on side effects like lumbar scoliosis and osteoarthritis. Most importantly added toe clearance should result in fewer trips and stumbles resulting in decrease in falls. All of this can possibly result in significantly less health economic burden. A large study on patients having undergone hip arthroplasty ( $n=753$ ) shows that leg length discrepancy reduces satisfaction and functional outcome after hip arthroplasty.<sup>47</sup> Reducing risks of falls reduces the degree of disability of amputees by improving their mobility and hence improves their quality of life.<sup>48</sup>

## REFERENCES

1. Winter DA, Patla AE, Frank JS, Walt SE. Biomechanical walking pattern changes in the fit and healthy elderly. *Phys Ther.* 1990; 70, 340–347.
2. Begg RK, Tirosh O, Said CM, Sparrow WA, Steinberg N, Levinger P, Galea MP. Gait training with real-time augmented toe-ground clearance information decreases tripping risk in older adults and a person with chronic stroke. *Front Hum Neurosci.* 2014;8:243. <https://doi.org/10.3389/fnhum.2014.00243>
3. Mills PM, Barrett RS. Swing phase mechanics of healthy young and elderly men. *Hum Mov Sci.* 2001; 20, 427–446. [https://doi.org/10.1016/S0167-9457\(01\)00061-6](https://doi.org/10.1016/S0167-9457(01)00061-6)
4. Moosabhojy MA, Gard SA. Methodology for determining the sensitivity of swing leg toe clearance and leg length to swing leg joint angles during gait. *Gait Posture.* 2006; 24, 493–501. <https://doi.org/10.1016/j.gaitpost.2005.12.004>
5. Berg WP, Alessio HM, Mills EM, Tong C. Circumstances and consequences of falls in independent community-dwelling older adults. *Age Ageing.* 1997; 26, 261–268.
6. Li W, Keegan TH, Sternfeld B, Sidney S, Quesenberry CP Jr, Kelsey JL. Outdoor falls among middle-aged and

older adults: a neglected public health problem. *Am J Public Health.* 2006; 96, 1192–1200. DOI: 10.2105/AJPH.2005.083055

7. Blake AJ, Morgan K, Bendall MJ, Dallosso H, Ebrahim SB, Arie TH, Fentem PH, Bassey EJ. Falls by elderly people at home: prevalence and associated factors. *Age Ageing.* 1988; 17, 365–372.

8. Crenshaw JR, Bernhardt KA, Achenbach SJ, Atkinson EJ, Khosla S, Kaufman KR, Amin S. The circumstances, orientations, and impact locations of falls in community-dwelling older women. *Arch Gerontol Geriatr.* 2017;73, 240–247. <https://doi.org/10.1016/j.archger.2017.07.011>

9. Yang Y, Van Schooten KS, Sims-Gould J, McKay HA, Feldman F, Robinovitch SN. Sex differences in the circumstances of falls among older adults in long-term care. *Innov Aging.* 2017; 431–432. <https://doi.org/10.1093/geroni/igx004.1549>

10. Ziegler-Graham K, MacKenzie EJ, Ephraim PL, Trivison TG, Brookmeyer R. Estimating the Prevalence of Limb Loss in the United States: 2005 to 2050. *Arch Phys Med Rehabil.* 2008; 89, 422–429. <https://doi.org/10.1016/j.apmr.2007.11.005>

11. Miller WC, Speechley M, Deathe B. The prevalence and risk factors of falling and fear of falling among lower extremity amputees. *Arch Phys Med Rehabil.* 2001; 82, 1031–1037. DOI: [10.1053/apmr.2001.24295](https://doi.org/10.1053/apmr.2001.24295)

12. Knoche W. Prothesen der unteren Extremität. Die Entwicklung vom Altertum bis 1930. ISBN:3-00-018655-7.

13. Eshraghi A, Osman NAA, Gholizadeh H, Karimi M, Ali S. Pistoning assessment in lower limb prosthetic sockets. *Prosthet Orthot Int.* 2012; 36, 15–24. <https://doi.org/10.1177/0309364611431625>

14. Lilja M, Johansson T, Öberg T. Movement of the tibial end in a PTB prosthesis socket: a sagittal X-ray study of the PTB prosthesis. *Prosthet Orthot Int.* 1993;17, 21–26.

15. Eshraghi A, Osman NAA, Gholizadeh H, Karimi M, Soodmand E. Gait Biomechanics of Individuals with Transtibial Amputation: Effect of Suspension System. *PLoS ONE* 9, e96988 (2014). <https://doi.org/10.1371/journal.pone.0096988>

16. Erikson U, James U. Roentgenological Study of Certain Stump—Socket Relationships in Above-knee Amputees with Special Regard to Tissue Proportions, Socket Fit and Attachment Stability. *Ups J Med Sci.* 1973;78, 203–214.

17. Winter, D. A. Foot trajectory in human gait: a precise and multifactorial motor control task. *Phys Ther.* 1992;72, 45–53.

18. Gard SA, Childress DS, Uellendahl JE. The Influence of Four-Bar Linkage Knees on Prosthetic Swing-Phase Floor Clearance. *J Prosthet Orthot.* 1996; 8, 34.

19. Sensinger JW, Intawachirarat N, Gard SA. Contribution of prosthetic knee and ankle mechanisms to swing-phase foot clearance. *IEEE Trans Neural Syst Rehabil Eng.* 2013; 21, 74–80. DOI: [10.1109/TNSRE.2012.2224885](https://doi.org/10.1109/TNSRE.2012.2224885)
20. Friberg O. Biomechanical significance of the correct length of lower limb prostheses: a clinical and radiological study. *Prosthet Orthot Int.* 1984;8, 124–129.
21. Gaunaud I, Gailey R, Hafner BJ, Gomez-Marin O, Kirk-Sanchez N. Postural asymmetries in transfemoral amputees. *Prosthet Orthot Int.* 2011; 35, 171–180. <https://doi.org/10.1177/0309364611407676>
22. Ehde DM, Smith DG, Czerniecki JM, Campbell KM, Malchow DM, Robinson LR. Back pain as a secondary disability in persons with lower limb amputations. *Arch Phys Med Rehabil.* 2001; 82, 731–734. <https://doi.org/10.1053/apmr.2001.21962>
23. Gailey R, Allen K, Castles J, Kucharik J, Roeder M. Review of secondary physical conditions associated with lower-limb amputation and long-term prosthesis use. *J Rehabil Res Dev.* 2008; 45, 15–29.
24. Defrin R, Benyamin SB, Aldubi RD, Pick CG. Conservative correction of leg-length discrepancies of 10mm or less for the relief of chronic low back pain. *Arch Phys Med Rehabil.* 2005; 86, 2075–2080. <https://doi.org/10.1016/j.apmr.2005.06.012>
25. Mahar RK, Kirby RL, MacLeod DA. Simulated leg-length discrepancy: its effect on mean center-of-pressure position and postural sway. *Arch Phys Med Rehabil.* 1985; 66, 822–824.
26. Biering-sørensen F. Physical measurements as risk indicators for low-back trouble over a one-year period. *Spine.* 1984; 9, 106–119.
27. Giles LGF, Taylor JR. Low-back pain associated with leg length inequality. *Spine.* 1981;6, 510–521.
28. Nolan L, Wit A, Dudziński K, Lees A, Lake M, Wychowański M. Adjustments in gait symmetry with walking speed in trans-femoral and trans-tibial amputees. *Gait Posture.* 2003; 17, 142–151. [https://doi.org/10.1016/S0966-6362\(02\)00066-8](https://doi.org/10.1016/S0966-6362(02)00066-8)
29. Kulkarni J, Adams J, Thomas E, Silman A. Association between amputation, arthritis and osteopenia in British male war veterans with major lower limb amputations. *Clin Rehabil.* 1998;12, 274–279.
30. Farrokhi S, Mazzone B, Yoder A, Grant K, Wyatt M. A Narrative Review of the Prevalence and Risk Factors Associated With Development of Knee Osteoarthritis After Traumatic Unilateral Lower Limb Amputation. *Mil Med.* 2016;181, 38–44. <https://doi.org/10.7205/MILMED-D-15-00510>
31. Bohl AA, Fishman PA, Ciol MA, Williams B, LoGerfo J, Phelan EA. A Longitudinal Analysis of Total 3-Year Healthcare Costs for Older Adults Who Experience a Fall Requiring Medical Care. *J Am Geriatr Soc.* 2010; 58, 853–860. <https://doi.org/10.1111/j.1532-5415.2010.02816.x>
32. Kaufman, K. Risk factors and costs associated with accidental falls among adults with above-knee amputations: a population-based study. *American Orthotic and Prosthetic Association* 2016. (Mayo Clinic). <http://www.aopanet.org/resources/research/>
33. Mundell B, Maradit Kremers H, Visscher S, Hoppe K, Kaufman K. Direct medical costs of accidental falls for adults with transfemoral amputations. *Prosthet Orthot Int.* 2017; 41, 564–570. <https://doi.org/10.1177/0309364617704804>
34. Burns ER, Stevens JA, Lee R. The direct costs of fatal and non-fatal falls among older adults — United States. *J Safety Res.* 2016; 58, 99–103. <https://doi.org/10.1016/j.jsr.2016.05.001>
35. March L, Smith EU, Hoy DG, Cross MJ, Sanchez-Riera L, Blyth F, Buchbinder R, Vos T, Woolf AD. Burden of disability due to musculoskeletal (MSK) disorders. *Best Pract Res Clin Rheumatol.* 2014; 28, 353–366. <https://doi.org/10.1016/j.berh.2014.08.002>
36. Struyf PA, Van Heugten CM, Hitters MW, Smeets RJ. The prevalence of osteoarthritis of the intact hip and knee among traumatic leg amputees. *Arch Phys Med Rehabil.* 2009; 90, 440–446. <https://doi.org/10.1016/j.apmr.2008.08.220>
37. Geertzen JH, Bosmans J C, Van Der Schans CP, Dijkstra PU. Claimed walking distance of lower limb amputees. *Disabil Rehabil.* 2005; 27, 101–104. <https://doi.org/10.1080/09638280400009345>
38. Norvell DC, Czerniecki JM, Reiber GE, Maynard C, Pecoraro JA, Weiss NS. The prevalence of knee pain and symptomatic knee osteoarthritis among veteran traumatic amputees and nonamputees. *Arch Phys Med Rehabil.* 2005; 86, 487–493. <https://doi.org/10.1016/j.apmr.2004.04.034>
39. Lemaire ED, Fisher FR. Osteoarthritis and elderly amputee gait. *Arch Phys Med Rehabil.* 1994;75, 1094–1099.
40. Hurley GRB, McKenney R, Robinson M, Zdravec M, Pierrynowski MR. The role of the contralateral limb in below-knee amputee gait. *Prosthet Orthot Int.* 1990;14, 33–42.
41. Rosenblatt NJ, Bauer A, Rotter D, Grabiner MD. Active dorsiflexing prostheses may reduce trip-related fall risk in people with transtibial amputation. *J Rehabil Res Dev.* 2014;51, 1229–1242. <http://dx.doi.org/10.1682/JRRD.2014.01.0031>
42. Rosenblatt NJ, Bauer A, Grabiner MD. Relating minimum toe clearance to prospective, self-reported, trip-related stumbles in the community. *Prosthet Orthot Int.* 2017;41(4):387-392. DOI:[10.1177/0309364616650085](https://doi.org/10.1177/0309364616650085)

43. Johnson L, De Asha AR, Munjal R, Kulkarni J, Buckley JG. Toe clearance when walking in people with unilateral transtibial amputation: effects of passive hydraulic ankle. *J Rehabil Res Dev.* 2014; 51, 429–437. DOI: [10.1682/JRRD.2013.05.0126](https://doi.org/10.1682/JRRD.2013.05.0126)

44. Lamers E. Effects of a new adaptive ankle prosthesis on level and sloped walking. 42nd Annu Meet Am Soc Biomech. Rochester MN USA (2018).

45. Johansson JL, Sherrill DM, Riley PO, Bonato P, Herr H. A clinical comparison of variable-damping and mechanically passive prosthetic knee devices. *Am J Phys Med Rehabil.* 2005; 84, 563–575. DOI: [10.1097/01.phm.0000174665.74933.0b](https://doi.org/10.1097/01.phm.0000174665.74933.0b)

46. Prinsen EC, Nederhand MJ, Sveinsdóttir HS, Prins MR, Van der Meer F, Koopman HFJM, Rietman JS. The influence of a user-adaptive prosthetic knee across varying walking speeds: A randomized cross-over trial. *Gait Posture.* 2017; 51, 254–260. DOI: [10.1016/j.gaitpost.2016.11.015](https://doi.org/10.1016/j.gaitpost.2016.11.015)

47. Röder C, Vogel R, Burri L, Dietrich D, Staub LP. Total hip arthroplasty: leg length inequality impairs functional outcomes and patient satisfaction. *BMC Musculoskelet Disord.* 2012; 13, 95. DOI: [10.1186/1471-2474-13-95](https://doi.org/10.1186/1471-2474-13-95)

48. Asano M, Rushton P, Miller WC, Deathe BA. Predictors of quality of life among individuals who have a lower limb amputation. *Prosthet Orthot Int.* 2008; 32, 231–243. DOI: [10.1080/03093640802024955](https://doi.org/10.1080/03093640802024955)

## AUTHORS BIOGRAPHY



**Mr. Knut Lechler**, Össur Medical Director Prosthetics, is a CPO-D at Össur ehf, Reykjavik, Iceland and has been in that position for 7 years. He graduated in 1991 as CPO (Master) and used to work as a freelancer before he started in the industry in 1999 (Flex-Foot Inc.). Since 2001 he is with Össur. Knut Lechler is an active member of the ISO-TC 168 WG1.



**Dr. Kristleifur Kristjansson**, Medical Officer and VP of Clinical Affairs, is a medical doctor and has been in his current position at Össur for 6 years. Dr. Kristjansson was a VP of Clinical Collaborations at deCODE genetics for 16 years and holds a consulting position at the University Hospital in Reykjavik Iceland as a pediatrician and geneticist.