

Zea mays L.

احمد عبد الواحد مرعي*

نوفل عدنان صبري*

عادل هايس عبد الغفور*

* كلية الزراعة / جامعة الانبار

2009 /
(ART-B40 S182 P-3 P-4 Syn23)
(RCBD)

.%91.12

(500)

. 500

(Zea mays L.)

(2004)

/ 4.222

/ 2.249

2005
(2005 FAO)

.(2002)

. 2010 / 10 / 8

. 2011 / 3 / 23

(1982 Narsinghani ,Singh)

2009

ART-B40	1
S 182	2
P-3	3
P-4	4
Syn23	5

(RCBD)
 (75 × 25)
 / 400 N:P (18 × 18)
 :
 .%40 (30)
 .%50 (60)
 500
 (δ²p) (δ²g) (h²_{b.s}) (δ²e)
 (rgij)
 -(1972 Das. ; 1998)

$$\delta^2 G = \frac{msv - mse}{r},$$

$$\delta^2 E = mse,$$

$$\delta^2 p = \delta^2 G + \delta^2 E$$

$$h^2_{b.s} = \frac{\delta^2 G}{\delta^2 p} \times 100$$

.1
.2009


$h^2_{b.s}$	δ^2P	δ^2E	δ^2G	
91.12	220.82	19.61	201.21	X1
77.72	204.22	45.50	158.72	X2
39.79	2.23	1.34	0.89	X3
74.24	2.01	0.52	1.49	X4 /
Zero	20.35	24.94	-4.59	X5 /
Zero	0.26	0.36	-0.10	X6 500
65.05	2699.81	943.63	1756.18	X7
71.64	101576	28804	72772	X8

.2

(r^2)	()	()	/		500 ()	()	
-0.399	0.823**	0.504	-0.351	0.330	0.277	-0.739*	()
	-0.359	-0.957**	0.114	-0.707*	-0.350	0.986**	(r^2)
		0.317	0.525	0.074	0.191	-0.686*	()
			0.173	0.183	0.178	-0.844**	()
				0.477	-0.064	0.677*	/
					-0.572	-0.011	/
						-0.458	() 5000

$$0.632 = r_{0.05} \quad *$$

$$0.765 = r_{0.01} \quad **$$



.3

(r^2)	()	()	/		5000 ()	()	
-0.354	0.896**	0.317	-0.264	0.120	-0.020	-0.597	()
	-0.250	-0.629	0.165	-0.006	-0.240	0.686*	(r^2)
		0.291	-0.428	0.103	0.127	-0.699*	()
			0.135	0.623	0.302	-0.621	()
				0.419	-0.310	0.368	/
					0.365	-0.216	
						-0.171	() 5000

$$0.632 = r_{0.05} \quad *$$

$$0.765 = r_{0.01} \quad **$$

.1998 .
 - .
 .107 : .
 .2001 .
 (1) .
 11: .(3)
 .2009 .
Zea mays L.
 - .
 .2010 .
 - .
 .1990 .
 .399 : . -
 .2009 .
Sorghum bicolor L.
 105: . -
 .2004 .
Sesamia cretica Led.
 120: . - . Zea mays L.
 .2002 .

- Akbar, M., Muhamad, S. Faqir. M. A., M. Y. A. and Rashid. 2008. Combining ability analysis in maize under normal and height temperature condition. J. Agric. Res. 46 (1): 27-38.
- Asrar, R. S. and V. Saleem and G. M. Subhani. 2007. Correlation and path coefficient analysis in maize (*Zea mays* L.). J. Agric. Res. , 45 (3):20-32
- Das, P.K. 1972. Studies on selection for yield in wheat: An application of genotypic and phenotypic correlation, path coefficient analysis and discriminate function J. Agric. Sci. 49:238-243.
- Devi, I. S., S. Muhamad and S. Mohamad. 2001. Character association and path coefficient analysis of grain yield and yield components in double cross of maize (Zea mays L.). Crop Res. Hisar. 21 (3): 335-359.
- F.A.O. 2005. Year Book. Production. V. 55.

- Kabdal, M. K., S. S. Verma, N. Ahmed and V. B. S. Panwar. 2003. Genetic variability and correlation studies of yield and its attributing characters in maize (*Zea mays* L.). Indian Agric. Sci. Dig. 23(2):137-139.
- Mohsan, Y. C., D. K. Singh and N. V. Rao. 2002. Path coefficient analysis for oil and grain yield in maize (*Zea mays* L.) genotypes. Nat. J. Pl. Impr. 4(1):75-77.
- Parh, D. K., M. A. Hamid, M. H. Rehman and M. Z. I. Talukdar. 1988. Correlation, path coefficient and selection indices in open-pollinated maize. Bangladesh. J. Agric. 15(1):69-74.
- Singh, S. P., A. A. Pianchi and V. G. Narsinghani. 1982. Character correlations and selection indices in F2 population of wheat. Indian J. Agric. Sci. 52: 420 - 424.
- Venugopal, M., N. A. Ansari and T. Rajanikanth. 2003. Correlation and path analysis in maize (*Zea mays* L.). Crop Res. Hisar. 25 (3): 525-529.
- Viola, G. M. Ganesh, S. S. Reddy and C. V. S. Kumar. 2003. Studies on correlation and path coefficient analysis of elite baby corn (*Zea mays* L.). Indian prog. Agric. 3(1-2):22-25.
- Yousaf, M. and M. Saleem. 2001. Correlation analysis of S1 families of maize for grain yield and its components. Pak. Int. J. Agric. Biol. 3(4):387-388.

VARIANCES ESTIMATION AND GENOTYPIC, PHENOTYPIC CORRELATION AND BROAD HERITABILITY PERCENTAGE IN MAIZE (*Zea mays* L.)

**Adel Hais Abudlgaffor Nawfal Adnan S. Ahmed Abdulwahed M.
College of Agriculture -Anbar University**

ABSTRACT

A field experiment was conducted at the farm of field crops in the foundation of agricultural Research / Abu Gharib during the autumn season of 2009. Five genotype of maize were used (ART-B40, S182, P-3, P-4, Syn23), as a randomized complete block design with three replications, in order to study variances, broad sense heritability percentage and genotypic and phenotypic correlation coefficient were estimated.

The results showed that the values of variances were varied among studied characters. The values of the genotypic and phenotypic variance were more than environment variance for all characters. The higher values for broad sense heritability appeared in plant height as 91.12%.

The genotypic correlation was positive and high significant between yield of plant and leaf area and positive signification with the number of rows/ear, while it was negative and highly significant between yield and negative and significant with (plant height, ear height, ear length). While it was negative and no significant among yield of plant and number of grain/row and the weight of 500 grains). Mean while the phenotypic correlation was positive and high significant too, between yield of plant and leaf area, was positive and no significant with number of rows/ear, while it was negative and high significant with ear height and negative and significant with plant height, ear length, while it was negative and no significant with number of grain/row and the weight of 500 grain.