



Correction

Dermatol Pract Concept. 2019;10(1):e2020027. November 10, 2019. DOI: <https://doi.org/10.5826/dpc.1001a27>

Alopecia areata incognita and diffuse alopecia areata: clinical, trichoscopic, histopathological, and therapeutic features of a 5-year study

In the article titled “Alopecia areata incognita and diffuse alopecia areata: clinical, trichoscopic, histopathological, and therapeutic features of a 5-year study” by Alessandrini et al [1] (<https://dpcj.org/index.php/dpc/article/view/dermatol-pract-concept-articleid-dp0904a05>), published on Octo-

ber 31, 2019, Table 2 included an error. The last row should have read Clobetasol 0.05% foam. These authors have no conflicts of interest to disclose, and they contributed significantly to this publication. This article was corrected online on December 31, 2019. The editors regret the error.

Reference

1. Alessandrini A, Starace M, Bruni F, Brandi N, Baraldi C, Misciali C, Fanti PA, Piraccini BM. Alopecia areata incognita and diffuse alopecia areata: clinical, trichoscopic, histopathological, and therapeutic features of a 5-year study. *Dermatol Pract Concept.* 2019;9(4):272-277 . DOI: <https://doi.org/10.5826/dpc.0904a05>.