

Assessment of some salivary enzymes levels in type 2 diabetic patients with chronic periodontitis (Clinical and biochemical study)

Ghasaq A. Abdul-wahab, B.D.S. ⁽¹⁾

Maha A. Ahmed, B.D.S., M.Sc. ⁽²⁾

ABSTRACT

Background: Diabetic patients have been reported to be more susceptible to gingivitis and periodontitis than healthy subjects. Many intracellular enzymes like (alkaline phosphatase- (ALP), aspartate aminotransferase- (AST) and alanine aminotransferase- (ALT) that are released outside cells into the gingival crevicular fluid (GCF) and saliva after destruction of periodontal tissue during periodontitis. This study was conducted to determine the periodontal health status and the levels of salivary enzymes (ALP, AST and ALT) of the study and control groups and to correlate the levels of these enzymes with clinical periodontal parameters in each study group.

Subjects, Materials and Methods: One hundred subjects were enrolled in the study, with an age range of (35-50) years, only males were included. The subjects were divided into study groups (group-I consists of 30 patients with controlled type 2 diabetes mellitus (T2DM), group-II consists of 30 patients with uncontrolled T2DM, group-III consists of 25 patients non-diabetics, all of them have chronic periodontitis (CP) and group-IV consists of 15 apparently-systemically healthy subjects and have healthy periodontium, as control group. Unstimulated saliva samples were collected for biochemical analysis of salivary enzymes (ALP, AST and ALT). The clinical periodontal parameters including: plaque index (PLI), gingival index (GI), bleeding on probing (BOP), probing pocket depth (PPD) and clinical attachment level (CAL) were recorded for all subjects at four sites per tooth except third molars.

Results: All clinical periodontal and biochemical parameters were highest in uncontrolled T2DM with CP patients and all enzymes levels revealed highly significant differences between all pairs of the study and control groups except AST enzyme level which demonstrated a non-significant difference between controlled T2 diabetics with CP and non-diabetics with CP. There were weak correlations between all clinical periodontal parameters and biochemical parameters except between PPD and ALT enzyme in non-diabetics with CP group and between CAL and AST enzyme in uncontrolled T2 diabetics with CP which demonstrated highly significant strong positive correlations.

Conclusion: It was concluded that T2DM and poor glycemic control have negative impact on periodontal health status. Salivary enzymes were considered as good biochemical markers of periodontal tissue destruction and useful in diagnosis, monitoring and efficient management of periodontal diseases and T2DM.

Key words: Enzymes, saliva, type 2 diabetes mellitus, periodontal diseases. (J Bagh Coll Dentistry 2015; 27(1):138-143).

INTRODUCTION

Diabetes Mellitus (DM) is an extremely important disease from a periodontal standpoint. It is a multi-systemic metabolic disorder characterized by abnormal carbohydrate, protein and lipid metabolism, the cardinal biochemical feature of this disease is elevated levels of glucose in the blood (hyperglycemia) ⁽¹⁾

Chronic hyperglycemia has been closely associated with an inflammatory response that has been linked to many complications (microvascular and macrovascular complications) observed in diabetes ⁽²⁾. T2DM is the more prevalent of the two major categories of overt diabetes and accounts for 90-95% of all diabetic cases and usually has an adult onset; it combines insulin resistance (IR) and insulin secretory defect ^(1,3).

Periodontal diseases (PDs) are one of the most widely spread diseases of mankind; they are considered an inflammatory disorder that damages tissues. The most common form of PD is called chronic periodontitis (CP) which described as a

multifactorial, irreversible and cumulative condition, initiated and propagated through the complex interactions between periodontopathic bacteria and the host defense system ⁽⁴⁻⁶⁾. So, it's considered the most important cause of tooth loss during adulthood ⁽⁷⁾. Systemic diseases are among risk factors of PD. DM and PD are thought to be associated biologically; DM is believed to promote periodontitis through an exaggerated inflammatory response to the periodontal microflora ⁽⁸⁾.

It has been shown that uncontrolled or poorly controlled DM has the greater incidence of severe PD compared with those patients who are well controlled or have no DM, which has been more prevalent in persons with T2DM ⁽⁹⁻¹¹⁾. Saliva is a complex fluid that influences oral health through specific and non-specific physical and chemical properties. When disruptions in the quality or quantity of saliva occur, there will often be detrimental effects on oral and systemic health. Saliva is a mirror of general health. The diverse salivary constituents provide sources for assessment and monitoring of health and disease states ⁽¹²⁾.

(1) Master Student, Department of Periodontics, College of Dentistry, University of Baghdad.

(2) Assistant Professor, Department of Periodontics, College of Dentistry, University of Baghdad.

Host responses to PD include the production of different enzymes that are released by stromal, epithelial or inflammatory cells. In addition to this, various enzymes associated with cell injury and cell death can be detected in GCF and saliva. Several enzymes are evaluated for early diagnosis of PD such as ALP, AST and ALT⁽¹³⁻¹⁵⁾. There has been correlation between T2DM, PD and numerous markers in saliva, such as intracellular enzymes⁽¹⁶⁾. These enzymes may be used to test the activity of PDs⁽¹⁷⁾.

SUBJECTS, MATERIALS AND METHOD

The subjects consisted of 100 males with age range of (35-50) years. The subjects recruited for the study were patients attending the specialized center for Endocrinology and Diabetes in Baghdad, as well as, patients from the department of Periodontics, at teaching hospital, College of Dentistry, University of Baghdad. All the individuals were informed about the purposes of the investigation and consented to its protocol. The subjects were divided into:

- 1) **Study group (G-I):** Consists of (30) males with CP and T2DM, well controlled, the HbA1c were <7%.
- 2) **Study group II (G-II):** Consists of (30) males with CP and uncontrolled T2DM (moderately and poorly controlled), the HbA1c were >7%.
- 3) **Study group III (G-III):** Consists of (25) males with CP and without history of any systemic diseases.
- 4) **Control group IV (G-IV):** Consists of (15) males without history of any systemic diseases and with healthy periodontium, this was defined by GI scores <0.5⁽¹⁸⁾ and without periodontal pockets or clinical attachment loss. This group represents a base line data for the levels of salivary ALP, AST and ALT.

Inclusion criteria include T2DM patients (≥ 5 years) on oral hypoglycemic medication, body mass index level ranges between 18.5 kg/m² - 24.9 kg/m²⁽¹⁹⁾, all subjects were presenting at least 20 teeth and CP in patients was defined as the presence of at least four sites with PPD ≥ 4 mm and clinical attachment loss of (1-2) mm or greater, this made according to the international classification system for PD⁽²⁰⁾.

Exclusion Criteria include T1 and T2 diabetic patients taking insulin therapy, presence of other systemic diseases other than diabetes, presence of retinopathy, neuropathy or diabetic foot, patients who have undergone periodontal treatment and course of anti-inflammatory or antimicrobial therapy 3 months prior to the study, gross oral

pathology such as oral cancer and smoking or alcohol drinking.

From each subject, (5ml) of unstimulated whole saliva was collected; The collected saliva was centrifuged at 4000 rpm for 15 minutes and then the clear supernatant saliva was collected and kept frozen and stored at -20 ° C until biochemical analysis of salivary enzymes.

Clinical periodontal parameters examination was performed after salivary sample collection by using Michigan O periodontal probe on four surfaces (mesial, buccal/ labial, distal and lingual/ palatal) of all teeth except third molar. The collected data include:-

1. Assessment of Soft Deposits by the Plaque Index System (PLI)⁽²¹⁾.
2. Assessment of Gingival Inflammation by the Gingival Index System (GI)⁽¹⁸⁾.
3. Assessment of Gingival Bleeding on Probing (BOP)
4. Assessment of Probing Pocket Depth (PPD)
5. Assessment of Clinical Attachment Level (CAL)

For ALP enzyme analysis we used kit manufactured by (BIOMEREIUX ® Sa), while for AST and ALT enzymes analysis we used kits manufactured by (RANDOX /UK). The activity of ALP was determined by measuring its absorbance at 510 nm, while the activities of AST and ALT were determined by measuring the absorbance at 505 nm both by the spectrophotometer.

Descriptive statistics in the form of mean, standard deviation, percentage and inferential statistics in the form of Games –Howell, LSD, Chi-square and Pearson correlation were used in this study. The level of significance was accepted at P< 0.05, highly significance at P< 0.01 and non-significant at P> 0.05.

RESULTS

A-Clinical periodontal parameters:

The results showed that all clinical periodontal parameters were highest in uncontrolled T2 diabetics with CP followed by non-diabetics with CP then controlled T2 diabetics with CP except for PPD which was highest in uncontrolled followed by controlled T2 diabetics both with CP then non-diabetic patients with CP, as shown in (table -1).

Highly significant differences were demonstrated by using Chi-square test for the comparisons between each two study groups regarding the percentages of BOP sites in addition, comparisons between all pairs of the study groups revealed significant and highly significant differences (p<0.01) regarding the

other clinical periodontal parameters except for PPD and CAL between controlled T2 diabetics with CP and non-diabetics with CP which were non-significant differences ($P>0.05$), as shown in (table-2).

B- Biochemical analysis:

The obtained results have shown that the mean concentrations of salivary enzymes (ALP, AST and ALT) were highest in uncontrolled T2 diabetics with CP followed by controlled T2 diabetics with CP then non-diabetics with CP and finally control group, as shown in (table-3). Inter groups comparisons for all enzymes levels revealed highly significant differences ($p < 0.01$) between all pairs of the study and control groups except AST enzyme level which demonstrated a non-significant difference ($P>0.05$) between controlled T2 diabetics with CP and non-diabetics with CP (table-4).

C-Correlation of ALP, AST and ALT enzymes levels with clinical periodontal parameters:

There were weak correlations between all clinical periodontal parameters and biochemical parameters except between PPD and ALT enzyme in non-diabetics with CP group and between CAL and AST enzyme in uncontrolled T2 diabetics with CP which demonstrated highly significant positive strong correlations. Significant positive correlations were revealed between PLI and AST enzyme in controlled T2 diabetics with CP, between GI and ALP enzyme in uncontrolled T2 diabetics with CP, as well as, CAL with both AST enzyme in controlled T2 diabetics with CP and ALT enzyme in uncontrolled T2 diabetics with CP and non-diabetics with CP, as shown in (table-5).

Table 1: Statistical description (Mean ± SD) of (PLI, GI, PPD and CAL) and percentages of sites according to BOP scores for the study and control groups

Groups	PLI	GI	BOP		PPD	CAL
	Mean±SD	Mean±SD	Score 0	Score 1	Mean ±SD	Mean ±SD
G- I	1.62 ± 0.59	1.61 ± 0.63	61.39 %	38.61 %	4.47±0.63	4.187±1.578
G- II	2.69 ± 0.37	2.79 ± 0.39	2.16 %	97.84 %	5.03±0.43	5.913±1.439
G- III	2.05 ± 0.68	2.00 ± 0.55	49.76 %	50.24 %	4.42±0.56	4.74±1.836
G- IV	0.12 ± 0.09	0.09 ± 0.08	-	-	-	-

Table 2: Inter- groups Comparisons of the mean values of PLI, GI, PPD, CAL and the Percentages of BOP sites between all pairs of the study groups

Groups		PLI		GI		BOP		PPD		CAL	
		P	Sig.	P	Sig.	P	Sig.	P	Sig.	P	Sig.
G- I	G- II	0.000	HS	0.000	HS	0.000	HS	0.000	HS	0.000	HS
	G- III	0.042	S	0.046	S	0.000	HS	0.753	NS	0.209	NS
G- II	G- III	0.000	HS	0.001	HS	0.000	HS	0.000	HS	0.009	HS

Table 3: Statistical description (mean level in IU/L ± SD) of ALP, AST and ALT enzyme for the study and control groups

Groups	ALP	AST	ALT
	Mean± SD	Mean± SD	Mean± SD
G- I	9.487 ± 1.983	14.533 ± 1.624	10.27 ± 0.89
G- II	14.587 ± 2.541	18.433 ± 1.165	13.25 ± 1.96
G- III	7.888 ± 1.15	13.56 ± 2.053	8.34 ± 1.24
G- IV	5.993 ± 0.823	9.533 ± 1.506	5.07 ± 1.52

Table 4: Inter groups comparisons of the mean concentrations (IU/L) of ALP, AST and ALT enzymes between all pairs of the study and control groups by using Games-Howell Method

Groups		ALP		AST		ALT	
		P-value	Sig.	P-value	Sig.	P-value	Sig.
G- I	G- II	0.000	HS	0.000	HS	0.000	HS
	G- III	0.003	HS	0.233	NS	0.000	HS
	G- IV	0.000	HS	0.000	HS	0.000	HS
G- II	G- III	0.000	HS	0.000	HS	0.000	HS
	G- IV	0.000	HS	0.000	HS	0.000	HS
G- III	G- IV	0.000	HS	0.000	HS	0.000	HS

Table 5: Person's Correlation Coefficient (r) between clinical periodontal parameters and the levels of salivary enzymes for each study group

ALP enzyme	PLI		GI		BOP		PPD		CAL	
	r	P	r	p	r	p	r	p	r	p
G-I	0.058	0.380	0.221	0.120	0.137	0.469	0.023	0.452	0.224	0.117
G-II	0.215	0.127	0.379	0.020	0.024	0.901	0.042	0.414	0.202	0.142
G-III	0.034	0.435	0.012	0.478	0.001	0.996	0.012	0.477	0.003	0.494
AST enzyme	r	P	r	p	r	p	r	p	r	p
G-I	0.420	0.010	-0.222	0.119	-0.245	0.192	-0.099	0.302	0.379	0.019
G-II	0.110	0.281	0.006	0.487	-0.254	0.088	-0.003	0.494	0.563	0.001
G-III	0.105	0.308	0.185	0.189	0.222	0.286	0.12	0.284	0.198	0.171
ALT enzyme	r	P	r	p	r	p	r	p	r	p
G-I	-0.125	0.256	0.047	0.402	0.114	0.547	-0.119	0.265	0.115	0.273
G-II	-0.137	0.236	0.165	0.192	-0.050	0.792	-0.015	0.468	0.414	0.011
G-III	0.243	0.121	-0.269	0.097	0.365	0.073	0.523	0.004	0.424	0.017

DISCUSSION

In the present study significant statistical differences were found between controlled T2 diabetic patients with CP and non-diabetics with CP and highly significant differences were found between uncontrolled T2 diabetics and both controlled T2 diabetics with CP and non-diabetics with CP regarding PLI and GI. These were in agreement with other studies⁽²²⁻²⁴⁾ and in disagreement with Ibrahim and Abaas⁽²⁵⁾ and Sharma et al⁽²⁶⁾. These are explained by the fact that patients with diabetes tend to be systemically compromised and that their oral environment is also compromised due to the reduction in the buffering capacity and volume of their saliva, increased salivary viscosity and the change in bacterial flora^(27, 28). All these factors lead to higher accumulation of plaque and calculus. Diabetes is often associated with increased gingival inflammation in response to bacterial plaque as the inflammatory reactions are intensified during poor metabolic control⁽⁸⁾.

The result of this study revealed that the uncontrolled diabetics have more sites with bleeding on probing than controlled and non-diabetic groups and this was in agreement with Offenbacher et al⁽²⁹⁾ and in disagreement with Kumar et al⁽³⁰⁾.

Regarding PPD and CAL the results represent highly significant differences were found between uncontrolled T2 diabetic patients with CP and both controlled T2DM and non-diabetic patients with CP. These results were in agreement with studies^(31, 32) and in disagreement with other study⁽³³⁾. This result may be explained by the fact that poor glycemic control, with the associated increase in advanced glycation end products, renders the periodontal tissues more susceptible to destruction. The cumulative effects of altered cellular response to local factors, impaired tissue

integrity and altered collagen metabolism as the collagen in diabetic patients is aged and more susceptible to breakdown⁽³⁴⁾, these undoubtedly play a significant role in the susceptibility of diabetic patients to infections and destructive PD. There were increased BOP, PPD, increased tooth mobility and greater loss of attachment as the individuals with diabetes are twice as likely to exhibit attachment loss as non-diabetic individuals⁽³⁵⁾.

Numerous markers in saliva have been proposed as a diagnostic test for periodontal disease such as (ALP, AST and ALT) enzymes. These intracellular enzymes are increasingly being released by sick periodontal tissues into the GCF and saliva where their activity can be estimated. From the present study findings revealed that the level of ALP was higher in diabetic groups than in non-diabetics and control groups, The observed high enzyme activity can be attributed to increase in inflammation and bone turnover rate as ALP enzyme is produced by polymorphonuclear leukocytes, osteoblasts, macrophages, fibroblasts and plaque bacteria within periodontal tissues or pockets. The increased ALP activity is probably a consequence of destructive process in the alveolar bone in the advanced stages of PD⁽¹³⁾. The present findings revealed that there were highly significant differences in salivary ALP levels between all pairs of the study groups and with control group. This is in accordance with many other studies^(16, 35).

The result of this study showed highly significant difference in salivary AST level between all pairs of the study groups and with the control group except between controlled T2 diabetics and non-diabetics both with CP, which was non-significant. This result was in agreement with studies^(16, 36), regarding ALT enzyme the results revealed that there were highly significant

differences in enzyme level between all pairs of the study groups and with control group, this was in agreement with previous studies^(16,37). There were weak non-significant positive correlations between ALP and clinical periodontal parameters (PLI, BOP, PPD and CAL) in all study groups, while there is a significant positive correlation between ALP and GI in uncontrolled T2 diabetics with CP group. These results were in agreement with Kalburgi *et al*⁽³⁵⁾ and Sanikop *et al*⁽³⁸⁾ and in disagreement with Kumar Sharma⁽³⁹⁾. The possible explanation of weak non-significant correlation may be due to limited human sample size. Regarding the AST enzyme there were a highly significant strong positive correlation between salivary AST and CAL within uncontrolled T2 diabetic patients with CP, and significant positive correlation between AST enzyme and both PLI and CAL parameters within controlled T2 diabetic patients with CP. These findings agreed with Abdul- Hadi⁽⁴⁰⁾ and in disagreement with other study⁽¹⁵⁾. The findings revealed that there were weak non-significant correlations between the ALT enzyme and the clinical periodontal parameters except between ALT enzyme and PPD in non-diabetics with CP which was highly significant strong positive correlation and between ALT enzyme and CAL in (uncontrolled T2 diabetics and non-diabetics) with CP which was significant positive correlations. These results agreed with Abdul- Hadi⁽⁴⁰⁾ and Raiet *et al*⁽⁴¹⁾ and in disagreement with Todorovic *et al*⁽¹⁵⁾. The significant positive correlations between ALT enzyme with PPD and CAL parameters indicated that the enzyme activity increased with increasing severity of periodontitis. An increase in AST and ALT enzymes activities reflect the negative effect of the (T2DM) on the periodontal tissue especially the soft tissue since these enzymes are intracellular included in the metabolic processes of cells and they were indicators of a higher level of cellular damage and a reflection of metabolic change in the inflamed gingiva^(13,14,17).

REFERENCES

1. Radhika T, Kannan R. Diabetes mellitus and oral health. *J Orofacial Sciences* 2012; 4(1): 7-10.
2. Southerland JH, Taylor GW, Offenbacher S. Diabetes and Periodontal Infection: Making the Connection. *Clinical Diabetes* 2005; 23(4): 171-178.
3. Naemiratch B, Manderson L. Lay Explanations of Type 2 Diabetes in Bangkok, Thailand. *Anthropology and Medicine* 2007; (14) 1: 83-94.
4. American Academy of Periodontology. Tobacco use and the periodontal patient, (position paper) *J Periodontol* 1999; 70:1419-1427.
5. El-Qadri SS, Ta'ani DQ. Assessment of periodontal knowledge and periodontal status of an adult population in Jordan. *Int J Dent Hygiene* 2004; 2: 132-6.
6. Kinane DF. Causation and pathogenesis of periodontal disease. *Periodontology* 2000 2001; 25: 8-20.
7. Preshaw PM. Diabetes and periodontitis: what's it all about? *Practical Diabetes* 2013; 30(1): 9-10.
8. Katagiri S, Nitta H, Nagasawa T, Izumi Y, Kanazawa M. Effect of glycemic control on periodontitis in type 2 diabetic patients with periodontal disease. *J of Diabetes Investigation* 2013; 4(3): 320-5.
9. Pag RC, Offenbacher HE, Seymour GJ, Kornmank S. Advanced in the Pathogenesis of Periodontitis; Summary of Developments Clinical Implication and Future Directions. *Periodontol* 2000 1997; 14: 216-48.
10. Drisko CH. Non-Surgical Periodontal Therapy. *Periodontol* 2000 2001; 25: 77-88.
11. Fowler FB, Breault LO, Guemic MF. Periodontal Disease and its Association with Systemic Disease. *Mil Med* 2001; 166: 85-90.
12. Zhang L, Henson BS, Camargo PM, Wong DT. The Clinical Value of Salivary Biomarkers for Periodontal Disease. *J Periodontol* 2009; 51: 25-37.
13. Kaufman E, Lamster IB. Analysis of Saliva for Periodontal Diagnosis. *J Clin Periodontol* 2000; 27: 453-465
14. Ozmeric N. Advances in Periodontal Disease Markers. *Clin Chim Acta* 2004; 343: 1-16.
15. Todorovic T, Dozic I, Vicente-Barrero M, Ljuskovic B, Pejovic J, Marjanovic M, Knezevic M. Salivary enzymes and periodontal disease. *Med Oral Patol Oral Cir Bucal* 2006; 11: 115-9.
16. Ikekpeazu EJ, Neboh EE, Maduka IC, Anyanwu EG, Okenyi NS. Periodontal disease and type 2 diabetes: effects on salivary enzyme activities. *Int J Diabetes* 2011; 31(1): 9-13.
17. Numabe Y, Hisano A, Kamoi K, Yoshie H, Ito K, Kurihara H. Analy-sis of Saliva for Periodontal Diagnosis and Monitoring. *J Periodontol* 2004; 40: 115-9.
18. Loe H. The gingival index, the plaque index and the retention index system. *J Periodontal* 1967; 38(6): 610-6.
19. World Health Organization. WHO expert consultation. Appropriate body- mass index for Asian populations and its implications for policy and intervention strategies. *The Lancet* 2004; 363: 157-63.
20. Lang NP, Bartold PM, Cullinan M et al. International classification workshop. Consensus report: Chronic periodontitis. *Annals Periodontol* 1999; 4: 53.
21. Silness P, Loe H. Periodontal disease in pregnancy. *Acta Odontol Scand* 1964; 22: 121.
22. Serrano C, Perez C, Rodríguez M. Periodontal conditions in a group of Colombian type 2 diabetic patients with different degrees of metabolic control. *Acta Odontol Latinoam* 2012; 25(1):132-9.
23. Tanwir F, Tariq A. Effect of glycemic control on periodontal status. *J Coll Physicians Surg Pak* 2012; 22(6):371-4.
24. Casarin RCV, Barbagallo A, Meulman BT, Santos VR, Sallum EA, Nociti FH, Duarte PM, Casati MZ, Gonc, alves RB. Subgingival biodiversity in subjects with uncontrolled type-2 diabetes and chronic periodontitis. *J Periodont Res* 2013; 48: 30-6.
25. Ibrahem LM, , Abaas RF, Periodontal health status and biochemical study of Saliva among diabetics and non-diabetics (Comparative study). *Must Dent J* 2007; 4(1):1-4.

26. Sharma R, Raj SS, Vinod K, Reddy YG, Desai V, Bailoor D. Comparison of oral health indicators in type 2 diabetes mellitus patients and controls. J of Indian Academy of Oral Medicine and Radiology 2011; 23(3): 168-72.
27. Hintao J, Teanpaisan R, Chongsuvivatwong V, Dahlen G, Rattarasarn C. Root surface and coronal caries in adults with type 2 diabetes mellitus. Community Dent Oral Epidemiol 2007; 35(4): 302-309.
28. Preetha P, Kim K, Eun S, Inglehart, Rohr M. Diabetes and Oral Health: The Importance of Oral Health-Related Behavior 2011; 85: 264-72.
29. Offenbacher S, Barros S, Singer R, et al. Periodontal disease at the biofilm-gingival interface. J Periodontol. 2007; 78: 1911-25.
30. Kumar A, Pandey MK, Singh A, Mittra P, Kumar P. Prevalence and severity of periodontal diseases in Type 2 Diabetes Mellitus of Bareilly region (India). Int J Med Sci Public Health 2013; 2: 77-83.
31. Stojanović N, Krunic J, Cicmil S, Vukotić O. Oral health status in patients with diabetes mellitus type 2 in relation to metabolic control of the disease. Srp Arh Celok Lek 2010; 138(7-8): 420-4.
32. Haseeb M, Khawaja KI, Ataullah K, Munir MB, Fatima A. Periodontal disease in type 2 diabetes mellitus. J Coll Physician Surg Pak 2012; 22(8): 514-8.
33. Mealey BL. Periodontal Disease and Diabetes: A Two-Way Street. JADA 2006; 137(10): 26-31.
34. Carranza, Newman, Taki and Klokkevold. Carranza's Clinical Periodontology. 11th ed. Elsevier, Saunders: 2012. (Chapter 4,p 41-42) and (chapter 27, p 305-308).
35. Kalburgi V, Jenifer HD, Warad S, Bhola S, Chaudhari HL. Comparison of salivary alkaline phosphatase levels among diabetics and non-diabetics with chronic periodontitis. J Oral Health Research 2010; 1(4):147-52.
36. Al-Rubae EA, Kadum HA, Al-Braich MS. Salivary aspartate amino transferase and alanine amino transferase of non-insulin-dependents (Type2) diabetic patients. J Fac Med Baghdad 2010; 521(2): 212-214.
37. Dhiraj T, Chhaya T. Salivary proteome in periodontal diagnosis. Int J Pharma and Bio Sci 2012; 3(2): 241-245.
38. Sanikop S, Patil S, Agrawal P. Gingival crevicular fluid alkaline phosphatase as a potential diagnostic marker of periodontal disease. J Indian Soc Periodontol 2012; 16(4): 513-8.
39. Kumar R, Sharma G. Salivary Alkaline Phosphatase level as Diagnostic marker for periodontal disease. J. Int Oral Health 2011; 3(5): 81-86.
40. Abdul-Hadi MJ. Evaluation of salivary enzymes activities among patients with chronic periodontitis. A master thesis/ Department of Periodontology. College of dentistry, University of Baghdad, 2009.
41. Rai B, Kharb S, Anand S.C. Salivary Enzymes and Thiocyanate: Salivary Markers of Periodontitis among Smokers and Non-smokers; a Pilot Study. Adv in Med Dent Sci 2007; 1(1): 1-4.

الخلاصة

الخلفية: ان مرضى السكري أكثر عرضة لالتهاب اللثة والنساج من الأشخاص الأصحاء. العديد من الانزيمات داخل الخلايا مثل (الكالايين فوسفاتيز, اسبارتيت امينو ترانسفيريز و الانين امينوترانسفيريز) تفرز خارج الخلايا الى السائل الشقي اللثوي واللغاب بعد تدمير الأنسجة اللثوية اثناء النساج. لذا اعدت هذه الدراسة لتحديد الحالة الصحية للثة ومستويات الانزيمات اللعابية لمجموعات الدراسة والظابطة، ثم لربط مستويات هذه الانزيمات مع مؤشرات ما حول الاسنان السريرية في كل مجموعة دراسة.

الأشخاص، المواد والطرق: التحق مائة شخص في الدراسة مع الفئة العمرية من (٢٥-٥٠) سنة، وكانوا ذكور فقط. تم تقسيم الأشخاص الى مجموعتين الدراسة (المجموعة الاولى تتكون من ٣٠ مريضاً مع السكري من النوع الثاني المسيطر عليه، المجموعة الثانية تتكون من ٣٠ مريضاً مع السكري من النوع الثاني الغير مسيطر عليه، والمجموعة الثالثة تتكون من ٢٥ مريضاً غير المصابين بالسكري، كل منهم لديه نساج مزمن). والمجموعة الرابعة تتكون من ١٥ شخصاً اصحاء والثة لديهم صحية، كمجموعة ظابطة. تم جمع عينات من اللعاب الغير محفز من كل شخص شارك في الدراسة لاجراء التحليل الكيمائي الحيوي للانزيمات اللعابية (الكالايين فوسفاتيز، الانين امينوترانسفيريز والاسبارتيت امينوترانسفيريز). مؤشرات ما حول الاسنان السريرية بما في ذلك مؤشر الصفيحة الجرثومية، مؤشر التهاب اللثة، مؤشر النزف عند التسبير، عمق جيوب اللثة و مستوى الانسجة الرابطة سريريا سجلت لكل شخص في الدراسة ولاربعة اسطح في كل سن باستثناء الرخي الثالثة.

النتائج: كل مؤشرات ما حول الاسنان السريرية والمؤشرات الكيميائية كانت اعلى لمرضى السكري من النوع الثاني الغير مسيطر عليه ولديهم نساج مزمن وكل مستويات الانزيمات اظهرت فروقات معنوية عالية بين كل ازواج المجاميع الدراسة والظابطة ماعدا مستوى الانزيم الاسبارتيت امينوترانسفيريز بين مجموعتي مرضى السكري من النوع الثاني المسيطر عليه مع نساج مزمن وغير المصابين بالسكري مع نساج مزمن الذي اظهر فرق غير معنوي. هناك علاقة ضعيفة بين مؤشرات ما حول الاسنان السريرية والمؤشرات الكيميائية ماعدا بين مؤشر عمق جيوب اللثة وانزيم الانين امينوترانسفيريز لدى مجموعة غير المصابين بالسكري من النوع الثاني مع نساج مزمن وبين مؤشر مستوى الانسجة الرابطة سريريا وانزيم الاسبارتيت امينوترانسفيريز في مجموعة مرضى السكري من النوع الثاني الغير مسيطر عليه مع نساج مزمن التي اظهرت علاقة ايجابية قوية معنوية عالية.

الاستنتاج: تم استنتاج انه مرض السكري النوع الثاني وسوء السيطرة على نسبة السكري في الدم لها تأثير سلبي على صحة ما حول الاسنان. تعتبر الانزيمات اللعابية كعلامات كيميائية حياتية جيدة للانسجة اللثوية المحطمة وهذا يوفر فرصة جيدة للتشخيص، المراقبة والسيطرة الجيدة على امراض اللثة والسكري من النوع الثاني.