

RESEARCH ARTICLE

## Rock Formation Acid Mine Drainage in Epithermal Gold Mineralization, Pandeglang, Banten Province

Dudi Nasrudin Usman<sup>1,\*</sup>, Sri Widayati<sup>1</sup>, Sriyanti<sup>1</sup>, Era Setiawan<sup>2</sup>

<sup>1</sup>Bandung Islamic University, Mining Engineering Department, Mining, Bandung City, Indonesia

<sup>2</sup>Cibaliung Sumberdaya Corporation, General Manager, Mining, Pandeglang – Banten, Indonesia

\* Corresponding author: dudi.n.usman@gmail.com

Tel.: +62-81-3175-66144; fax: +62-22-426-3895

Received : Oct 7, 2019; Accepted: Nov 19, 2019.

DOI: 10.25299/jgeet.2019.4.4.3903

### Abstract

Mine acid water is acidic water and contains iron and sulfate, which is formed under natural conditions when geological strata containing pyrites are exposed to an oxidizing atmosphere or environment. One of the impacts of the mineralization zone where there is a mining process is the potential for the formation of acid mine drainage, especially in the Cibaliung gold mineralization area and its surroundings, Pandeglang Regency, Banten Province. Acid-forming sulfide minerals include pyrite (FeS<sub>2</sub>), headquarters (FeS<sub>2</sub>), picolites (FexSx), calcocytos (CuS), covellite (CuS), chalcocopyrite (CuFeS<sub>2</sub>), molybdenite (MoS), mulenite (NiS), chalcocytos (CuS), covellite (CuS), chalcocopyrite (CuFeS<sub>2</sub>), molybdenite (MoS), mulenite (NiS), chalcocytos (CuS), covellite (CuS), chalcocopyrite (CuFeS<sub>2</sub>), molybdenite (MoS), mulenite (NiS), galena (PbS) ) and sphalerite (ZnS). Of all these minerals, pyrite is the most dominant sulfide in acid formation.

Alkaline mine water (alkaline mine drainage) is mine water that has an acidity level (pH) of 6 or more, containing alkalinity but still containing dissolved metals that can produce acids. The quality of mine water, acid or alkali, depends on the presence or absence of acid mineral content (sulfides) and alkaline materials in the geological strata.

Acid water formation tends to be more intensive in mining areas. This can be prevented by avoiding exposure to sulfide-containing materials in the free air. Acid-forming sulfide minerals include pyrite (FeS<sub>2</sub>), headquarters (FeS<sub>2</sub>), picolites (FexSx), calcocytos (CuS), covellite (CuS), chalcocopyrite (CuFeS<sub>2</sub>), molybdenite (MoS), mulenite (NiS), chalcocytos (CuS), covellite (CuS), chalcocopyrite (CuFeS<sub>2</sub>), molybdenite (MoS), mulenite (NiS), chalcocytos (CuS), covellite (CuS), chalcocopyrite (CuFeS<sub>2</sub>), molybdenite (MoS), mulenite (NiS), galena (PbS) ) and sphalerite (ZnS). Of all these minerals, pyrite is the most dominant sulfide in acid formation. Formation of potential acidic water also occurs in tailings which are residues/processing residues containing sulfide minerals. The formation of acid mine drainage does not always develop in every sulfide-ore mining. In certain types of ore deposits, there are neutralizing agents which prevent the formation of acid mine drainage.

**Keywords:** Acid Mine Drainage, Epithermal gold, Sulphide Minerals, Source Rock, Cibaliung.

### 1. Introduction

AAT is formed as a result of oxidation of certain sulfide minerals contained in rocks, which react with oxygen in the air in an aqueous environment (Sayoga, 2007). In the initial stages of acid mine drainage is the presence of water in a green mining pit. As a result of these mining activities, can potentially produce acid mine drainage it is necessary to consider how the impact on the surrounding community.

Control of acid mine drainage is something that needs to be done during mining activities and after mining activities end, because Acid Mine Water (AAT) can cause a decrease in the quality of water, surface water, and groundwater, in addition, if flowed into the river will have an impact on the community stay along the river and will disturb the biota that lives on land as well as the biota in the waters. Acid Mine Water (AAT) can reduce water pH from public waters so that it will kill aquatic biota.

Alkaline mine water (alkaline mine drainage) is mine water that has an acidity level (pH) of 6 or more, containing alkalinity but still containing dissolved metals that can produce acids. The quality of mine water, acid or alkali, depends on the presence or absence of acid mineral content (sulfides) and alkaline materials in the geological strata. It generally contains a lot of sulfides and contains little alkaline material that tends to form acid mine water. Conversely, materials that contain a lot of alkalies, even though they contain sulfide material with a lot of concentration, often produce alkaline water (net machine water).

### 2. System and Characteristics of Epithermal Deposits

Epithermal ore deposits are deposits formed in near-surface hydrothermal environments that have relatively low temperatures and pressures and are associated with sub-aerial calc-alkali magmatism.



Among the types of mine water above there is the possibility of a transition type so that the appropriate data collection and analysis of metal concentrations,

water pH, and oxygen status need to be done to determine the type or characteristics of mine water.





Fig 3. Vein density Zone of gold mineralization in the area of Mangku Alam, Cimanggu, Pandeglang Banten

## 6. Reaction Formation of Acid Mine Water

### 6.1 Stages of the Establishment of Mine Acid Water


Stages of the reaction process of acid mine drainage.

#### Phase I:



The first reaction is the weathering reaction of pyrite (pyrite) accompanied by an oxidation process. Sulfur is oxidized to sulfate and iron-ferrous is released.


#### Phase II:



The second reaction is the conversion from ferrous to ferric materials which consume one mole of acidity. Slow reaction rate at pH <5 and abiotic conditions. Thiobacillus bacteria will accelerate the oxidation process. Thiobacillusferrooxidans is a gram-negative, acidophilic, autotrophic bacterium capable of using iron or various reduced sulfur compounds as an energy source (Vishniac in Wijayanti, 2002).


This microorganism has shown its great role in producing acid mine drainage. These microorganisms are also able to oxidize ferrous ions in ferrous form into ferric.

#### Phase III :



The third reaction is the hydrolysis of iron. Hydrolysis is a reaction that separates water molecules. Three moles of acidity is produced from this reaction. The formation of ferric hydroxide precipitation depends on pH, ie more at pH above 3.5.

#### Phase IV :



The fourth reaction is the continued oxidation of pyrite by ferric iron. This is a propagation reaction that takes place very quickly and will stop if the pyrite or ferric iron runs out. The oxidizing agent in this reaction is iron-ferry.

## 6.2 Characteristics of Differences in Rock Formation on the Potential or Absence of Acid Mine Water.

### 6.2.1 Based on Average Comparison Test

In each lithology can indicate the presence or absence of differences in the value of gold content (Au) as the main mineral, so it is necessary to know whether there are differences or similarities in lithology differences or there are other factors that influence it, a quantitative analysis based on the values of the levels of the main minerals and associated minerals based on drilling assay data.

Comparative test analysis of 3 average vectors that can be used to see whether there are differences in mineral content (Au, Ag, Cu, Pb, Zn, and As) in the Breccia to Andesitic (BRAN), Quartz Vein (VEIN) and Breccia mineralized zones Vein (VNBR) is a Multivariate Analysis of Variance (Manova). For this reason, it is necessary to conduct a multivariate normality test of the data.




Fig 4. Plot graph between Square Distance (dj2) and the quantile value of the Chi-Square distribution of each dj2 in the BRAN Zone.

Based on the picture above, the points on the scatterplot have a pattern that tends to be exponential, so it can be said that the BRAN Zone data is not normally distributed multivariate. Next, the correlation analysis between the value of Square Distance (dj2) and the quantile value of the Chi-Square distribution is hypothesized as follows.

H0: There is no relationship between the value of Square Distance (dj2) and the quantile value of the Chi-Square distribution (Data not normally distributed multivariate)

H1: There is a relationship between the value of Square Distance (dj2) and the quantile value of the Chi-Square distribution (multivariate normally distributed data)

Table 1. Multivariate Normality Test for the BRAN Zone Data

Mineralization Zone	R	p-value	Result
BRAN	0,923	0,000	<b>There is a Correlation</b>

Based on table 2. above it appears that the value of Square Distance (dj2) and the quantile value of the Chi-Square distribution have a strong relationship. So it can be concluded that the BRAN Zone data has a multivariate normal distribution.




Fig 5. Plot graph between Square Distance (dj2) and the quantile value of the Chi-Square distribution of each dj2 in the VEIN Zone

Next, the correlation analysis between the value of Square Distance (dj2) and the quantile value of the Chi-Square distribution is hypothesized as follows.

H0: There is no relationship between the value of Square Distance (dj2) and the quantile value of the Chi-Square distribution (Data not normally distributed multivariate)

H1: There is a relationship between the value of Square Distance (dj2) and the quantile value of the Chi-Square distribution (multivariate normally distributed data)

The test results (Table 2.) show that the value of Square Distance (dj2) and the quantile value of the Chi-Square distribution have a strong relationship, so it can be concluded that the VEIN Zone data has a multivariate normal distribution.

Table 2. VEIN Zone Data Multivariate Normality Test

Mineralization Zone	R	p-value	Result
VEIN	0,962	0,000	<b>There is a Correlation</b>




Fig 6. Plot graph between Square Distance (dj2) and the quantile value of the Chi-Square distribution of each dj2 in the VNBR Zone.

Next, the correlation analysis between the value of Square Distance (dj2) and the quantile value of the Chi-Square distribution is hypothesized as follows.

H0: There is no relationship between the value of Square Distance (dj2) and the quantile value of the Chi-Square distribution (Data not normally distributed multivariate)

H1: There is a relationship between the value of Square Distance (dj2) and the quantile value of the Chi-Square distribution (multivariate normally distributed data)

Based on (Table 5.4) above it can be seen that the value of Square Distance (dj2) and the quantile value of the Chi-Square distribution have a strong relationship, so it can be concluded that the VNBR Zone data has a multivariate normal distribution.

**Tabel 3. Data Zone of multivariate normality test VNBR**

Mineralization Zone	r	p-value	Result
VNBR	0,962	0,000	<b>There is a Correlation</b>

Based on the results of testing multivariate normality assumptions in each mineralized zone, it can be concluded that the assumption of multivariate normality is fulfilled. Therefore, the comparison test analysis of 3 average vectors used is Multivariate Analysis of Variance (Manova).

### 6.3 Manova Average Comparison Test

Hypothesis testing was carried out with a significance level ( $\alpha$ ) of 5% to determine the ratio of mineral content (Au, Ag, Cu, Pb, Zn, and As) in the Breccia to Andesitic (BRAN), Quartz Vein (VEIN) and Breccia Vein mineralized zones. (VNBR) with the hypothesis formulation as follows. H0: There were no significant (significant) differences between mineral levels (Au, Ag, Cu, Pb, Zn, and As) in the Breccia to Andesitic (BRAN), Quartz Vein (VEIN) and Breccia Vein (VNBR) mineralized zones.

H1: There is a significant (significant) difference between mineral content (Au, Ag, Cu, Pb, Zn, and As) in the Breccia to Andesitic (BRAN), Quartz Vein (VEIN), and Breccia Vein (VNBR) mineralized zones.

Test criteria: Reject H0 if p-value  $\leq$  0.05, accept in other cases.

Based on (Table 4.) it can be seen that by using the Wilks' Lambda test statistic obtained a p-value of 0,000 < 0.05, then H0 is rejected.

**Table 4. Comparison Test of Average Manova**

Zona	Statistical Test	F Count	p-value	Adjective
BRAN, VEIN, dan VNBR	<i>Wilks' Lambda</i>	6,212	0,000	<b>H<sub>0</sub>Reject</b>

So it can be concluded that there are significant (significant) differences between mineral content (Au, Ag, Cu, Pb, Zn, and As) in the Breccia to Andesitic (BRAN), Quartz Vein (VEIN), and Breccia Vein (VNBR) mineralized zones. Furthermore, to find out which

mineralized zones have different levels of minerals (Au, Ag, Cu, Pb, Zn, and As), further testing is done using T2 Hotelling.

The content of a precipitate indicates the presence of an element in the mineralized zone, where the presence of Cu, Pb, Zn and As in mineralization which could be the potential for acid mine formation in a sulfide mineralized zone.

### 6.4 Conclusion

There is a significant (significant) difference between mineral levels (Au, Ag, Cu, Pb, Zn, and As) in the Breccia to Andesitic (BRAN), Quartz Vein (VEIN) and Breccia Vein (VNBR) mineralized zones. Furthermore, to find out which mineralized zones have different levels of minerals (Au, Ag, Cu, Pb, Zn, and As). The content of a precipitate indicates the presence of an element in the mineralized zone, where the presence of Cu, Pb, Zn and As in mineralization which could be the potential for acid mine formation in a sulfide mineralized zone.

Mineralization which could be a potential formation of acid mine drainage in a zone of sulfide mineralization. Many parameters inside rocks and host rock to make occurrence about Mine Water Drainage. The statistical method can be to get a solution to a correlation between parameter in rocks and ore.

### Acknowledgments

Thank you to all those who have supported this research on Acid Mine Water. Especially to PT Cibaliung Sumberdaya, we are very grateful to have been given the ease and fluency in conducting research. To LPPM Unisba who has given permission and set us as one of the winners of the grant so that we can conduct research and all parties that we cannot mention one by one

### References

- DudiNasrudin Usman, Nana Sulaksana, FebriHirnawan, Iyan Haryanto, &NurdinSyaefulBahri (2017). Analysis of the Solid Structure of Late Miocene Gold Mineralization Zones. Snapp Proceedings: Science, Technology, 7 (1), 392-397. LPPM - Bandung Islamic University.
- DudiNasrudin Usman, Nana Sulaksana, FebriHirnawan, Iyan Haryanto. 2018. Gold-Silver Mineralization in the Neo-Tectonism of Honje Formation and Cipacar Formation, in Cibaliung Block, Banten Province. Journal of Geoscience, Engineering, Environment and Technology. Vol. 3 No. 4 2018. DOI: 10.24273 / jgeet.2018.3.4.1852
- Sayoga, R. G. 2007. Mine Water Management: Important Aspects in Mining with Environmental Insights. Scientific Speech, assembly of ITB Professors. Department of Mining Engineering ITB.
- Wijanyanti, Retno. 2002. Oxidation of Ferrous Ion in Mine Water with Iron-Oxidizing Bacteria. Mineral Technology Research and Development Center, Bandung



© 2016 Journal of Geoscience, Engineering, Environment and Technology. All rights reserved. This is an open access article distributed under the terms of the CC BY-SA License (<http://creativecommons.org/licenses/by-sa/4.0/>).

---