

## Production of Exopolysaccharides by Strains of *Streptococcus thermophilus*

UMI PURWANDARI<sup>1\*</sup> AND TODOR VASILJEVIC<sup>2</sup>

<sup>1</sup>Department of Agroindustrial Technology, Faculty of Agriculture, Universitas Trunojoyo Madura, Indonesia, PO Box 2, Kamal, Madura 69162, Indonesia; <sup>2</sup>School of Molecular Sciences, Faculty of Health, Engineering and Science, Institute for Sustainability and Innovation, Victoria University, Werribee Campus, PO Box 14428, Melbourne City MC, Victoria 8001, Australia.

Two strains of *Streptococcus thermophilus* producing capsular and capsular-ropy exopolysaccharides (EPS) were examined for their growth and EPS production in M17 medium supplemented with glucose, galactose or lactose and incubated at 30, 37 or 42°C for 24 hours. Growth parameters (viable cells, OD, lactate production, pH) and EPS production were determined. Flow behavior of the EPS dispersions was assessed as a function of concentration and temperature. Culture growth during incubation was affected by types of sugar, temperature and time. Growth was enhanced by glucose, lactose and higher incubation temperature. EPS concentration in the medium was greater in the presence of glucose and galactose. Despite the restricted growth conditions, the capsular strain produced comparable levels of EPS to the capsular-ropy strain even under sub-optimal incubation temperature

Key words: *Streptococcus thermophilus*, exopolysaccharides, growth conditions

Considerable attention has been given to bacterial exopolysaccharides (EPS) in foods arising their ability to provide potential health benefits to consumers (Looijesteijn *et al.* 2001) and their application as thickening agents in processed products (Cerning 1990). Although certain bacterial species can produce considerable amounts of EPS, yoghurts prepared with strains *Streptococcus thermophilus* resulted in less than 0.1% EPS in the final product (Cerning 1990). Nevertheless, EPS play an important role in the development of yoghurt texture, with the type of EPS exerting a greater effect than their concentration (Vaningelgem *et al.* 2004). In general, two strain-dependent types of EPS (Mozzi *et al.* 2006) have been frequently assessed for their effects on yoghurt texture, namely 'ropy' and 'capsular'. Certain bacterial strains may even produce a mix of these two types in various proportions (Zisu and Shah 2003; Zisu and Shah 2005). Furthermore, combinations of the two types of EPS producing cultures improved not only the total EPS production but also yoghurt texture (Marshall and Rawson 1999).

Numerous studies have been conducted in order to enhance our understanding of the factors governing EPS production and the mechanisms by which they affect the yoghurt texture. Most of the EPS produced during yoghurt fermentation are heteropolysaccharides, and their production may be unpredictable due to a plasmid-related instability (Boels *et al.* 2001). EPS production, thickening properties, molecular mass and structural conformation are greatly affected by environmental factors (Ruas-Madiedo *et al.* 2005). In some species, these factors only affected EPS yield, because their monosaccharide composition remained unchanged (Looijesteijn and Hugenholtz 1999). Strain selection, temperature, pH and growth stage (Aslim *et al.* 2005), nitrogen source (Degeest *et al.* 2002), and metabolizable sugars (Mozzi *et al.* 2003) are among the

factors influencing EPS production. EPS production by some *S. thermophilus* strains was coupled to growth (Vin *et al.* 2005). However, in some mesophilic bacterial strains, EPS were mainly produced at sub-optimal growth conditions (Cerning 1990). Depending on the strain and growth conditions, maximum yield of EPS may be achieved in the exponential (Duenas *et al.* 2003) or stationary growth phase (Gancel and Novel 1994). At the end of growth phase, there are some indications that EPS undergoes undesirable enzymatic degradation (Pham *et al.* 2000; Degeest *et al.* 2002). The type of carbon source apparently governs the total amount of EPS produced with a clear strain-dependence (Mozzi *et al.* 2001). For example, *S. thermophilus* LY03 produced more EPS with lactose than with glucose (Degeest and de Vuyst 2000a). Nitrogen compounds in the growth medium are necessary during formation of sugar nucleotides, essential precursors for EPS assembly. However, the type of nitrogen source appears to be less important in EPS production than is the carbon source (Duenas *et al.* 2003).

Although several types of EPS have been reported to influence yoghurt texture, and some mechanisms on their interactions with milk proteins has been proposed, there is a lack of information on the rheological properties of EPS produced by yoghurt cultures. Such information can be useful for predicting the possible interaction of EPS with milk component(s) to influence the texture of yoghurt. Therefore, this study aimed to examine EPS production by two strains of *S. thermophilus* which produced capsular-ropy and capsular EPS. We reported the effect of fermentation conditions including temperature, time, and carbon source on its production.

### MATERIALS AND METHODS.

**Bacterial Strains.** The two strains of *S. thermophilus* examined in the study were kindly provided by Australian Starter Culture Research Centre (Werribee, Australia). *S. thermophilus* ST 1275 produces mainly ropy with a smaller portion of capsular EPS, while ST 285 produces only

\*Corresponding author: Tel/Fax: +62-31-3013234,  
E-mail: umipurwandari@trunojoyo.ac.id

capsular EPS (Zisu and Shah 2003; 2005). The frozen (-80°C) cultures in 300 mL L<sup>-1</sup> glycerol were activated by growing them twice in M17 medium (Amyl Media, Merck Pty Ltd., Kilsyth, Victoria, Australia) at 37°C for at least 24 hours.

**Medium.** The medium used in these experiments was basal M17 medium supplemented with galactose, glucose or lactose at 20 g L<sup>-1</sup>. The cultures were inoculated into 30 mL of the sterile medium in 50 mL Falcon tubes (Falcon, Blue Max, Becton Dickinson and Company, Franklin Lakes, N.J., USA) and incubated at 30, 37 or 42°C for 24 hours, during which samples were periodically collected for analysis of all growth parameters and determination of EPS concentration.

**Bacterial Counts.** Determination of viable cell counts was carried out by serial dilutions in sterile 1 g L<sup>-1</sup> peptone water, and incubation at the corresponding temperature for 24 hours to obtain sufficient cell growth. Cell concentration was additionally determined based on optical density at 650 nm.

**EPS Determination.** EPS assessed in the rheological studies were extracted from the culture grown in M17 medium by precipitation with ethanol (van Geel-Schutten *et al.* 1998). Initially, fermentation was stopped by adding 20 g L<sup>-1</sup> TCA and keeping in a cool room (4°C) overnight. This was followed by protein removal by centrifugation at 11 000 × g and 4°C for 10 min (Model J2-HS, Beckman, Fullerton, California, USA). The supernatant was collected and mixed with two volumes of ethanol. The mixture was allowed to stand for twelve hours for complete precipitation of EPS. This procedure was repeated twice, after which the EPS was freeze-dried (Dynavac freeze drier, Dynavac Eng. Pty. Ltd., Melbourne, Australia). The

concentration of EPS was assessed using the phenolsulphuric acid method (Dubois *et al.* 1956). Lactic acid concentration was determined by an HPLC method (Donkor *et al.* 2005). The method uses dilute 0.01 mol/L H<sub>2</sub>SO<sub>4</sub> as a mobile phase with flow rate of 0.6 mL min<sup>-1</sup>, and 300 x 7.8 mm ion-exchange Aminex HPX-87H column (Bio-Rad Laboratories, Australia) coupled to a UV detector (Varian Analytical Instruments, Walnut Creek, CA, USA) at 220 nm. The column temperature was maintained at 65°C.

**Statistical Analysis.** A randomized split plot in time block design was applied to the design of the fermentation experiments. This was set up with four main effects: strain (two levels), sugar types (three levels), temperatures (three levels), and fermentation time (six levels: 0, 3, 6, 9, 12, 24 hours). This design was replicated twice with at least two sub samplings. Results were analyzed using a General Linear Model procedure. The level of significance was set at P=0.05. The best fit correlational analysis for rheological data was carried out using the Rheoplus software (v2.81, Anton Paar).

## RESULTS

**Bacterial Counts.** Viable cell counts were significantly (P<0.05) affected by all factors: strain, temperature, type of sugar, fermentation time and their interactions. The capsular strain consistently showed slower growth compared to the capsular-ropy culture, regardless of the growth conditions. The maximum cell numbers of the capsular strain were 2.2x10<sup>8</sup> CFU mL<sup>-1</sup>, achieved in glucose-M17 medium at 42°C. In contrast, the capsular-ropy culture grew to 1.6x10<sup>10</sup> CFU mL<sup>-1</sup> in lactose-M17 medium at 37°C (Fig 1a, b). Higher temperatures (37 and 42 °C) resulted in greater cell growth than at 30°C for both strains and all types of sugar.




Fig 1 Viable cell counts of capsular-ropy (a) and capsular (b) strain of *Streptococcus thermophilus* cultivated in a M17 medium supplemented with glucose (solid line), galactose (dashed line) or lactose (dotted line) at 30°C (□), 37°C (○) or 42°C (▽) for 24 h.

Table 1 Cell growth parameters of the capsular-ropy and capsular strain of *Streptococcus thermophilus* as affected by temperature and sugar source during growth phase

Strain	Temperature, (°C)	Time, (hour)	Type of sugar	pH	Lactic acid concentration, (mg mL <sup>-1</sup> )	OD*
Capsular-ropy	30	8	galactose	6.9 <sup>EB</sup>	0.53 <sup>2A</sup>	0.07 <sup>CA</sup>
			lactose	5.9 <sup>BB</sup>	0.67 <sup>BA</sup>	0.86 <sup>BA</sup>
			glucose	6.3 <sup>BB</sup>	0.87 <sup>CA</sup>	0.72 <sup>BA</sup>
		24	galactose	6.9 <sup>CA</sup>	0.54 <sup>2A</sup>	0.09 <sup>BA</sup>
			lactose	5.9 <sup>BB</sup>	0.66 <sup>BB</sup>	0.95 <sup>BA</sup>
			glucose	5.5 <sup>BB</sup>	0.86 <sup>CA</sup>	1.73 <sup>CA</sup>
	37	8	galactose	6.8 <sup>CA</sup>	0.55 <sup>2AB</sup>	0.16 <sup>AA</sup>
			lactose	5.8 <sup>BB</sup>	0.68 <sup>BA</sup>	1.23 <sup>BB</sup>
			glucose	5.1 <sup>AA</sup>	0.88 <sup>CA</sup>	2.52 <sup>CB</sup>
		24	galactose	6.7 <sup>CA</sup>	0.68 <sup>BC</sup>	0.20 <sup>BA</sup>
			lactose	5.8 <sup>BB</sup>	0.57 <sup>2A</sup>	1.08 <sup>BA</sup>
			glucose	5.0 <sup>AA</sup>	0.89 <sup>CA</sup>	2.08 <sup>CB</sup>
42	8	galactose	6.6 <sup>BB</sup>	0.58 <sup>BB</sup>	0.29 <sup>BA</sup>	
		lactose	5.3 <sup>2A</sup>	0.77 <sup>BB</sup>	1.75 <sup>BC</sup>	
		glucose	5.1 <sup>AA</sup>	0.89 <sup>CA</sup>	2.53 <sup>CB</sup>	
	24	galactose	6.5 <sup>BA</sup>	0.60 <sup>BB</sup>	0.28 <sup>BA</sup>	
		lactose	5.3 <sup>2A</sup>	0.74 <sup>BC</sup>	1.67 <sup>BB</sup>	
		glucose	5.1 <sup>AA</sup>	0.89 <sup>CA</sup>	2.78 <sup>CB</sup>	
Capsular	30	8	galactose	6.9 <sup>CA</sup>	0.55 <sup>2A</sup>	0.24 <sup>CA</sup>
			Lactose	5.9 <sup>BA</sup>	0.65 <sup>BA</sup>	0.94 <sup>BB</sup>
			glucose	5.3 <sup>2A</sup>	0.84 <sup>CB</sup>	2.12 <sup>BB</sup>
		24	Galactose	6.8 <sup>CA</sup>	0.56 <sup>2A</sup>	0.27 <sup>CB</sup>
			lactose	5.9 <sup>BA</sup>	0.66 <sup>BA</sup>	1.01 <sup>AB</sup>
			glucose	5.3 <sup>2A</sup>	0.83 <sup>CB</sup>	2.11 <sup>CB</sup>
	37	8	galactose	6.8 <sup>CA</sup>	0.54 <sup>2A</sup>	0.20 <sup>BA</sup>
			lactose	5.9 <sup>BA</sup>	0.64 <sup>BA</sup>	0.57 <sup>BA</sup>
			glucose	5.4 <sup>2A</sup>	0.84 <sup>CB</sup>	1.69 <sup>CB</sup>
		24	galactose	6.6 <sup>CA</sup>	0.57 <sup>2A</sup>	0.14 <sup>AA</sup>
			lactose	5.9 <sup>BA</sup>	0.66 <sup>BA</sup>	0.66 <sup>BB</sup>
			glucose	5.4 <sup>2A</sup>	0.84 <sup>CB</sup>	1.51 <sup>CA</sup>
42	8	galactose	6.9 <sup>CA</sup>	0.59 <sup>2B</sup>	0.20 <sup>BA</sup>	
		lactose	5.9 <sup>BA</sup>	0.66 <sup>BA</sup>	0.57 <sup>BA</sup>	
		glucose	5.4 <sup>2A</sup>	0.76 <sup>CA</sup>	1.59 <sup>CA</sup>	
	24	galactose	6.9 <sup>CA</sup>	0.57 <sup>2A</sup>	0.16 <sup>AA</sup>	
		lactose	6.1 <sup>BB</sup>	0.62 <sup>BA</sup>	0.44 <sup>BA</sup>	
		glucose	5.4 <sup>2A</sup>	0.78 <sup>CA</sup>	1.60 <sup>CA</sup>	
SEM**				0.101	0.020	0.108

\*Different small caps in a column denote significant (P<0.05) difference among means of the same culture, temperature and time but different sugar types. The different capital letters depict significant (P<0.05) difference among means of the same culture, sugar type and time but different temperature. \*\*SEM, standard error of means, with P=0.05.

M17 medium containing glucose or lactose was more growth supportive than that with galactose for both strains and in all temperatures.

Positive effects of higher fermentation temperature, glucose or lactose on cell growth was also confirmed by other growth parameters such as optical density, pH and lactic acid production (Table 1). Statistically, OD readings were only influenced by the type of sugar and its interactions with other factors (P<0.05). Fermentation time and the type of strain individually had no effect on cell growth as measured by OD, but their interactions did (P<0.05). Sugar type and its interaction with other factors significantly (P<0.05) affected the final pH of the medium. Lactic acid concentration was similarly influenced by the factors examined. The types of sugar substantially (P<0.05) influenced lactate concentration, where glucose produced most lactic acid (~0.80-0.87±0.020 mg mL<sup>-1</sup>), whereas galactose produced the lowest levels ~0.55-0.62±0.02 mg mL<sup>-1</sup>. In most cases, from the 8th hour onwards, lactate concentration was either increased or remained unchanged.

**EPS Concentration.** The EPS production was only affected by the type of sugar and the fermentation time, and their interactions with other factors (P<0.05). The amount of EPS produced by the two strains was not significantly different (P>0.05). Maximum EPS concentration produced by the capsular-ropy strain was 441.0±36.2 mg L<sup>-1</sup> in galactose-M17 at 42°C after 12 hours. Similarly, the capsular strain produced 563.0±36.2 mg L<sup>-1</sup> with glucose at 30°C after 12 hours (Fig 2a, b). EPS production of the capsular strain at all temperatures was relatively high and the difference among them was negligible. Notably in our study, EPS production was significantly (P<0.05) affected by incubation time though not statistically significant (P>0.05), higher temperature tended to enhance EPS production by the capsular-ropy strain in all types of sugar (Fig 2a,b). On the other hand, the capsular strain appeared to prefer lower




Fig 2 Exopolysaccharides (EPS) concentration, expressed as glucose equivalent, produced by the capsular-ropy (a) and the capsular (b) strain of *Streptococcus thermophilus* grown in a M17 medium supplemented with glucose (solid line), galactose (dashed line) or lactose (dotted line), at 30°C (□), 37°C (○) or 42°C (▽) for 24 h.



temperature (30°C) and produced more EPS when grown on glucose or lactose (Fig 2a, b). M17 medium containing galactose poorly supported cell growth, however, the EPS concentration in this medium was relatively high. In medium with lactose at all temperatures, and glucose or galactose at low temperature (30°C), maximum EPS concentration was reached after 8 hours. Afterwards, the EPS concentration was often slightly reduced towards the end of fermentation. The pattern of EPS production over time by the capsular strain was somewhat similar to that of the capsular-ropy counterpart, except that EPS concentration in galactose medium at low temperature (30°C) remained stationary after 4 hours. Moreover, EPS concentrations in lactose medium at low temperatures (30 and 37°C) were low, and did not show any considerable change until the end of fermentation.

### DISCUSSION

**Bacterial Counts.** Our results are in agreement with previous work showing comparable cell numbers of *S. thermophilus* grown in lactose- and glucose-containing medium (Degeest and de Vuyst 2000b). In contrast to these findings, van den Bogaard *et al.* (2000) found that *S. thermophilus* grew poorly on glucose (Poolman 2002). Growth of the *S. thermophilus* strain in the glucose based medium requires the activity of phosphoglucomutase whose activity may be suppressed in the presence of lactose (Levander and Radstrom 2001). Degeest *et al.* (2000b) also showed that galactose did not support the growth of a *S. thermophilus* strain.

**EPS Concentration.** Our experiment showed that both strains produced considerable amounts of EPS in M17 medium, even higher than as reported by others (Aslim *et al.* 2006). As in this experiment, no difference in EPS production of both ropy and non-ropy strains was found by Mozzi *et al.* (2006). Most of *S. thermophilus* strains possess a Leloir system for EPS synthesis (Mora *et al.* 2002). The three sugars examined in our study supported EPS production, as has reported elsewhere (Chervaux *et al.* 2000), with galactose frequently related to EPS production (Mozzi *et al.* 2001). Our result was in contrast to previous report that glucose poorly supported EPS production compared to lactose (Degeest *et al.* 2000a), but similar to a more recent work (Shene *et al.* 2008) that EPS production of *S. thermophilus* strain was higher on glucose-containing medium. The growth-associated nature of EPS production of *S. thermophilus* strains has been reported previously, with the EPS produced mainly during logarithmic phase (Ruas-Madiedo *et al.* 2005; Shene *et al.* 2008). The maximum production of EPS seemingly varies with strains and is not always linear with temperature of fermentation, as also shown by previous report (Shene *et al.* 2008).

Galactose was a predominant (Goh *et al.* 2005) and a consistent (Goh *et al.* 2005; Mozzi *et al.* 2006) primary unit in the structure of the EPS backbone in several *S. thermophilus* strains, although most of them did not utilize galactose (Mora *et al.* 2002) or metabolized galactose only at the end of growth phase (de Vin *et al.* 2005). However,

several reports found discrepancies between theoretical values and actual galactose concentration in the medium, suggesting a flux to other metabolites including lactic acid or even assimilation of galactose into EPS in the Gal<sup>-</sup> strains of *S. thermophilus* (de Vin *et al.* 2005). Galactose appeared to contribute to EPS anabolism rather than the cell growth (Mozzi *et al.* 2001). The availability of galactose might be initiated by the activation of galactose symporter in the absence of lactose (de Vin *et al.* 2005) or low lactose/galactose ratio (Poolman 2002). Moreover, galactose catabolism is also possible by the activation of some enzymes in Leloir pathway such as galactokinase in a lactose-depleted environment (de Vin *et al.* 2005). The EPS concentration in the medium was reduced at the end of the growth phase. This could be related to shortage of ATP required for EPS polymerization (Welman *et al.* 2006). Moreover, cell lysis (Mozzi *et al.* 2003) could lead to enzymatic degradation of EPS (Pham *et al.* 2000) towards the late stages of the growth.

### REFERENCES

- Aslim B, Beyatli Y, Yuksekdogan ZN. 2006. Productions and monomer compositions of exopolysaccharides by *Lactobacillus delbrueckii* subsp *bulgaricus* and *Streptococcus thermophilus* strains isolated from traditional home-made yoghurts and raw milk. *Int J Food Sci Tech* 41:973-9.
- Aslim B, Yuksekdogan ZN, Beyatli Y, Mercan N. 2005. Exopolysaccharide production by *Lactobacillus delbrueckii* subsp *bulgaricus* and *Streptococcus thermophilus* strains under different growth conditions. *World J Microbiol Biotechnol* 21:673-7.
- Boels IC, Kranenburg R van, Hugenholtz J, Kleerebezem M, Vos WM de. 2001. Sugar catabolism and its impact on the biosynthesis and engineering of exopolysaccharide production in lactic acid bacteria. *Int Dairy J* 11:723-32.
- Cerning J. 1990. Exocellular polysaccharides produced by lactic acid bacteria. *FEMS Microbiol Lett* 87: 113-30.
- Chervaux C, Ehrlich SD, Maguin E. 2000. Physiological study of *Lactobacillus delbrueckii* subsp *bulgaricus* strains in a novel chemically defined medium. *Appl Environ Microbiol* 66:5306-11.
- de Vin F, Radstrom P, Herman L, Vuyst L de. 2005. Molecular and biochemical analysis of the galactose phenotype of dairy *Streptococcus thermophilus* strains reveals four different fermentation profiles. *Appl Environ Microbiol* 71:3659-67.
- Degeest B, de Vuyst L. 2000a. Correlation of activities of the enzymes  $\alpha$ -phosphoglucomutase, UDP-galactose 4-epimerase, and UDP-glucose pyrophosphorylase with exopolysaccharide biosynthesis by *Streptococcus thermophilus* LY03. *Appl Environ Microbiol* 66:3519-27.
- Degeest B, de Vuyst L. 2000b. Correlation of activities of the enzymes  $\alpha$ -Phosphoglucomutase, UDP-galactose 4-epimerase, and UDP-glucose pyrophosphorylase with exopolysaccharide biosynthesis by *Streptococcus thermophilus* LY03. *Appl Environ Microbiol* 66:3519-27.
- Degeest B, Mozzi F, Vuyst L de. 2002. Effect of medium composition and temperature and pH changes on exopolysaccharide yields and stability during *Streptococcus thermophilus* LY03 fermentations. *Int J Food Microbiol* 79:161-74.
- Donkor ON, Henriksson A, Vasiljevic T, Shah NP. 2005. Probiotic strains as starter cultures improve angiotensin-converting enzyme inhibitory activity in soy yogurt. *J Food Sci* 70:M375-81.
- Dubois M, Gilles KA, Hamilton K, Rebers PA, Smith F. 1956. Colorimetric method for the determination of sugars and related substances. *Anal Chem* 28:350-6.
- Duenas M, Munduate A, Perea A, Irastorza A. 2003. Exopolysaccharide production by *Pediococcus damnosus* 2.6 in a semidefined medium under different growth conditions. *Int J Food Microbiol* 87:113-20.
- Gancel F, Novel G. 1994. Exopolysaccharide production by *Streptococcus salivarius* ssp *thermophilus* cultures: two distinct modes of polymer production and degradation among clonal variants. *J Dairy Sci* 77:689-95.
- Goh KKT, Haisman RD, Singh H. 2005. Examination of exopolysaccharide produced by *Lactobacillus delbrueckii* subsp *bulgaricus* using confocal laser scanning and scanning electron microscopy techniques. *J Food Sci* 70:M224-9.
- Levander F, Radstrom P. 2001. Requirement for phosphoglucomutase in exopolysaccharide biosynthesis in glucose- and lactose-utilizing *Streptococcus thermophilus*. *J Appl Environ Microbiol* 67:2734-8.
- Looijesteijn PJ, Hugenholtz J. 1999. Uncoupling of growth and exopolysaccharide production by *Lactococcus lactis* subsp. *cremoris* Nizo B40 and optimization of its synthesis. *J Biosci Bioeng* 88:178-82.
- Looijesteijn PJ, Trapet L, Vries E de, Abee T, Hugenholtz J. 2001. Physiological function of exopolysaccharides produced by *Lactococcus lactis*. *Int J Food Microbiol* 64:71-80.
- Marshall VM, Rawson HL. 1999. Effects of exopolysaccharide-producing strains of thermophilic lactic acid bacteria on the texture of stirred yoghurt. *Int J Food Sci Technol* 34:137-43.
- Mora D, Fortina MG, Parini C, Ricci G, Gatti M, Giraffa G, Manachini PL. 2002. Genetic diversity and technological properties of *Streptococcus thermophilus* strains isolated from dairy products. *J Appl Microbiol* 93:278-87.
- Mozzi F, Giori GS de, Font de Valdez G. 2003. UDP-galactose 4-epimerase: A key enzyme in exopolysaccharide formation of *Lactobacillus casei* CRL 87 in controlled pH batch cultures. *J Appl Microbiol* 94:175-83.
- Mozzi F, Rollán G, Giori GS de, G, Font de Valdez G. 2001. Effect of galactose and glucose on the exopolysaccharide production and the activities of biosynthetic enzymes in *Lactobacillus casei* CRL 87. *J Appl Microbiol* 91:160-7.
- Mozzi F, Vaningelgem F, Hebert EM, Van Der Meulen R, Moreno MRF, Font de Valdez G, Vuyst de L. 2006. Diversity of heteropolysaccharide-producing lactic acid bacterium strains and their biopolymers. *Appl Environ Microbiol* 72:4431-5.
- Pham PL, Dupont I, Roy D, Lapointe G, Cerning J. 2000. Production of exopolysaccharide by *Lactobacillus rhamnosus* R and analysis of its enzymatic degradation during prolonged fermentation. *Appl Environ Microbiol* 66:2302-10.
- Poolman B. 2002. Transporters and their roles in lab cell physiology. *nt J Gen Mol Microbiol* 82:147-64.
- Ruas-Madiedo P, Altling AC, Zoon P. 2005. Effect of exopolysaccharides and proteolytic activity of *Lactococcus lactis* subsp *cremoris* strains on the viscosity and structure of fermented milks. *Int Dairy J* 15:155-64.
- Shene C, Canquil N Bravo S, Rubilar M. 2008. Production of the exopolysaccharides by *Streptococcus thermophilus*: Effects of growth conditions on fermentation kinetics and intrinsic viscosity. *Int J Food Microbiol* 124:279-84.
- Van den Bogaard PTC, Kleerebezem M, Kuipers OP, Vos WM. 2000. Control of lactose transport,  $\beta$ -galactosidase activity, and glycolysis by CCPA in *Streptococcus thermophilus*: evidence for carbon catabolite repression by a non-phosphoenolpyruvate-dependent phosphotransferase system sugar. *J Bacteriol* 182:5982-9.
- Van Geel-Schutten GH, Flesch F, Brink B, Smith MR, Dijkhuizen L. 1998. Screening and characterization of *Lactobacillus* strains producing large amounts of exopolysaccharides. *Appl Microbiol Biotechnol* 50:697-703.
- Vaningelgem F, Zamfir M, Mozzi F, Adriani T, Vancanneyt M, Swings J, Vuyst L de. 2004. Biodiversity of exopolysaccharides produced by *Streptococcus thermophilus* strains is reflected in their production and their molecular and functional characteristics. *Appl Environ Microbiol* 70:900-12.
- Welman AD, Maddox IS, Archer RH. 2006. Metabolism associated with raised metabolic flux to sugar nucleotide precursors of exopolysaccharides in *Lactobacillus delbrueckii* subsp *bulgaricus*. *J Indust Microbiol Biotechnol* 33:391-400.
- Zisu B, Shah NP. 2003. Effects of pH, temperature, supplementation with whey protein concentrate, and adjunct cultures on the production of exopolysaccharides by *Streptococcus thermophilus* 1275. *J Dairy Sci* 86:3405-15.
- Zisu B, Shah NP. 2005. Textural and functional changes in low-fat mozzarella cheeses in relation to proteolysis and microstructure as influenced by the use of fat replacers, pre-acidification and EPS starter. *Int Dairy J* 15:957-72.