

## On a Property of the $n$ -dimensional Cube

Rafayel Kamalian<sup>1</sup>, Arpine Khachatryan<sup>2</sup>

<sup>1</sup> Institute for Informatics and Automation Problems,  
National Academy of Sciences of the Republic of Armenia, 0014, Armenia,  
email: rrkamalian@yahoo.com

<sup>2</sup> Ijevan Branch of Yerevan State University, 4001, Armenia,  
email: khachatryanarpine@gmail.com

We show that in any subset of vertices of the  $n$ -dimensional cube which contains at least  $2^{n-1} + 1$  vertices ( $n \geq 4$ ), there are four vertices that induce a claw, or there are eight vertices that induce the cycle of length eight.

We consider finite graphs  $G = (V, E)$  with vertex set  $V$  and edge set  $E$ . The graphs contain no multiple edges or loops. The  $n$ -dimensional cube is denoted by  $Q_n$ , and a claw is the complete bipartite graph  $K_{1,3}$ . Moreover, the vertex of a degree three in a claw is called a claw-center. Non-defined terms and concepts can be found in [1].

The main result of the paper is the following:

**Theorem 1.** *Let  $n \geq 4$  and let  $V' \subseteq V(Q_n)$ . If  $|V'| \geq 2^{n-1} + 1$ , then at least one of the following two conditions holds:*

- (a) *there are four vertices in  $V'$  that induce a claw;*
- (b) *there are eight vertices in  $V'$  that induce a simple cycle.*

**Proof.** Our proof is by induction on  $n$ . Suppose that  $n = 4$ . Clearly, without loss of generality, we can assume that  $|V'| = 9$ . Consider the following partition of the vertices of  $Q_4$ :

$$V_1 = \{(0, \alpha_2, \alpha_3, \alpha_4) : \alpha_i \in \{0, 1\}, 2 \leq i \leq 4\}, V_2 = \{(1, \alpha_2, \alpha_3, \alpha_4) : \alpha_i \in \{0, 1\}, 2 \leq i \leq 4\}.$$

Clearly, the subgraphs of  $Q_4$  induced by  $V_1$  and  $V_2$  are isomorphic to  $Q_3$ . Define:

$$V'_1 = V_1 \cap V', V'_2 = V_2 \cap V'.$$

We shall assume that  $|V'_1| \geq |V'_2|$ . We shall complete the proof of the base of induction by considering the following cases:

- Case 1:  $|V'_1| = 8$  and  $|V'_2| = 1$ . Clearly, any vertex from  $V'_1$  is a claw-center.
- Case 2:  $|V'_1| = 7$  and  $|V'_2| = 2$ . It is not hard to see that  $V'_1$  contains a claw-center.
- Case 3:  $|V'_1| = 6$  and  $|V'_2| = 3$ . Again, it is a matter of direct verification that  $V'$  contains a claw-center.
- Case 4:  $|V'_1| = 5$  and  $|V'_2| = 4$ . Consider the subgraph  $G_1$  of  $Q_4$  induced by  $V'_1$ . Clearly, if  $G_1$  contains a vertex of a degree three, then this vertex is a claw-center. Therefore, without

loss of generality, we can assume that any vertex in  $G_1$  has a degree at most two. It is not hard to see that this implies that  $G_1$  contains no isolated vertex. Moreover, since  $|V'_1| = 5$ , we can conclude that  $G_1$  is a connected graph, and, consequently, it is the path of length four.

Now, let  $a_1, a_2, a_3$  be the internal vertices of  $G_1$ , and let  $b_1, b_2$  be the end-vertices of  $G_1$ . Clearly, we can assume that neither of  $a_1, a_2, a_3$  has a neighbour in  $V'_2$ . Since  $|V_2| = 8$  and  $|V'_2| = 4$ , we have that there are five possibilities for  $V'_2$ . We invite the reader to check that in four of these cases one can find a claw-center in  $V'_2$ , and in the final case  $V'$  has a vertex  $z$  such that  $V' \setminus \{z\}$  induces a simple cycle.

Now, let us assume that the statement is true for  $n - 1$ , and a subset  $V'$  of the vertices of  $Q_n$  satisfies the inequality  $|V'| \geq 2^{n-1} + 1$ . Consider the following partition of the vertices of  $Q_n$ :

$$V_1 = \{(0, \alpha_2, \dots, \alpha_n) : \alpha_i \in \{0, 1\}, 2 \leq i \leq n\}, V_2 = \{(1, \alpha_2, \dots, \alpha_n) : \alpha_i \in \{0, 1\}, 2 \leq i \leq n\}.$$

Clearly, the subgraphs of  $Q_n$  induced by  $V_1$  and  $V_2$  are isomorphic to  $Q_{n-1}$ . Moreover, it is not hard to see that at least one of the following two inequalities is true:  $|V_1 \cap V'| \geq 2^{n-2} + 1$  and  $|V_2 \cap V'| \geq 2^{n-2} + 1$ . Thus the proof follows from the induction hypothesis.

For the case of  $n = 3$  we have:

**Proposition 1.** *Let  $V' \subseteq V(Q_3)$  and let  $|V'| \geq 6$ . Then at least one of the following two conditions holds:*

- *there are four vertices in  $V'$  that induce a claw;*
- *there are six vertices in  $V'$  that induce a simple cycle.*

**Acknowledgement.** We would like to thank Zhora Nikoghosyan and Vahan Mkrtchyan for their attention to this work.

## References

- [1] West D.B. *Introduction to Graph Theory*. Prentice-Hall, New Jersey, 1996.