

A REVIEW OF *Ceratocystis sensu stricto* WITH SPECIAL REFERENCE TO THE SPECIES COMPLEXES *C. coerulescens* AND *C. fimbriata*

Mauricio Marín Montoya¹ y Michael J. Wingfield²

ABSTRACT

The genus *Ceratocystis sensu stricto* Ellis & Halst. includes many economically important plant pathogens of both angiosperms and gymnosperms, worldwide. Diseases caused by these fungi include vascular wilts, sap stains on logs and lumber, stem cankers and rots of roots, stems and fruits. Most *Ceratocystis* species are well adapted to dispersal by insects. Many species produce volatile metabolites with fruity odors that attract a wide range of insects to infected tissues. However, the degree of this association between insects and *Ceratocystis* species is highly variable, ranging from a mutualistic relationship such as that between *C. polonica*, *C. laricicola* and *C. rufipennis* and their specific bark beetle vectors, to non-specific associations, as are found in the case of *C. paradoxa*, *C. fagacearum* and *C. fimbriata*, with nitidulid beetles, flies and ambrosia beetles. The high degree of morphological similarity found amongst these fungi probably results from convergent evolution driven by selection for insect dispersal, and has made it difficult to clearly define the boundaries between species of *Ceratocystis*. During the course of the last decade, the use of phylogenetic studies based on comparisons of DNA sequence data has contributed substantially to clarify the taxonomy of this group of fungi. These studies have led to the conclusion that some *Ceratocystis* species thought to be single entities in the past actually represent complexes of cryptic species. This review is primarily intended to provide an introduction to experimental studies presented on species of *Ceratocystis sensu stricto*. However, in order to provide a firm background to this fungal genus, a brief outline is presented on related fungi that can broadly be referred to as the ophiostomatoid fungi.

Key words: Ophiostomatoid fungi, phylogenetic analysis, cryptic species, forest pathology.

RESUMEN

REVISIÓN DE *Ceratocystis sensu stricto* CON ESPECIAL REFERENCIA A LOS COMPLEJOS DE LAS ESPECIES *C. coerulescens* Y *C. fimbriata*

El género *Ceratocystis sensu stricto* incluye un alto número de hongos fitopatógenos de plantas angiospermas y gimnospermas en diversas regiones del mundo. Entre las principales enfermedades

¹ Profesor Asistente, Universidad Nacional de Colombia, Sede Medellín. Escuela de Biociencias. A.A. 3840, Medellín, Colombia. <mamarinm@unalmed.edu.co>

² Director. Forestry and Agricultural Biotechnology Institute (FABI). Department of Microbiology and Plant Pathology. University of Pretoria, Pretoria, South Africa. <mike.wingfield@fabi.up.ac.za>

Recibido: Agosto 18 de 2005; aceptado: Febrero 16 de 2006.

causadas por estos hongos se destacan los marchitamientos vasculares, manchado de maderas, chancros y pudriciones radiculares, de tallos y frutos. Muchas especies producen metabolitos volátiles con olores que atraen una amplia variedad de insectos a los tejidos vegetales infectados, sin embargo, el grado de asociación entre los insectos y las diferentes especies de **Ceratocystis** es altamente variable, presentándose asociaciones mutualistas tales como aquellas entre **C. polonica**, **C. laricicola** y **C. rufipenni** e insectos escoltídos, así como también relaciones no específicas como en los casos de **C. paradoxa**, **C. fagacearum** y **C. fimbriata** con nitidúlidos, moscas y diversos coleópteros. El alto grado de similitud morfológica encontrada entre estos hongos hace difícil la definición de los límites entre las especies de **Ceratocystis**. Durante la pasada década, la utilización de estudios filogenéticos basados en comparaciones de DNA ha contribuido sustancialmente a clarificar la taxonomía de este grupo de hongos, así por ejemplo, dichos estudios han permitido concluir que algunas especies de **Ceratocystis** realmente representan complejos de especies crípticas. Esta revisión tiene como objetivo principal proveer una introducción a los estudios experimentales desarrollados para las especies de **Ceratocystis** *sensu stricto* y brevemente discutir aspectos relacionados con los denominados hongos ophiostomatoides.

Palabras claves: Hongos ophiostomatoides, análisis filogenético, especies crípticas, patología forestal.

The ophiostomatoid fungi include more than 100 species of ascomycetes that share morphological characteristics and adaptation to dispersal by insects and other arthropods (Wingfield *et al.* 1993, Harrington 1993a, Kile 1993). These fungi are found on a wide variety of substrates in tropical, subtropical and temperate regions of the world. Ecologically, they occupy a wide range of niches, ranging from saprotrophs to primary pathogens of both angiosperm and gymnosperm plants. Some species are agents of sap stain on conifers and hardwood trees, while others are aggressive pathogens of economically important agronomic crops (e.g. sweet potato, tobacco, coffee, citrus, cacao, pineapple, sugarcane, rubber) or forest trees (e.g. oak, elm, eucalyptus and plane trees) (Upadhyay 1981 1993, Wingfield *et al.* 1993). In addition, several ophiostomatoid fungi cause virulent

systemic diseases in humans and animals (Summerbell *et al.* 1993).

Morphologically, the ophiostomatoid fungi are characterised by spherical ascomatal bases with elongated necks, deliquescent asci borne irregularly throughout the centrum of the ascocarps, and single-celled ascospores that lack germ pores, often have ornamented sheaths and are produced in a mucilaginous matrix that exudes in sticky droplets through ostioles at the apices of the necks. The many different anamorph states found among the ophiostomatoid fungi all have holoblastic conidium development, but display a variety of conidiophore and conidial forms (Hunt 1956, Upadhyay 1981 1993, Spatafora and Blackwell 1994). The sticky spore masses (including the conidial masses produced by some species) facilitate insect dispersal, as they

easily adhere to the insect bodies (Malloch and Blackwell 1993). The high degree of morphological similarity found amongst these fungi is probably a result of convergent evolution, driven by selection for insect dispersal (Spatafora and Blackwell 1994).

The morphological similarity among the ophiostomatoid fungi made it difficult to establish valuable taxonomic characters to delimitate them at both lower (species, genera) and higher (families, orders) taxonomic levels (Spatafora and Blackwell 1994). Consequently, controversial and often confusing taxonomic debates have surrounded their classification. This confusion stems back to even 1891, when Halsted and Fairchild misinterpreted the ascocarps of the first described species, *Ceratocystis fimbriata* Ellis & Halst. as pycnidia (Nannfeldt 1932, Hunt 1956, Griffin 1968, De Hoog 1974, Benny and Kimbrough 1980, Upadhyay 1981, De Hoog and Scheffer 1984, Von Arx and Van Der Walt 1988). Indeed the term "ophiostomatoid fungi" to some extent illustrates this confusion and emerged from a meeting of scientists in 1990 who recognised that genera such as *Ophiostoma* and *Ceratocystis* were phylogenetically unrelated but morphologically and ecologically similar (Wingfield *et al.* 1993).

Fifteen genera have historically been grouped within what are considered to be ophiostomatoid fungi. Some of these genera such as *Ceratocystis* Ellis &

Halstead and *Ophiostoma* H. & P. Sydow, have been widely accepted, whereas others have had limited recognition and have later been considered as synonyms or invalid genera (e.g. *Ceratocystiopsis* pad. & Kendrick, *Endoconidiophora* Münch, *Europhium* Parker) (Wingfield, Seifert and Webber 1993). To complicate matters further, more than 15 hyphomycete genera with a wide range of conidiophores, conidium ontogeny, and conidial forms, have been linked to the ophiostomatoid fungi (Upadhyay and Kendrick 1975, Upadhyay 1981, 1993).

A major theme of the taxonomic debates surrounding the ophiostomatoid fungi has been whether they represent one or several genera, and more specifically whether *Ceratocystis* and *Ophiostoma* should be treated as congeneric (Hunt 1956, Griffin 1968, Olchoweki and Reid 1974, Upadhyay 1981 1993) or as distinct genera (De Hoog 1974, De Hoog and Scheffer 1984, Wingfield, Van Wyk y Marasas 1988). However, more recent phylogenetic analyses based on DNA sequence comparisons have demonstrated unequivocally that despite their morphologically similar teleomorphs, these genera are phylogenetically unrelated even at the ordinal level, with *Ceratocystis* residing in the Microascales and *Ophiostoma* being more closely related to species in the Diaporthales (Hausner *et al.* 1993a, b, Spatafora and Blackwell 1994, Wingfield, Viljoen, y Wingfield 1999). Currently, five genera are accepted to belong to

the wider group referred to as ophiostomatoid fungi. These include *Ceratocystis*, *Ophiostoma*, *Ceratocystiopsis*, *Gondwanamyces* Marais & Wingfield and *Cornuvesica* Viljoen & Winfield (Wingfield, Seifert and Webber 1993, Upadhyay 1993, Marais *et al.* 1998, Viljoen, Wingfield and Wingfield 2000). This is despite the fact that they represent polyphyletic lineages.

***Ceratocystis*.** The name *Ceratocystis sensu lato* has been used collectively for all the ophiostomatoid fungi, while *Ceratocystis sensu stricto* is used exclusively for species with anamorphs in the form genus *Thielaviopsis* Went emend. Paulin, Harrington *et al.*, McNew (Wingfield *et al.* 1993, Paulin-Mahady, Harrington, y McNew 2002). These anamorphs are distinguished by conidium development of ring wall building (enteroblastic conidiogenesis) (Minter, Kirk y Sutton 1983, Minter 1987), and were recently transferred from the genus *Chalara* (Corda) Rabenh. to *Thielaviopsis* based on analysis of sequence data from the ribosomal RNA (rRNA) operon (Paulin-Mahady, Harrington y McNew 2002). Species of *Ceratocystis sensu stricto* can also be distinguished from other ophiostomatoid fungi by the lack of cellulose and rhamnose in their cell walls (Spencer and Gorin 1971, Jewell 1974, Weijman and De Hoog 1975) and their sensitivity to the antibiotic cycloheximide (Harrington 1981).

***Ophiostoma*.** Fungi residing in the genus *Ophiostoma* are characterised by having cellulose and rhamnose in their cell walls and being tolerant to cycloheximide in culture (Harrington 1981, De Hoog and Scheffer 1984). The anamorphs of these species are accommodated in several hyphomycete genera, such as *Leptographium* Lagerberg & Melin, *Pesotum* Crane & Schocknecht, *Sporothrix* Hektoen & Perkins: Nicot & Mariat, *Knoxdaviesia* Wingfield, van Wyk & Marasas and *Hyalorhinocladiella* Upadhyay & Kendrick (Wingfield *et al.* 1993, Okada *et al.* 1998, Hausner, Reid y Klassen 2000, Jacobs and Wingfield 2001). In all these genera conidium development occurs exogenically through apical wall building (holoblastic conidiogenesis) (Minter, Kirk y Sutton 1982).

***Ceratocystiopsis*.** The genus *Ceratocystiopsis* was established to accommodate ophiostomatoid fungi with elongate or falcate ascospores surrounded by a falcate sheath with attenuated ends (Upadhyay and Kendrick 1975, Upadhyay 1981, Wingfield 1993). Anamorphs of these species are formed by both enteroblastic and holoblastic conidiogenesis and have included *Sporothrix*, *Hyalorhinocladiella*, *Chalara* and *Knoxdaviesia* states. However, the validity of this genus remains uncertain, as morphological and molecular data have shown that most species should be included within *Ophiostoma* (Wingfield 1993, Hausner, Reid y Klassen 1993a 1993).

b) but that others with *Knoxdaviesia* and *Chalara* anamorphs reside in genera such as *Gondwanamyces* and *Cornuvesica*, respectively (Marais *et al.* 1998, Viljoen *et al.* 2000) that are unrelated to *Ophiostoma*.

***Gondwanamyces* and *Cornuvesica*.** The genus *Gondwanamyces* was recently established to accommodate two ophiostomatoid species with *Knoxdaviesia* anamorphs (*Ceratocystiopsis proteae* Wingfield, Van Wyk & Marasas and *Ophiostoma capense* Wingfield & Van Wyk) (Marais *et al.* 1998). *Cornuvesica* was erected to include the former species *Ceratocystiopsis falcata* (Wright & Cain) Upadhyay, which has a *Chalara*-like anamorph but is phylogenetically unrelated to *Ceratocystis sensu stricto* (Hausner, Reid y Klassen 1993-b, Viljoen, Wingfield y Wingfield 2000).

Ceratocystis sensu stricto. The genus *Ceratocystis sensu stricto* includes many economically important plant pathogens that especially occur in tropical and sub-tropical regions of the world. Most species require wounds in order to infect their hosts, and their pathogenicity varies widely from weakly virulent fungi to very aggressive pathogens. Some species have even a saprotrophic life style (*Ceratocystis moniliformis* (Hedgc.) C. Moreau). Diseases caused by these fungi include vascular wilts, sap and lumber stains, stem cankers and rots of roots, stems and fruits

(Kile 1993, Harrington, Wingfield and Kile 1996).

The most virulent *Ceratocystis* species generally cause vascular wilts and stem cankers of trees and other woody plants. For example, *Ceratocystis fagacearum* (Bretz) Hunt causes the disease known as oak wilt of *Quercus* spp. in North America (Henry *et al.* 1944), *Ceratocystis fimbriata* causes wilt and cankerstain diseases on a wide range of hosts including cocoa (*Theobroma cacao* L.) (Schieber 1969), coffee (*Coffea arabica* L.) (Pontis 1951) and plane trees (*Platanus* spp.) (Walter 1946, Panconesi 1981), *Chalara australis* Walker & Kile cause a serious wilt disease of *Nothofagus cunninghamii* (Hook.) Oerst (Kile and Walker 1987) and *Ceratocystis polonica* (Siemaszko) Moreau displays relatively high levels of virulence to Norway spruce (*Picea abies* L. Karst.) and other spruce species in Eurasia (Siemaszko 1939, Solheim 1986 1988, Krokene and Solheim 1998, Kirisits 2001).

Weaker pathogens include agents of sap stain of lumber such as *Ceratocystis coerulescens* (Münch) Bakshi on conifers and *Ceratocystis douglasii* (Davidson) Wingfield & Harrington on Douglas-fir (*Pseudotsuga menziesii* (Mirb.) Franco) (Münch 1907, Harrington, Wingfield and Kile 1996, Wingfield, Harrington and Solheim 1997, Harrington and Wingfield 1998). Other weak pathogens include species producing lumber stain in hardwoods such as *Ceratocystis vires-*

ens (Davidson) C. Moreau on maple (*Acer saccharum* Marsh.) (Davidson 1944) and *Ceratocystis eucalypti* Yuan & Kile on *Eucalyptus* spp. (Kile *et al.* 1996).

Likewise, various annual crop plants are also seriously affected by *Ceratocystis* species. *Ceratocystis paradoxa* (Dade) Moreau causes tissue rot on sugar cane (*Saccharum officinarum* L.), *Musa* spp. and *Cocos nucifera* L. (Kile 1993). Similarly, *Ceratocystis adiposa* (Hedgcock) Moreau incites tissue rot on sugar cane (Butler 1906) and *Ceratocystis radicola* (Bliss) Moreau causes a root rot on date palms (*Phoenix dactylifera* L.) (Bliss 1941).

Members of the genus *Ceratocystis* have several means of transmission and dispersal, in addition to being vectored by insects. Species producing chlamydo-spores are usually soil-borne (Rossetto and Ribeiro 1990, Kile 1993), whereas those that sporulate aeri-ally are dispersed by wind and water splash (Baker and Thomas 1946, Vigouroux and Stojadinovic 1990). Species that cause vascular wilts and woodstain can also be transmitted through root grafts (Gibbs and French 1980, Accordi 1986) and by pruning tools. This is especially important for the spread of *Ceratocystis* pathogens in agricultural crops and forest plantations (Walter, Rex and Schreiber 1952, Matasci and Gessler 1997).

Most *Ceratocystis* species are well adapted to dispersal by insects. Many

species produce volatile metabolites with sweet fruity odours that attract a wide range of insects to infected tissues (Hanssen 1993, Kile 1993). However, the intimacy at the association between insects and *Ceratocystis* species is highly variable. Species such as *C. polonica*, *Ceratocystis laricicola* Redfern & Minter and *Ceratocystis rufipenni* Wingfield, Harrington & Solheim, seem to have a mutualistic relationship with their bark beetle vectors (Redfern *et al.* 1987, Solheim 1988, Wingfield, Harrington and Solheim 1997). On the other hand, species such as *C. paradoxa* (Chang and Jensen 1974), *C. fagacearum* (Juzwik and French 1983, Appel, Kurdyla and Lewis Junior 1990), *C. fimbriata* (Crone and Bachelder 1961, Hinds 1972) and *C. moniliformis* (Hinds and Davidson 1972) have non-specific associations with fungivorous or sap-feeding insects such as nitidulid beetles (Coleoptera: Nitidulidae), flies (Diptera: Drosophilae) and ambrosia beetles (Peplinski and Merrill 1974, Kile 1993).

Different modes of reproduction are found among *Ceratocystis* species, including heterothallism, homothallism and exclusive asexual reproduction (Harrington, Steimel and Kile 1998). Genetic studies have shown that sexuality in these species is controlled by two mating-type genes designated MAT-1 and MAT-2 (Harrington and McNew 1997, 1998). In strictly heterothallic species, such as *C. eucalypti* and *C. fagacearum*,

sexual progeny are only produced upon mating of strains representing different mating types (Kile *et al.* 1996, Harrington, Steimel and Kile 1998). Other species, such as *Ch. australis* and *Chalara neocaledoniae* Kiffer & Delon, are known only by their anamorph states, and supposedly only reproduce asexually (Kile *et al.* 1996, Harrington, Steimel and Kile 1998). However, most *Ceratocystis* species, including among others, *C. fimbriata*, *C. coerulescens*, *C. virescens*, *C. polonica* and *Ceratocystis pinicola* Harrington & Wingfield, are homothallic and also undergo unidirectional mating type switching (Harrington and McNew 1997, 1998). In these species, self-fertile strains that contain both mating type idiomorphs give rise to both self-fertile and self-sterile strains. The latter contain only the MAT-1 gene, and these MAT-1 strains can cross with the self-fertile strains. Progeny of selfing and crossing events segregate in a 1:1 ratio, with half the ascospores being self-fertile (MAT-1 and MAT-2) and the other half self-sterile

(MAT-1). It has been shown that self-sterile strains are produced as the result of a deletion of the MAT-2 idiomorph, which then results in the expression of only the MAT-1 mating type (Witthuhn *et al.* 2000).

The close morphological similarity between many *Ceratocystis* species contributes to poorly defined species boundaries within the genus (Harrington, Wingfield and Kile 1996). Phylogenetic studies using DNA sequence data from the rRNA operon (Wingfield *et al.* 1994, Visser *et al.* 1995, Wingfield *et al.* 1996, Witthuhn *et al.* 1998, 1999, Barnes *et al.* 2003), mating type genes (Witthuhn *et al.* 2000, Harrington *et al.* 2002) and microsatellite DNA regions (Barnes *et al.* 2001) have suggested that some *Ceratocystis* species that have been regarded as single entities in the past, actually represent complexes of cryptic species. Two of these species complexes, the *C. coerulescens* and *C. fimbriata* complex, are treated in more detail below (Figure 1).




Figure 1. Phylogeny of the *Ceratocystis coerulescens* species complex and its relationship with *Ceratocystis fimbriata*. The phylogenetic tree was produced from the DNA sequence of a 565 bp region of the ribosomal RNA operon using PAUP (Witthuhn et al. 1998).

***Ceratocystis coerulescens* complex.** Eleven species have been grouped in the *Ceratocystis coerulescens* complex. These species have similar morphology and are closely related based on DNA sequence data and isozyme profiles (Harrington, Wingfield and Kile 1996, Harrington and Wingfield 1998, Witthuhn et al. 1999, 2000). Seven species cause sap stain on conifers in the family Pinaceae, while four species are pathogens or incite sap stain on hardwood trees. Phylogenetic analysis using sequences of the rRNA operon and MAT-2 HMG box have

shown that the species occurring on conifers form a single monophyletic clade, with the hardwood species as their basal ancestor (Witthuhn et al. 1998, 2000). It has thus been suggested that adaptation to conifer hosts occurred once in the evolutionary history of *Ceratocystis* (Witthuhn et al. 1998, 2000).

Ceratocystis coerulescens was the first species described in the group colonizing conifers (Münch 1907). This fungus causes blue-stain of spruce and pine sapwood in Europe and North America (Münch 1907,

Harrington, Wingfield and Kile 1996, Harrington and Wingfield 1998). *Ceratocystis pinicola* occurs in Britain and is an agent of bluestain on pine logs and lumber (Gibbs 1993, Harrington and Wingfield 1998). *Ceratocystis rufipenni* was described from Engelmann spruce (*Picea engelmannii* Parry ex Engelm.) and white spruce (*Picea glauca* (Moench) Voss) infested by the North American spruce bark beetle, *Dendroctonus rufipennis* Kirby in British Columbia (Wingfield, Harrington and Solheim 1997). It is virulent on Sitka spruce (*Picea sitchensis* (Bong.) Carr.) and can kill trees that have been inoculated in sufficiently high dosages (Solheim and Safranyik 1997). *Ceratocystis resinifera* Harrington & Wingfield has been found in continental Europe and North America and colonizes wounds of live spruce trees (Harrington, Wingfield and Kile 1996, Harrington and Wingfield 1998). *Ceratocystis douglasii* has originally been recognized as a form of *C. coerulescens* (Davidson 1953), but has subsequently been described as a new species (Wingfield, Harrington and Solheim 1997). This fungus stains the sapwood of Douglas-fir in western North America, and appears to be restricted to the natural range of its host (Wingfield, Harrington and Solheim 1997, Harrington and Wingfield 1998). *Ceratocystis laricicola* infests larch (*Larix* spp.) and has a mutualistic association with the bark beetle *Ips cembrae* Heer in Europe and Japan (Redfern *et al.* 1987, Yamaoka *et al.*

1998, Kirisits 2001, Stauffer *et al.* 2001). The remaining species in the conifer subgroup is *C. polonica*, which is a pathogen and agent of blue-stain on Norway spruce and other spruce species in Europe and Japan (Solheim 1986, Yamaoka *et al.* 1997). This *Ceratocystis* species is primarily vectored by the aggressive bark beetle *Ips typographus* L. (Siemaszko 1939, Furniss, Solheim and Christiansen 1990). In addition, *Ips typographus* f. *japonicus* Nijima in Japan (Yamaoka *et al.* 1997) as well as *Ips amitinus* Eichhoff (Kirisits 2001) and *Ips duplicatus* Sahlberg (Krokene and Solheim 1996) in Europe have recently been identified as important vectors of *C. polonica*.

The four *Ceratocystis* species occurring on hardwoods include *C. virescens* that causes sapstreak in maple and other hardwood species in eastern North America (Davidson 1944, Harrington and Wingfield 1998), *C. eucalypti* that is a wound colonist on *Eucalyptus* spp. in Tasmania and Victoria, Australia (Kile *et al.* 1996), and two asexual species occurring in Australia, *Ch. australis* and *Ch. neocaledoniae*. The first of these is a virulent pathogen causing myrtle wilt on *Nothofagus cunninghamii* in Tasmania (Kile *et al.* 1996). The latter species produces a vascular wilt disease on coffee (*Coffea robusta* Linden ex Wild.) and guava trees (*Psidium guajava* L.) (Kile and Walker 1987).

The *Ceratocystis* species occurring on hardwoods and conifers can be

distinguished morphologically, based on conidiophore characteristics. The hardwood group has tapering and proliferating phialides, while the conifer species lack these phialide characters (Harrington, Wingfield and Kile 1996). The conifer species can be reliably subdivided into two morphologically well-defined groups: *C. coerulescens sensu lato*, where the ascospores are surrounded by an even sheath that is wider at the ends than at the sides, and *C. polonica sensu lato*, where ascospore sheaths are wider at the sides than at the ends (Harrington and Wingfield 1998). Within *Ceratocystis coerulescens sensu lato* on conifers, small morphological differences exist (Harrington and Wingfield 1998), however, unequivocal separation of the entire conifer colonizing species is practically only possible through isozyme and DNA sequence comparisons (Harrington, Wingfield and Kile 1996, Witthuhn *et al.* 1998, 2000, Harrington *et al.* 2002).

Ceratocystis polonica sensu lato.

As discussed above, species in the *C. coerulescens* complex have been widely studied during the course of the past decade. These studies have included several phenotypic and genotypic characters (Harrington, Wingfield and Kile 1996, Harrington and McNew 1998, Witthuhn *et al.* 1998, 2000, Harrington and Wing-

field 1998, Harrington *et al.* 2002). They have added considerable resolution to the taxonomy of nine of these species, but problems have remained in distinguishing between *C. polonica* and *C. laricicola*.

Ceratocystis polonica was first isolated from Norway spruce in Poland and described by Siemaszko 1939 as *Ophiostoma polonicum*. Visser *et al.* 1995 reported the presence of a *Chalara* (*Thielaviopsis*) anamorph in cultures of *O. polonicum*, and this characteristic, together with DNA sequence comparisons, clearly showed that this fungus is a species of *Ceratocystis*. *Ceratocystis laricicola*, on the other hand, was correctly described by Redfern *et al.* 1987. It was, however, not compared with *C. polonica*, probably due to the fact that *C. polonica* was not recognised as having a *Chalara* state at the time when *C. laricicola* was described (Harrington and Wingfield 1998). Later studies showed the two species were morphologically indistinguishable (Harrington and Wingfield 1998), had identical ITS sequences (Witthuhn *et al.* 1998), and a low level of isozyme variation (Harrington, Wingfield and Kile 1996). However, they represent distinct ecological entities (Siemaszko 1939, Solheim 1986, Redfern *et al.* 1987) and are true biological species, as interspecific crosses between MAT-1 and MAT-2 strains of both species fail to produce viable progeny (Harrington and McNew 1998, Harrington *et al.* 2002).

These two fungi, together with *C. rufipenni* are especially important for forestry, because they are associated with aggressive tree-killing bark beetles. *Ceratocystis polonica* is a mutualistic associate of the Eurasian spruce bark beetle, *I. typographus*, which is a highly destructive pest of Norway spruce throughout Eurasia (Postner 1974, Christiansen and Bakke 1988). Since the 18th century *I. typographus* has caused large-scale outbreaks with millions of killed trees (Klimetzek and Yue 1997) in Scandinavia and central Europe (Postner 1974, Christiansen and Bakke 1988, Führer 1996, Kirisits 2001). The most recent and still ongoing outbreak of *I. typographus* in central and western Europe started around 1992 and has been triggered by destructive windthrows, followed by years with warm and dry climate (Führer 1996, Kirisits 2001). From 1985 to 1994, the *I. typographus* – *C. polonica* complex has been responsible for losses of about 14,1 million cubic meters of spruce wood in Austria, Switzerland and the German province Baden-Württemberg alone (Führer 1996).

Ceratocystis laricicola is vectored by the eight spined larch bark beetle, *I. cembrae* (Redfern *et al.* 1987, Yamaoka *et al.* 1998, Stauffer *et al.* 2001, Kirisits 2001), which is distributed across continental Europe and Asia (Postner 1974, Pfeffer 1995) and has been introduced to Scotland and Denmark (Crooke and Bevan 1957, Redfern *et al.* 1987, Stauffer *et al.* 2001). This bark beetle is regarded as

a secondary although important pest of larch and plays a particularly important role in regions outside the natural range of European larch (Schimitschek 1931, Postner 1974, Stauffer *et al.* 2001). For example, *I. cembrae* and *C. laricicola* have become principal pests of larch in Scotland, after their introduction in 1955 with imported timber from Germany (Crooke and Bevan 1957, Redfern *et al.* 1987).

Mass inoculation experiments that aim to mimic natural mass-attacks by bark beetles, have demonstrated that both *C. polonica* (Christiansen 1985, Solheim 1988, Kirisits 1998, Krokene and Solheim 1998, Yamaoka, Takahashi and Iguchi 2000, Kirisits and Offenthaler 2002) and *C. laricicola* (Redfern *et al.* 1987, Yamaoka *et al.* 1998, Kirisits 2001) are able to kill their respective host trees. It is thought that through the mutualistic association between these bark beetles and the specific *Ceratocystis* spp. that they carry, the fungus increases the impact of each beetle attack and thereby contributes to exhaust the host defences and makes the tree suitable for bark-beetle reproduction (Whitney 1982, Harrington 1993b, Paine, Raffa and Harrington 1997). Reciprocally, these fungi benefit by being dispersed and introduced into suitable host trees by their beetle vectors (Whitney 1982, Malloch and Blackwell 1993, Krokene 1996). Ironically, the less virulent pathogenic *Ophiostoma* spp. associated with these insects (Solheim 1986,

Krokene 1996, Krokene and Solheim 1998, Kirisits 2001, Viiri 2002) gain the same dispersal advantage, simply by being present in this environment.

Mechanisms of fungal colonization and host responses have been particularly well studied in the *C. polonica* – Norway spruce system. Fungal succession studies have shown that *C. polonica* is the first species to invade the sapwood after *I. typographus* has attacked a tree (Solheim 1992). This is probably due to its rapid growth and its ability to tolerate the low oxygen levels found in living sapwood (Solheim 1991). However, not all isolates of *C. polonica* have the same ability to kill trees and to induce intensive blue-stain in the sapwood of its host tree. Isolates with low levels of virulence have been reported from independent mass inoculation experiments in Austria (Kirisits and Anglberger 1999, Baier *et al.* 2002) and Norway (Krokene and Solheim 2001). Kirisits and Anglberger 1999 considered the low level of virulence as a general characteristic of the particular isolate used in their experiment. Likewise, in the Norwegian study loss of pathogenicity was attributed to senescence of the isolates after several transfers of cultures during routine storage or, alternatively, to long periods of storage of the isolates (Krokene and Solheim 2001). However, mycoviral infections, such as those reported in the forest tree pathogens *Ophiostoma novoulmi* (Rogers, Buck, and Braiser 1987, Cole *et al.* 1998) and

Cryphonectria parasitica (Nuss 1992, 2000), might have been involved but were not considered.

Spruce trees infected by *C. polonica* respond in several ways to contain the fungus. Responses such as cellular proliferation, mobilization of polyphenolic parenchyma cells, formation of traumatic resin ducts, lignification (Nagy *et al.* 2000, Evensen *et al.* 2000, Krokene, Solheim and Christiansen 2003) as well as translocation of starch reserves (Christiansen and Ericsson 1986) have been shown to be part of the defence mechanisms. Furthermore, experiments have shown that preinoculation of spruce trees with non lethal dosages of *C. polonica* inoculum induces disease resistance that protects trees against subsequent mass infections of the same fungus (Christiansen *et al.* 1999, Krokene and Solheim 2001). The mechanisms underlying this induced resistance differ from the systemic acquired resistance (SAR) present in angiosperms, since it is non-systemic and non-specific (Krokene, Solheim and Christiansen 2003). The induced resistance of Norway spruce is due to anatomical and biochemical reactions in the phloem and the sapwood, in which polyphenolic parenchyma cells and traumatic resin ducts are involved (Krokene, Solheim and Christiansen 2003). Different necrotizing fungi, chemical elicitors and even mechanical wounding can activate a similar defense response compared to that induced by pre-

treatment with *C. polonica* (Christiansen *et al.* 1999, Krokene and Solheim 2001, Franceschi, Krekling and Christiansen 2002).

The *Ceratocystis fimbriata* complex. *Ceratocystis fimbriata* was first reported causing rot of sweet potato (*Ipomoea batatas* L.) in 1890 in New Jersey, USA (Halsted 1890). It is the type species of the genus *Ceratocystis* and as in the case of other *Ceratocystis* species, its taxonomy was unclear for more than fifty years. Saccardo 1892 reclassified this fungus as *Sphaeronaema fimbriatum*, while Zimmerman 1900 described a strain of *C. fimbriata* causing coffee canker in Java (Indonesia) as a separate species, *Rostrella coffea* Zimmermann. Elliott 1923 renamed *C. fimbriata* as *Ceratostomella fimbriata*, but when Nannfeldt 1932 erected the family Ophiostomataceae, the species was transferred to the genus *Ophiostoma*, and later, based on the presence of endoconidia (now recognised as the *Thielaviopsis* state) to *Endoconidiophora* (Davidson 1935). The name *Ceratocystis fimbriata* was recognized again by Bakshi 1950, who revived the generic name *Ceratocystis* and considered *Ophiostoma*, *Endoconidiophora*, *Rostrella*, *Linostoma* and *Gros-mannia* as synonyms of *Ceratocystis*. When Hunt 1956 accepted Bakshi's 1950 amendment, the name *C. fimbriata* became widely accepted (Griffin 1968, Von Arx 1974, Upadhyay 1981).

Morphologically, *C. fimbriata* is characterised by perithecia with long necks that taper towards the tips and terminate in 8 to 15 convergent ostiolar hyphae. Perithecial bases are dark and globose, surrounded by a dense network of hyphae. Asci are evanescent in the early stages of development, while ascospores are characteristically hat-shaped and exuded through the perithecial necks in sticky masses (Hunt 1956, Upadhyay 1981). The fungus also produces chains of cylindrical conidia and aleurioconidia (chlamydospores) that play an important role in the survival of this fungus in the soil (Pontis 1951, Webster and Butler 1967, Accordi 1989). The anamorph of *C. fimbriata* has for many years been accommodated in *Chalara* (Hunt 1956, Nag Raj and Kendrick 1975, Upadhyay 1981), but recently, based on a phylogenetic analysis of DNA sequence data, has been recognised as best residing in *Thielaviopsis* (Paulin-Mahady *et al.* 2002).

Ceratocystis fimbriata is one of the most aggressive plant pathogens in the genus *Ceratocystis*, causing wilt, canker-stain diseases as well as tissue rot on a wide variety of perennial as well as agronomic crop plants, worldwide (Kile 1993). More than 30 economically important plants are attacked by this pathogen (Baker and Harrington 2000) including, among others, plane trees (Walter, Rex and Schreiber 1952, Panconesi 1981), *Eucalyptus* spp. (Roux *et al.* 1999, Roux *et al.* 2001), aspen (*Populus tre-*

muloides Michx.) (Zalasky 1965, Hinds 1972), rubber (*Hevea brasiliensis* (Willd: Juss.) Mull-Arg.) (Silveira *et al.* 1994), mango (*Mangifera indica* L.) (Batista 1960), almond (*Prunus amygdalus* Batsch.) (De Vay *et al.* 1968, Teviotdale and Harper 1991), *Syngonium* spp. (Walker *et al.* 1988) and pimento trees (*Pimenta officinalis* L.) (Leather 1966).

Ceratocystis fimbriata is particularly important in Latin America, where it causes substantial losses in three of the main perennial crops that are grown in this region: coffee, cocoa and citrus (*Citrus* spp.). Coffee canker has been reported from Colombia (Obregon 1936), Venezuela (Pontis 1951), Guatemala (Szkolnik 1951) and Costa Rica (Echandi and Segall 1956). Symptoms of diseased plants include yellowing of foliage, dieback and wilt. When the outer surface of the stem bark of infected trees is removed, dark lesions are obvious and these usually extend downwards towards the roots and girdle the trunks, causing tree death (Pontis 1951, Castro 1991). Coffee canker has become one of the most important threats to coffee production in Colombia, where more than 800000 hectares of this crop are planted (Castro 1998). The principal mode of ingress for *C. fimbriata* into coffee trunks is through mechanical wounds (Pontis 1951). The fungus is easily transported on agricultural tools or with soil. Thus, a major factor associated with spread of coffee canker in Colombia is infection through wounds

on the lower stems made by the shoes of farmers needing to secure themselves on the steep slopes on which coffee is cultivated (Castro 1991).

Ceratocystis fimbriata was first reported on cacao in Ecuador in 1918, causing a serious wilt disease (Desrosiers 1958). This disease was subsequently found in Venezuela (Malaguti 1952), Colombia (Idrobo 1958), Trinidad (Goberdhan 1959), Costa Rica (Havord 1962) and the Dominican Republic (Schieber 1969). Recently, cacao wilt disease has also been detected in Brazil, in plantations from South Bahia (Bezerra 1997). Cacao wilt is frequently associated with wounds made by machetes and other tools. The mycelium and spores enter the xylem and ray parenchyma cells through such fresh wounds, and cause dark reddish to purple staining that may develop into a vascular wilt. Externally, the disease is easily recognized because wilted leaves become dry, curled and remain attached to the trees for several weeks (Montes de Oca 1975). The incidence of cacao wilt has been reduced significantly in Latin America in recent years, especially because resistant varieties have become widely available for planting (Delgado and Echandi 1965, Ocampo, Mafla and Victoria 1982, Simmonds 1994).

The first report of *C. fimbriata* in citrus orchards was made in 1981, when it caused serious dieback on lime trees (*Citrus aurantifolia*

(Christm.) Swing) in the central region of Colombia (Instituto Colombiano Agropecuario 1993). Three years later, it was also found on lemon trees (*Citrus limon* Burmman) in Argentina (Contreras and Marmeloz 1984). By 1994, about 10% of the citrus trees in Colombia had been killed by the pathogen (Mourichon 1994). The disease is characterised by a general yellowing of the foliage, followed by heavy defoliation and prominent shoot proliferation. Internally, brown to reddish lesions are evident in the stele of the trunk, and, as the disease progresses, lesions converge and become flame shaped, black in colour, and trees are eventually killed (Mourichon 1994).

The taxonomic complexity of what has been referred to as *C. fimbriata* in Latin America became evident in a recent study based on ITS sequence comparisons and microsatellite analysis (Barnes *et al.* 2001). Isolates of *C. fimbriata* from coffee in Colombia were found to reside in two genetically isolated groups that might represent different species. However, apart from this study and scattered information that is not widely available in the scientific literature, very little is known regarding the biology, ecology or genetic background of *C. fimbriata* in Latin America. For example, in countries such as Colombia and Costa Rica, several native hosts (e.g. *Inga* sp., *Herrania* sp., *Annona muricata* L., *Caryodendron orinocense* Karst.) are infected by this pathogen (Montes de

Oca 1975, Pardo-Cardona 1995 and Buriticá 1999), but their role as alternative hosts and sources of inoculum for susceptible agricultural crops remains uncertain.

Ceratocystis fimbriata has long been recognised as a variable fungus consisting of distinct, host-specialized strains (Webster and Butler 1967, Harrington 2000). The best studied of these host specific races is *C. fimbriata forma specialis platani*, which causes a canker stain disease of plane trees in the Northern hemisphere (Walter, Rex and Schreiber 1952, Santini and Capretti 2000). Similarly, reciprocal inoculation experiments have demonstrated the existence of host-specific isolates on pimento (Leather 1966), sweet potato (Kojima and Uritani 1976), *Syngonium* (Vogelzang and Scott 1991) and cocoa (Baker and Harrington 2000). Earlier genetic analyses indicated that such host-specialized isolates could represent distinct, but closely related species (Olsen and Martin 1949, Webster and Butler 1967). The view that the host is an important factor separating strains of *C. fimbriata sensu lato* has recently also been confirmed based on DNA sequence data (Wingfield *et al.* 1996, Witthuhn *et al.* 1998, Santini and Capretti 2000, Barnes *et al.* 2001, 2003). Witthuhn *et al.* 1999 found that isolates of *C. fimbriata* from *Populus* spp., *Prunus* spp. and *Platanus* spp. could be separated based on RFLP analysis of the ITS regions of the rRNA operon. Similar

results were obtained by Santini and Capretti 2000 and Barnes *et al.* 2001, using RAPD / minisatellite markers and microsatellite markers, respectively. Thus, *C. fimbriata* strains from different hosts and geographical regions grouped in separated clades in accordance with their respective hosts and origins.

The view that *C. fimbriata* may represent a complex of cryptic species has gained further support by the recent description of two new *Ceratocystis* species that probably used to be treated within *C. fimbriata*. *Ceratocystis albofundus* Wingfield, De Beer & Morris was reported causing wilt and dieback of *Acacia mearnsii* Wildeman in South Africa (Morris, Wingfield and De Beer 1993, De Beer 1994). Detailed morphological and molecular comparisons showed that the fungus represents a distinct *Ceratocystis* species that is characterised by light-coloured perithecial bases, dark necks, divergent ostiolar hyphae and distinct ITS sequences (Wingfield *et al.* 1996, Roux *et al.* 1999). The other new species from wounds on *Eucalyptus* in Australia has recently been described as *Ceratocystis pirilliformis* Barnes & Wingfield (Barnes *et al.* 2003). This species has hatshaped ascospores similar to those found in *C. fimbriata*, but differs from this species in its pear-shaped perithecial bases. Phylogenetic analysis of the rRNA ITS regions also supported the definition of this fungus as a new *Ceratocystis* species (Barnes *et al.* 2003).

CONCLUSIONS

Selection pressure for insect dispersion has resulted in convergent morphological characteristics in the ophiostomatoid fungi, which has made it difficult to establish reliable taxonomic characters for this group of fungi. Thus, the classification of these fungi has been the subject of substantial debate since the first ophiostomatoid species was described in 1890. In recent years, phylogenetic analyses using DNA sequence data have provided new clarity to classification of these fungi and have facilitated more accurate identification of them.

The genus *Ceratocystis sensu stricto* is reserved for those ophiostomatoid fungi that have *Thielaviopsis* anamorphs, that are intolerant to the antibiotic cycloheximide in culture and that lack cellulose and rhamnose in their cell walls. Some of these species are economically important plant pathogens of both agricultural crops and forest plantation trees. As species of *Ceratocystis* have been more intensively studied, it has become evident that some of the species in fact represent a complex of several host-specialized forms and cryptic taxa. This has important implications, especially in designing for quarantine measures against these pathogens.

Nine of the eleven species thus far identified in the *C. coerulescens* complex have been unequivocally delimited using a wide variety of

phenotypic and genotypic characteristics. Nevertheless, problems have been experienced in separate *C. polonica* and *C. laricicola*. This is because they are morphologically indistinguishable, have similar isozyme profiles and identical ITS sequences. Both species are associated with aggressive conifer bark beetles and are able to kill their hosts, which justifies a more intensive study of them. Improved taxonomy of these species and an extended knowledge regarding their population characteristics would expand our understanding of their epidemiological relationships with insect vectors and host trees.

Ceratocystis fimbriata represents one of the most virulent and economically important vascular pathogens in many agricultural crops and forest trees. Genetic and pathological studies have suggested that this fungus comprises a complex of several host-specialized forms. Two of these "forms" have already been described as new species based on molecular and detailed morphological comparisons. It is clear that further investigations are required to delimitate other species within this complex of important plant pathogens.

Much literature has been generated for *Ceratocystis sensu stricto*. During the last decade, research dealing with these fungi was especially focused in understanding the phylogenetic relationships amongst *Ceratocystis* species. However, a few studies deal

with the population biology of *Ceratocystis* species. The knowledge of the level of variability and degree of genetic structure of natural populations of plant pathogens is important to establish hypothesis regarding their origin, movement and reproduction strategies. This information is likely to be required in the planning and implementation of management strategies against diseases caused by species of *Ceratocystis*, including generation of resistant plant material, development of quarantine strategies and appropriate use of chemical control measures.

ACKNOWLEDGEMENTS

We wish to express our gratitude to Prof. Brenda Wingfield, Dr. Oliver Preisig and Dr. Thomas Kirisits for their critic review of this manuscript. Financial support was provided by the National Research Foundation (NRF) of South Africa, the University of Pretoria, the Forestry and Agricultural Biotechnology Institute (FABI), the members of the Tree Pathology Cooperative Program (TPCP), the National Center of Coffee Research (Cenicafé, Colombia) and the Institute of Forest Entomology, Forest Pathology and Forest Protection, Universität für Bodenkultur Wien (IFFF-BOKU), Austria.

REFERENCES

- Accordi, S. M. 1986. Diffusione de *Ceratocystis fimbriata* f. *platani* attraverso le anastomosi radicali. In: Informatore Fitopatologico. Vol. 36, no. 11; p. 53-58.
- _____. 1989. Sopravivenza nel terreno di *Ceratocystis fimbriata* f. sp. *platani*. In: Informatore Fitopatologico. Vol. 39, no. 5; p. 57-62.
- Appel, D. N., Kurdyla, T. and Lewis, R., Jr. 1990. Nitidulids as vectors of the oak wilt fungus and other *Ceratocystis* spp. in Texas. In: European Journal of Forest Pathology. Vol. 20; p. 412-417.
- Baier, P., Führer, E., Kirisits, T. and Rosner, S. 2002. Defence reactions of Norway spruce against bark beetles and the associated fungus *Ceratocystis polonica* in secondary pure and mixed species stands. In: Forest Ecology and Management. Vol. 159; p. 73-86.
- Baker, K. F. and Thomas, E. H. 1946. Failure of bud and graft unions of rose induced by *Chalaropsis thielavioides*. In: Phytopathology. Vol. 36; p. 281-291.
- _____ and Harrington, T. C. 2000. Host-specialized *Ceratocystis fimbriata* on cacao, sweet potato and sycamore. In: Inoculum. Vol. 51, no. 6; p. 3-7.
- Baker, K. F. and Thomas, E. H. 2001. *Ceratocystis fimbriata*. Available in Internet: <http://www.public.iastate.edu/~tcharrin/CABIinfo.html>. [Consulted: October 2003]
- Bakshi, B. K. 1950. Fungi associated with ambrosia beetles in Great Britain. In: Transactions of the British Mycological Society. Vol. 33, no. 1-2; p. 111-120.
- Barnes, I., Gaur, A., Burgess, T., Roux, J., Wingfield, B. D. and Wingfield, M. J. 2001. Microsatellite markers reflect intraspecific relationships between isolates of the vascular wilt pathogen *Ceratocystis fimbriata*. In: Molecular Plant Pathology. Vol. 2, no. 6; p. 319-325.
- _____, Roux, J., Wingfield, B. D., Dudzinski, M. J., Old, K. M. and Wingfield, M. J. 2003. *Ceratocystis pirilliformis*, a new species from *Eucalyptus nitens* in Australia. In: Mycologia. Vol.95; no. 5; p. 865-871.
- Batista, A. C. 1960. *Ceratocystis fimbriata* Ell. & Halst. sobre *Mangifera indica* L. In: Instituto de Micologia da Universidade do Recife. p. 1-46. (Publicação no. 244).
- Benny, G. L. and Kimbrough, J. W. 1980. A synopsis of the orders and families of Plectomycetes with keys to genera. In: Mycotaxon. Vol. 12, no. 1; p. 1-91.
- Bezerra, J. L. 1997. *Ceratocystis fimbriata* causing death of budded cocoa seedlings in Bahia, Brazil. In: Incoped Newsletter. Vol. 1; p. 6.
- Bliss, D. E. 1941. A new species of *Ceratostomella* on the date palm. In: Mycologia. Vol. 33, no. 2; 468-482.

- Buriticá, P. 1999. Patógenos y enfermedades de las plantas de importancia económica en Colombia. Bogotá: ICA. 329 p.
- Butler, E. J. 1906. Fungus diseases of sugar cane in Bengal. India Department of Agriculture In: Memories, Botanical Series, no.1; p.1-53.
- Castro, B. 1991. Nuevas recomendaciones para el control de la llaga macana del cafeto. In: Avances Técnicos Cenicafé, no. 160; p. 1-4.
- _____. 1998. Evaluación del daño económico ocasionado por la llaga macana del cafeto. In: Cenicafé. Informe anual de actividades 1997-1998. Centro Nacional de Investigaciones de Café, Cenicafé, Chinchiná. 10 p.
- Chang, V. C. S. and Jensen, L. 1974. Transmission of the pineapple di-sease organism of sugarcane by nitidulid beetles in Hawaii. In: Journal of Economical Entomology. Vol. 67, no. 2; p. 190-192.
- Christiansen, E. 1985. *Ips/Ceratocystis* infection of Norway spruce: what is a deadly dosage? In: Zeitschrift Für Angewandte Entomologie. Vol. 99; p. 6-11.
- Christiansen, E. and Ericsson, A. 1986. Starch reserves in *Picea abies* in relation to defense reaction against a bark beetle transmitted blue-stain fungus, *Ceratocystis polonica*. In: Canadian Journal of Forest Research. Vol. 16; p. 78-83.
- _____. and Bakke, A. 1988. The spruce bark beetle of Eurasia. p. 479-503. In: Dynamics of forest insect populations: pattern, causes, implications. New York: Plenum Press.
- _____, Krokene, P., Berryman, A. A., Franceschi, V. R., Krekling, T., Lieutier, F., Lonneborg, A. and Solheim, H. 1999. Mechanical injury and fungal infection induce acquired resistance in Norway spruce. In: Tree Physiology. Vol. 19; p. 399-403.
- Contreras, J. and Marmeloz, L. 1984. *Ceratocystis fimbriata* Ellis & Halst., a new lemon tree pathogen. p. 432-435. In: V International Citrus Congress. (5: 1984 : São Paulo, Brazil). Proceedings. Sao Paulo: International Society of Citriculture.
- Crone, L. J. and Bachelder, S. 1961. Insect transmission of canker stain fungus, *Ceratocystis fimbriata* f. sp. *platanii*. In: Phytopathology. Vol. 51; p. 576.
- Crooke, M. and Bevan, D. 1957. Note of the first British occurrence of *Ips cembrae* Heer (Col. Scolytidae). In: Forestry. Vol. 30; p. 21-28.
- Davidson, R. W. 1935. Fungi causing stain in logs and lumber in the southern states, including five new species. In: Journal of Agricultural Research. Vol. 50; p. 789-807.
- _____. 1944. Two American hardwood species of *Endoconidiophora*

described as new. In: Mycologia. Vol. 34; p. 650-662.

_____. 1953. Two common lumber staining fungi in the western United States. In: Mycologia. Vol. 45; p. 579-586.

De Beer, C. 1994. *Ceratocystis fimbriata* with special reference to its occurrence as a pathogen of *Acacia mearnsii* in South Africa. Bloemfontein, South Africa. Thesis Magister Scientiae. University of the Orange Free State.

De Hoog, G. S. 1974. The genera *Blastobotrys*, *Sporothrix*, *Calcarisporium* and *Calcarisporiella* gen. nov. In: Studies in Mycology. Vol. 7; p. 1-84.

_____ and Scheffer R. J. 1984. *Ceratocystis* versus *Ophiostoma*: a reappraisal. In: Mycologia. Vol. 76; p. 292-299.

De Vay, J.E, Lukezic, F.L., English, H., Trujillo, E.E. and Moller, W.J. 1968. *Ceratocystis* canker of deciduous fruit trees. In: Phytopathology. Vol. 58; p. 949-954.

Delgado, J. C. and Echandi, E. 1965. Evaluación de la resistencia de especies y clones de cacao al mal del machete provocado por *Ceratocystis fimbriata*. In: Turrialba. Vol. 15; p. 286-289.

Desrosiers, R. 1958. El problema de la *Cerastomella* en el Ecuador. p. 67-70. In: Conferencia Interamericana de

cacao. Palmira, Colombia: Ministerio de Agricultura, División de Investigaciones Agropecuarias.

Echandi, E. and Segall, R. H. 1956. Trunk, branch and stem canker of coffee trees. In: Plant Disease Reporter. Vol. 40; p. 916-918.

Elliott, J. A. 1923. The ascigerous stage of the sweet potato black rot fungus. In: Phytopathology. Vol. 13; p. 56.

Evensen, P. C., Solheim, H., Høiland, K. and Stenersen, J. 2000. Induced resistance of Norway spruce, variation of phenolic compounds and their effects on fungal pathogens. In: European Journal of Forest Pa-hology. Vol.30, no. 2; p. 97-108.

Franceschi, V. R., Krekling, T. and Christiansen, E. 2002. Application of methyl jasmonate on *Picea abies* (Pinaceae) stems induces defense related responses in phloem and xylem. In: American Journal of Botany. Vol. 89, no. 1; p. 578-586.

Führer, E. 1996. Entomologische aspekte der umwandlung montaner fichtenforste. In: Entomologia Generalis. Vol. 21, no.1-2; p. 1-15.

Furniss, M. M., Solheim, H. and Christiansen, E. 1990. Transmission of blue-stain fungi by *Ips typo-graphus* (Coleoptera: Scolytidae) in Norway spruce. In: Annals of the Entomological Society of America. Vol. 83; no. 2; p. 712-716.

- Gibbs, J. N. and French, D. W. 1980. Transmission of oak wilt. In: USDA Forestry Service Research Paper NC, no. 185; 17 p.
- _____. 1993. The biology of ophiostomatoid fungi causing sapstain in trees and freshly cut logs. pp. 153-160. In: Wingfield, M.J., Seifert, K.A. and Webber J.A., eds. *Ceratocystis* and *Ophiostoma*: taxo-nomy, ecology and pathogenicity. St. Paul, Minnesota: APS Press.
- Goberdhan, L. 1959. The present situation of Ceratostomella disease of cacao in Trinidad. In: Caribbean Commonwealth Publication Exchange Service, Cocoa. Vol. 89, 4 p.
- Griffin, H. D. 1968. The genus *Ceratocystis* in Ontario. In: Canadian Journal of Botany. Vol. 46, no. 5; p. 689-718.
- Halsted, B. D. 1890. Some fungus diseases of the sweet potato. In: New Jersey Agricultural College Experiment Station Bulletin, no. 76; p. 7-14.
- Halsted, B. D. and Fairchild, D.G. 1891. Sweet-potato black rot (*Ceratocystis fimbriata*, Ell. & Halst.). In: Journal of Mycology. Vol. 7; p. 1-11.
- Hanssen, H. P. 1993. Volatile metabolites produced by species of *Ophiostoma* and *Ceratocystis*. pp. 161-172. In: Wingfield, M.J., Seifert, K.A. and Webber J.A., eds. *Ceratocystis* and *Ophiostoma*: taxo-nomy, ecology and pathogenicity. St. Paul, Minnesota: APS Press.
- Harrington, T. C. 1981. Cycloheximide sensitivity as a taxonomic character in *Ceratocystis*. In: Mycologia. Vol. 73; p. 1123-1129.
- _____. 1993a. Biology and taxonomy of fungi associated with bark beetles. p. 37-58. In: Beetle-pathogen interactions in conifer forests San Diego: Academic Press.
- _____. 1993b. Diseases of conifers caused by species of *Ophiostoma* and *Leptographium*. pp. 161-172. In: Wingfield, M.J., Seifert, K.A. and Webber J.A., eds. *Ceratocystis* and *Ophiostoma*: taxonomy, ecology and pathogenicity. St. Paul, Minnesota: APS Press.
- Harrington, T. C. Steimel, J., Wingfield, M.J. and Kile, G.A. 1996. Isozyme variation and species delimitation in the *Ceratocystis coerulescens* complex. In: Mycologia. Vol. 88; p. 104-113.
- _____ and McNew, D. L. 1997. Self-fertility and unidirectional mating-type switching in *Ceratocystis coerulescens*, a filamentous ascomycete. In: Current Genetics. Vol. 32; p. 52-59.
- Harrington, T. C. and McNew, D. L. 1998. Partial interfertility among the *Ceratocystis* species on conifers. In: Fungal Genetics and Biology. Vol. 25; p. 44-53.

- _____ and Wingfield, M. J. 1998. The *Ceratocystis* species on conifers. In: Canadian Journal of Botany. Vol. 76; p. 1446-1457.
- _____, Steimel, J. and Kile, G. 1998. Genetic variation in three *Ceratocystis* species with outcrossing, selfing and asexual reproductive strategies. In: European Journal of Forest Pathology. Vol. 28; p. 217-226.
- _____. 2000. Host specialization and speciation in the American wilt pathogen *Ceratocystis fimbriata*. In: Fitopatologia Brasileira. Vol. 25S; p. 262-263.
- _____, Pashenova, N. V., McNew, D. L., Steimel, J. and Konstantinov, M. Yu. 2002. Species delimitation and host specialization of *Cerato-cystis laricicola* and *C. polonica* to larch and spruce. In: Plant Disease. Vol. 86, no. 4; p. 418-422.
- Hausner, G., Reid, J. and Klassen, G. R. 1993a. On the phylogeny of *Ophiostoma*, *Ceratocystis* s.s. and *Microascus*, and relationships within *Ophiostoma* based on partial ribosomal DNA sequences. In: Canadian Journal of Botany. Vol. 71; p. 1249-1265.
- Hausner, G., Reid, J. and Klassen, G. R. 1993b. On the subdivision of *Ceratocystis* s.l., based on partial ribosomal DNA sequences. In: Canadian Journal of Botany. Vol. 71; p. 52-63.
- _____, Reid, J. and Klassen, R. 2000. On the phylogeny of members of *Ceratocystis* s.s. and *Ophiostoma* that possess different anamorphic states, with emphasis on the anamorph genus *Leptographium*, based on partial ribosomal DNA sequences. In: Canadian Journal of Botany. Vol. 78; p. 903-916.
- Havord, G. 1962. Problemas esenciales en las investigaciones en cultivos perennes. 6 p. In: Seminario sobre Diseños Estadísticos y Técnicas Experimentales con Cultivos Perennes. (1962: Turrialba, Costa Rica). Turrialba, Costa Rica: IICA.
- Henry, B. W., Moses, C.S., Richards, C.A. and Riker, A.J. 1944. Oak wilt, its significance, symptoms and cause. In: Phytopathology. Vol. 34; p. 636-647.
- Hinds, T. E. 1972. Insect transmission of *Ceratocystis* species associated with aspen cankers. In: Phytopathology. Vol. 62; p. 221-225.
- _____ and Davidson, R.W. 1972. *Ceratocystis* species associated with aspen cankers. Phytopathology. Vol. 62; p. 221-225.
- Hunt, J. 1956. Taxonomy of the genus *Ceratocystis*. In: Lloydia. Vol.19; p.1-58.
- Idrobo, M. S. 1958. El complejo *Xyleborus-Ceratostomella* en Colombia. p. 73-79. In: Conferencia Interamericana de Cacao. Ministerio de Agricultura, División de Investigaciones Agropecuarias, Palmira, Colombia.

Instituto Colombiano Agropecuario, ICA. 1993. La enfermedad pudrición basal de los cítricos. pp. 1-5. In: Programa de Frutales ICA. Acta de reunión sobre la enfermedad conocida como pudrición basal de los cítricos. Palmira, Colombia: ICA.

Jacobs, K. and Wingfield, M. J. 2001. *Leptographium* species: tree pathogens, insects associates, and agents of blue-stain. St. Paul, Minnesota, USA: APS Press.

Jewell, T. R. 1974. A qualitative study of cellulose distribution in *Ceratocystis* and *Europhium*. In: Mycologia. Vol. 66; p. 139-146.

Juzwik, J. and French, D. W. 1983. *Ceratocystis fagacearum* and *C. piceae* on the surfaces of free-flying and fungus-mat-inhabiting nitidulids. In: Phytopathology. Vol. 73; p. 1164-1168.

Kile, G. A. and Walker, J. 1987. *Chalara australis* sp. nov. (Hyphomycetes), a vascular pathogen of *Nothofagus cunninghamii* (Fagaceae) in Australia and its relationship to other *Chalara* species. In: Australian Journal of Botany. Vol. 35; p. 1-32.

_____. 1993. Plant diseases caused by species of *Ceratocystis sensu stricto* and *Chalara*. pp. 173-183. In: Wingfield, M. J., Seifert, K. A. and Webber J. A., eds. *Ceratocystis* and *Ophiostoma*: taxonomy, ecology and pathogenicity. St. Paul, Minnesota: APS Press.

Kile, G. A., Harrington, T. C., Yuan, Z. Q., Dudzinski, M. J. and Old, K. M. 1996. *Ceratocystis eucalypti* sp. nov., a vascular stain fungus from eucalypts in Australia. In: Mycological Research. Vol.100; p. 571-579.

Kirisits, T. 1998. Pathogenicity of three blue-stain fungi associated with the bark beetle *Ips typographus* to Norway spruce in Austria. In: Österreichische Zeitschrift für Pilzkunde. Vol. 7; p. 191-201.

Kirisits, T. and Anglberger, H. 1999. Report on a strain of the pathogenic blue-stain fungus *Ceratocystis polonica* with low virulence. In: Österreichische Zeitschrift für Pilzkunde. Vol. 8; p. 157-167.

_____. 2001. Studies on the association of ophiostomatoid fungi with bark beetles in Austria with special emphasis on *Ips typographus* and *Ips cembrae* and their associated fungi *Ceratocystis polonica* and *Ceratocystis laricicola*. Vienna, Austria, 69 h. *Rerum Naturalium Technicarum Doc-tor Thesis*. Universität für Bodenkultur Wien (BOKU). Institute of Forest Entomology, Forest Pathology and Forest Protection.

Kirisits, T. and Offenthaler I. 2002. Xylem sap flow of Norway spruce after inoculation with the blue-stain fungus *Ceratocystis polonica*. In: Plant Pathology. Vol. 51; 359-364.

- Klimetzek, K. D. and Yue, C. 1997. Climate and forest insect outbreaks. In: *Biologia*, Bratislava. Vol. 52, no. 2; p. 153-157.
- Kojima, M. and Uritani, I. 1976. Possible involvement of furanoterpenoid phytoalexins in establishing host-parasite specificity between sweet potato and various strains of *Ceratocystis fimbriata*. In: *Physiological Plant Pathology*. Vol. 8, no. 1; p. 97-111.
- Krokene, P. 1996. The role of blue-stain fungi in tree-killing by bark beetles. Oslo. 116 h. Doctor Science thesis. University of Oslo. Division of Zoology, Department of Biology.
- _____ and Solheim, H. 1996. Fungal associates of five bark beetle species colonizing Norway spruce. In: *Canadian Journal of Forest Research*. Vol. 26; p. 2115-2122.
- _____ and _____ 1998. Phytopathogenicity of four blue-stain fungi associated with aggressive and nonaggressive bark beetles. In: *Phytopathology*. Vol. 88; p. 39-44.
- Krokene, P., Christiansen, E., Solheim, H., Franceschi, V.R. and Berryman, A.A. 1999. Induced resistance to pathogenic fungi in Norway spruce. In: *Plant Physiology*. Vol. 121; p. 565-569.
- _____ and Solheim, H. 2001. Loss of pathogenicity in the blue-stain fungus *Ceratocystis polonica*. In: *Plant Pathology*. Vol. 50; p. 497-502.
- _____, Solheim, H. and Christiansen, E. 2001. Induction of disease resistance in Norway spruce (*Picea abies*) by necrotizing fungi. In: *Plant Pathology*. Vol. 50; p., 230-233.
- _____, Solheim, H. and Christiansen, E. 2003. Inducible anatomical defense responses in Norway spruce stems and their possible role in induced resistance. In: *Tree Physiology*. Vol. 23, no. 3; p. 191-197.
- Leather, R. I. 1966. A canker and wilt disease of pimento (*Pimenta officinalis*) caused by *Ceratocystis fimbriata* in Jamaica. In: *Transactions of the British Mycological Society*. Vol. 49; p. 213-218.
- Malaguti, G. 1952. *Ceratostomella fimbriata* en el cacao de Venezuela. In: *Acta Científica Venezolana*. Vol. 3; p. 94-97.
- Malloch, D. and Blackwell, M. 1993. Dispersal biology of the ophiostomatoid fungi. p. 195-206. In: Wingfield, M.J., Seifert, K.A. and Webber J.A., eds *Ceratocystis and Ophiostoma: taxonomy, ecology and pathogenicity*. St. Paul, Minnesota: APS Press.
- Marais, G. J., Wingfield, M. J., Viljoen, C. D. and Wingfield, B. D. 1998. A new ophiostomatoid genus

- from *Protea* infructescences. In: Mycologia. Vol. 90; p. 136-141.
- Matasci, M. and Gessler, C. 1997. Ein pilz bedroht die existenz der platane. In: Acta Veterinaria Hungarita. Vol. 45; p. 69-75.
- Minter, D. W., Kirk, P. M. and Sutton, B. C. 1983. Thalic phialides. In: Transactions of the British Mycological Society. Vol. 80; p. 39-66.
- _____, _____ and Sutton, B. C. 1982. Holoblastic phialides. In: Transactions of the British Mycological Society. Vol. 79; p. 75-93.
- Minter, D. W. 1987. The significance of conidiogenesis in pleoanamorphy. p. 241-262. In: Pleomorphic fungi. Tokyo, Amsterdam: Elsevier.
- Montes de Oca, H. A. 1975. Estudio de la resistencia a *Ceratocystis fimbriata* Ellis & Halst., en híbridos y clones de cacao (*Theobroma cacao* L.), utilizando dos métodos de evaluación. Turrialba. Thesis Magister Scientiae. Instituto Interamericano de Ciencias Agrícolas de la OEA, Costa Rica.
- Morris, M. J., Wingfield, M. J. and De Beer, C. 1993. Gummosis and wilt of *Acacia mearnsii* in South Africa caused by *Ceratocystis fimbriata*. In: Plant Pathology. Vol. 42; p. 814-817.
- Mourichon, X. 1994. Serious citrus dieback in Colombia caused by *Ceratocystis fimbriata*. In: Fruits. Vol. 49, no. 5/6; p. 415-416.
- Münch, E. 1907. Die Blaufaule des nadelholzes. I-II. In: Naturwissenschaftliche Zeitschrift für Forst- und Landwirtschaft. Vol. 5; p. 531-573.
- Nag Raj, T. R. and Kendrick, W. B. 1975. A monograph of *Chalara* and allied genera. Waterloo: Wilfred Laurier University Press. 200 p.
- Nagy, N. E., Franceschi, V. R., Solheim, H., Krekling, T. and Christiansen, E. 2000. Wound-induced traumatic resin duct development in stems of Norway spruce (Pinaceae): anatomy and cytochemical traits. In: American Journal of Botany. Vol. 87; p. 302-313.
- Nannfeldt, J. A. 1932. Studien über die morphologie und Systematik der nichtlichenisierten inoperculaten Dyscomyceten. In: Nova Acta Regiae Societatis Scientiarum Upsaliensis. Ser. 4. Vol. 8; p. 1-368.
- Nuss, D. L. 1992. Biological control of chestnut blight: an example of virus-mediated attenuation of fungal pathogenesis. In: Microbiological Reviews. Vol. 56, no. 4; p. 561-576.
- Nuss, D. L. 2000. Hypovirulence and chestnut blight: from the field to the laboratory and back. p. 149-170. In: Fungal pathology. Netherlands: Kluwer Academic Publishers.

- Oregon, R. 1936. Un amarillamiento del cafeto. In: Boletín Agrícola de la Sociedad Antioqueña de Agricultura, no. 9; 725-732.
- Ocampo, R. F., Mafla, B. G. and Victoria, K. J. I. 1982. Mejoramiento genético para resistencia a *Ceratocystis fimbriata* en cacao (*Theobroma cacao* L.). p. 725-730. In: VIII International Cocoa Research Conference. (1982: Cartagena, Colombia). Proceedings. Cartagena: Cocoa Producers' Alliance, Cartagena, Colombia.
- Okada, G., Seifert, K. A., Takematsu, A., Yamaoka, Y., Miyazaki, S. and Tubaki, K. 1998. A molecular phylogenetic reappraisal of the *Graphium* complex based on 18S rDNA sequences. In: Canadian Journal of Botany. Vol.76; p. 1495-1506.
- Olchowecki, A. and Reid, J. 1974. Taxonomy of the genus *Ceratocystis* in Manitoba. In: Canadian Journal of Botany. Vol. 52; p. 1675-1711.
- Olsen, E. O. and Martin, W. J. 1949. Relationship of *Ceratostomella fimbriata* from *Hevea* rubber tree and sweet potato. In: Phytopathology. Vol. 39; p.17-30.
- Paine, T. D., Raffa, K. and Harrington, T. C. 1997. Interactions among scolytid bark beetles, their associated fungi, and live host conifers. In: Annual Review of Entomology. Vol. 42; p. 179-206.
- Panconesi, A. 1981. *Ceratocystis fimbriata* of plane trees in Italy: biological aspects and control possibility. In: European Journal of Forest Pathology. Vol. 11, no. 1; p. 385-395.
- Pardo-Cardona, V. M. 1995. Hongos fitopatógenos de Colombia. Medellín: Universidad Nacional de Colombia.
- Paulin-Mahady, A. E., Harrington, T. C. and McNew, D. 2002. Phylogenetic and taxonomic evaluation of *Chalara*, *Chalaropsis*, and *Thielaviopsis* anamorphs associated with *Ceratocystis*. In: Mycologia. Vol. 94; p. 62-72.
- Peplinski, J. D. and Merrill, W. 1974. Nonsurvival of *Ceratocystis fagacearum* in frass of oak bark beetles and ambrosia beetles. In: Phytopathology. Vol. 64; p. 1528-1530.
- Pfeffer, A. 1995. Zentral- und Westpaläarktische Borken- und Kernkäfer. Basel: c/o aturhistorisches Museum Basel. 310 p.
- Pontis, R. E. 1951. A canker disease of the coffee tree in Colombia and Venezuela. In: Phytopathology. Vol. 41; p. 179-184.
- Postner, M. 1974. Scolytidae (Ipidae) Borkenkäfer. In: Schwenke, W., ed. Die Forstschädlinge Europas. Hamburg, Berlin: Paul Parey Verlag.

- Redfern, D. B., Stoakley, J.T., Steele, H. and Minter, D.W. 1987. Dieback and death of larch caused by *Ceratocystis laricicola* sp. nov. following attack by *Ips cembrae*. In: Plant Pathology Vol. 36; p. 467-480.
- Rogers, H. J., Buck, K. W. and Braiser, C. M. 1987. A mitochondrial target for double-stranded RNA in diseased isolates of the fungus that causes Dutch elm disease. In: Nature. Vol. 329; p. 558-560.
- Rossetto, C. J. and Ribeiro, I. J. A. 1990. Seca da mangueira. Part 12: Recomendacoes de controle. In: Revista de Agricultura. Vol. 65; p. 173-180.
- Roux, J., Wingfield, M. J., Wingfield, B. D., Bouillet, J. P. and Alfenas, A. C. 1999. A serious new disease of *Eucalyptus* caused by *Ceratocystis fimbriata* in Central Africa. In: Forest Pathology. Vol.30, 175-184.
- _____, Coutinho, T. A., Mujuni Byabashaija, D. and Wingfield, M. J. 2001. Diseases of plantation *Eucalyptus* in Uganda. In: South African journal of Science. Vol. 97, no.1/2; p. 16-18.
- Saccardo, P. A. 1892. Em: *Sylloge fungorum*. Vol.10; p. 215-216.
- Santini, A. and Capretti, P. 2000. Analysis of the Italian population of *Ceratocystis fimbriata* f. sp. *platani* using RAPD and minisatellite markers. In: Plant Pathology. Vol. 49; p. 461-467.
- Schieber, E. 1969. Enfermedad del cacao "mal de machete" provocada por *Ceratocystis fimbriata* en la Republica Dominicana. In: Turrialba. Vol. 19, no. 3; p. 340-344.
- Schimitschek, E. 1931. Der achtzählige Lärchenborkenkäfer *Ips cembrae* Heer – Zur Kenntnis seiner Biologie und Ökologie sowie seines Lebensvereins. In: Zeitschrift für Angewandte Entomologie. Vol. 17; p. 253-344.
- Siemaszko, W. 1939. Zespoly grzybów towarzyszących kornikom polskim. In: Planta Polonica. Vol. 7; p. 1-54.
- Silveira, A., Cardoso, R., Brignani, N., Ortolani, A. and Godoy, G. 1994. Caracterizacao do prejuizo provocado pelo mofo cinzento (*Ceratocystis fimbriata*) em painés de seringueira (*Hevea brasiliensis*). In: Summa Phytopathologica. Vol. 20; p. 196-199.
- Simmonds, N. W. 1994. Horizontal resistance to cocoa diseases. In: Cocoa Growers' Bulletin no. 47; p. 42-52.
- Solheim, H. 1986. Species of Ophiostomataceae isolated from *Picea abies* infested by the bark beetle *Ips typographus*. In: Nordic Journal of Botany. Vol. 6; p. 199-207.

- Solheim, H. 1988. Pathogenicity of some *Ips typographus*-associated blue-stain fungi to Norway spruce. In: Meddelelser fra Norsk Institutt for Skogforskning. Vol. 40; p. 1-11.
- Solheim, H. 1991. Oxygen deficiency and spruce resin inhibition of growth of blue stain fungi associated with *Ips typographus*. In: Mycological Research. Vol. 95; p. 1387-1392.
- _____. 1992. Fungal succession in sapwood of Norway spruce infested by the bark beetle *Ips typographus*. In: European Journal of Forest Pathology. Vol. 22; p. 136-148.
- Solheim, H. and Safranyik, L. 1997. Pathogenicity of the spruce beetle associated blue-stain fungi, *Ceratocystis rufipenni* and *Leptographium abietinum* to Sitka spruce. In: Canadian Journal of Forest Research. Vol. 27; p. 1336-1341.
- Spatofora, J. W. and Blackwell, M. 1994. The polyphyletic origins of ophiostomatoid fungi. In: Mycological Research. Vol. 98; p. 1-9.
- Spencer, J. F. T. and Gorin, P. A. J. 1971. Systematics of the genera *Ceratocystis* and *Graphium*. Proton magnetic resonance spectra of the mannose-containing polysaccharides as an aid in classification. In: Mycologia. Vol. 63; p. 387-402.
- Stauffer, C., Kirisits, T., Nussbaumer, C., Pavlin, R. and Wingfield, M. J. 2001. Phylogenetic relationships between the European and Asian eight spined larch bark beetle populations (Coleoptera, Scolytidae) inferred from DNA sequences and fungal associates. In: European Journal of Entomology. Vol. 98; p. 99-105.
- Summerbell, R. C., Kane, J., Krajden, S. and Duke, E. E. 1993. Medically important *Sporothrix* species and related ophiostomatoid fungi. pp. 185-192. In: Wingfield, M. J., Seifert, K. A. and Webber J. A., eds. *Ceratocystis* and *Ophiostoma*: taxonomy, ecology and pathogenicity. St. Paul, Minnesota: APS.
- Szkolnik, M. 1951. Coffee trunk and stem canker in Guatemala. In: Plant Disease Reporter. Vol. 35; p. 500-501.
- Teviotdale, B. L. and Harper, D. H. 1991. Infection of pruning and small bark wounds in almond by *Ceratocystis fimbriata*. In: Plant Disease. Vol. 75; p. 1026-1030.
- Upadhyay, H. P. 1993. Classification of the ophiostomatoid fungi. p. 7-13. In: Wingfield, M. J., Seifert, K. A. and Webber J. A., eds. *Ceratocystis* and *Ophiostoma*: taxonomy, ecology and pathogenicity. APS Press, St. Paul, Minnesota.
- _____. 1981. A monograph of *Ceratocystis* and *Ceratocystiopsis*. Athens, USA: University of Georgia.
- _____ and Kendrick, W. B. 1975. Prodrromus for a revision of *Cera-*

- tocystis* (Microascales, Ascomycetes) and its conidial states. In: Mycologia. Vol. 67; p.798-805.
- Vigouroux, A. and Stojadinovic, B. 1990. Possibilities d'infection du platane pour *Ceratocystis fimbriata* f. *platani* après contamination de l'eau ou se développent des recines blesses. In: European Journal of Forest Pathology. Vol. 17; p. 181-184.
- Viiri, H. 2002. Interactions between host trees and fungi associated with the spruce bark beetle (*Ips typographus*). Suonenjoki, 48 h. Philosophiae Doctor Thesis. University of Joensuu, Suonenjoki, Finland. The Finnish Forest Research Institute.
- Viljoen, C. D., Wingfield, B. D. and Wingfield, M. J. 2000. Computer aided systematic evaluation of morphological characters of the ophiostomatoid fungi. In: Mycotaxon. Vol. 74; p. 217-239.
- Visser, C., Wingfield, M. J., Wingfield, B. D. and Yamaoka, Y. 1995. *Ophiostoma polonicum* is a species of *Ceratocystis sensu stricto*. In: Systematic and Applied Microbiology. Vol. 18; p. 403-409.
- Vogelzang, B. K. and Scott, E. S. 1991. *Ceratocystis fimbriata*, causal agent of basal rot of *Syngonium* cultivars, and host range studies of isolates of *C. fimbriata* in Australia. In: Australasian Plant Pathology. Vol. 19, 82-89.
- Von Arx, J. A. 1974. The genera of fungi sporulating in pure culture. 2ed. Cramer, Vaduz. 315 p.
- _____ and van der Walt, J. P. 1988. Ophiostomatales and Endomycetales. pp. 167-176. In: The expanding realm of yeast-like fungi. Proceedings of an International Symposium on the Perspectives of Taxonomy, Ecology and Phylogeny of Yeast and Yeast-like Fungi. The Netherlands: Elsevier Science.
- Walker, J.M., Tesoriero, L., Pascoe, I. and Forsberg, L. I. 1988. Basal rot of *Syngonium* cultivars and the first record of *Ceratocystis fimbriata* from Australia. In: Australasian Plant Pathology. Vol. 17; p. 22-23.
- Walter, J. M. 1946. Canker stain of plane trees. In: USDA Circular No. 742.
- _____, Rex, E. G. and Schreiber, R. 1952. The rate of progress and destructiveness of canker stain of plane trees. In: Phytopathology. Vol. 42; p. 236-239.
- Webster, R. and Butler, E. 1967. A morphological and biological concept of the species *Ceratocystis fimbriata*. In: Canadian Journal of Botany. Vol. 45; p. 1457-1468.
- Weijman, A. C. M. and De Hoog, G. S. 1975. On the subdivision of the genus *Ceratocystis*. In: Antonie van Leeuwenhoek. Vol. 41; p. 353-360.

Whitney, H. S. 1982. Relationships between bark beetles and symbiotic organisms. p. 183-211. In: Bark beetles in North American conifers. Austin, USA: University of Texas Press.

Wingfield, B. D., Grant, W. S. Wolfaardt, J. F. and Wingfield, M. J. 1994. Ribosomal RNA sequence phylogeny is not congruent with ascospore morphology among species in *Ceratocystis sensu stricto*. In: Molecular Biology and Evolution. Vol. 11; p. 376-383.

_____, Viljoen, C. D. and Wingfield, M. J. 1999. Phylogenetic relationships of ophiostomatoid fungi associated with *Protea infructescences* in South Africa. In: Mycological Research. Vol. 103, no. 12; p. 1616-1620.

Wingfield, M. J., van Wyk, P. S. and Marasas, W. F. O. 1988. *Ceratocystiopsis proteae* sp. nov., with a new anamorph genus. In: Mycologia. Vol. 80; p. 23-30.

_____. 1993. Problems delineating the genus *Ceratocystiopsis*. pp. 21-25. In: Wingfield, M.J., Seifert, K.A. and Webber J.A., eds. *Ceratocystis* and *Ophiostoma*: taxonomy, ecology and pathogenicity. St. Paul, Minnesota: APS Press,

Wingfield, M. J., Seifert, K. A. and Webber, J. F. (eds). 1993. *Ophiostoma* and *Ceratocystis*: Taxonomy,

ecology and pathogenicity. St. Paul, Minnesota, USA: APS Press.

_____, De Beer, C., Visser, C. and Wingfield, B. D. 1996. A new *Ceratocystis* species defined using morphological and ribosomal DNA sequence comparisons. In: Systematic and Applied Microbiology. Vol. 19; p. 191-202.

_____, Harrington, T. C. and Solheim, H. 1997. Two new species in the *Ceratocystis coerulescens* complex from conifers in western North America. In: Canadian Journal of Botany. Vol. 75; p. 827-834.

Witthuhn, R. C., Wingfield, B. D., Wingfield, M. J., Wolfaardt, M. and Harrington, T. C. 1998. Monophyly of the conifer species in the *Ceratocystis coerulescens* complex based on DNA sequence data. In: Mycologia. Vol. 90; p. 96-101.

Witthuhn, R. C., Wingfield, B. D., Wingfield, M. J. and Harrington, T. C. 1999. PCR-based identification and phylogeny of species of *Ceratocystis sensu stricto*. In: Mycological Research Vol. 103, no. 6; p. 743-749.

Witthuhn, R. C., Harrington, T. C., Steimel, J. P., Wingfield, B. D. and Wingfield, M. J. 2000. Comparison of isozymes, rDNA spacer regions and MAT-2 DNA sequences as phylogenetic characters in the analysis of the *Ceratocystis coerulescens complex*. In: Mycologia. Vol. 92, no. 4; p. 447-452.

Witthuhn, R. C., Wingfield, M. J., Takahashi, I. and Solheim, H. 1997. Ophiostomatoid fungi associated with the spruce bark beetle *Ips typographus* f. *japonicus* in Japan. In: Mycological Research. Vol. 101, no. 10; p. 1215-1227.

Yamaoka, Y., Wingfield, M., Ohsawa, M. and Kuroda, Y. 1998. Ophiostomatoid fungi associated with *Ips cembrae* in Japan and their pathogenicity to Japanese larch. In: Mycoscience Vol. 39, no. 1; p. 367-378.

_____, Takahashi, I. and Iguchi, K. 2000. Virulence of ophiostomatoid fungi associated with the spruce bark

beetle *Ips typographus* f. *japonicus* in Yezo spruce. In: Journal of Forest Research. Vol. 35, no. 5; p. 87-94.

Zalasky, H. 1965. Process of *Ceratocystis fimbriata* infection in Aspen. In: Canadian Journal of Botany. Vol. 43; p. 1157-1162.

Zimmerman, A. 1900. Ueber den Krebs von *Coffea arabica*, verursacht durch *Rostrella coffeae* gen. et sp. n. In: Bulletin of the Institute of Botanical Gardens Buitenzorg. Vol. 4; p. 19-22.