



LIPI

REINWARDTIA

A JOURNAL ON TAXONOMIC BOTANY, PLANT SOCIOLOGY AND ECOLOGY

ISSN 0034 – 365 X | E-ISSN 2337 – 8824 | Accredited 792/AU3/P2MI-LIPI/04/2016



2017 16 (1)

REINWARDTIA

A JOURNAL ON TAXONOMIC BOTANY, PLANT SOCIOLOGY AND ECOLOGY

Vol. 16 (1): 1 – 48, June 15, 2017

Chief Editor

Kartini Kramadibrata (Mycologist, Herbarium Bogoriense, Indonesia)

Editors

Dedy Darnaedi (Taxonomist, Herbarium Bogoriense, Indonesia)

Tukirin Partomihardjo (Ecologist, Herbarium Bogoriense, Indonesia)

Joeni Setijo Rahajoe (Ecologist, Herbarium Bogoriense, Indonesia)

Marlina Ardiyani (Taxonomist, Herbarium Bogoriense, Indonesia)

Himmah Rustiami (Taxonomist, Herbarium Bogoriense, Indonesia)

Lulut Dwi Sulistyaningsih (Taxonomist, Herbarium Bogoriense, Indonesia)

Topik Hidayat (Taxonomist, Indonesia University of Education, Indonesia)

Eizi Suzuki (Ecologist, Kagoshima University, Japan)

Jun Wen (Taxonomist, Smithsonian Natural History Museum, USA)

Barry J Conn (Taxonomist, School of Life and Environmental Sciences, The University of Sydney, Australia)

David G. Frodin (Taxonomist, Royal Botanic Gardens, Kew, United Kingdom)

Graham Eagleton (Wagstaffe, NSW, Australia)

Secretary

Rina Munazar

Layout

Liana Astuti

Illustrators

Subari

Wahyudi Santoso

Anne Kusumawaty

Correspondence on editorial matters and subscriptions for Reinwardtia should be addressed to:

HERBARIUM BOGORIENSE, BOTANY DIVISION,
RESEARCH CENTER FOR BIOLOGY– INDONESIA INSTITUTE OF SCIENCES
CIBINONG SCIENCE CENTER, JLN. RAYA JAKARTA – BOGOR KM 46,
CIBINONG 16911, P.O. Box 25 CIBINONG
INDONESIA

PHONE (+62) 21 8765066; Fax (+62) 21 8765062

E-MAIL: reinwardtia@mail.lipi.go.id

<http://e-journal.biologi.lipi.go.id/index.php/reinwardtia>

1	2	3
4	5	6
7	8	9
10	11	12

Cover images: *Catanthera keris* Veldk. (1. Inflorescences; 2. Close up flower; 3. Flower bud), *Medinilla squillula* Veldk. (4. Habit; 5. Branches; 6. Fascicle of uniflorous infructescences), *Medinilla uninervis* Veldk. (7. Habit. Note 1-nerved leaves; 8. infructescence; 9. Immature and mature fruits), *Medinilla zoster* Veldk. (10. Habit; 11. Inflorescences; 12. Flower). Photo credits: Bangun 223, Lowry & Phillipson 7287, Mahroji, Fabanyo & Soleman 69, Callmander, *et al.* 1067.

The Editors would like to thank all reviewers of volume 16(1):

Agus Susatya - University of Bengkulu, Bengkulu, Indonesia
Agus Sutanto - Indonesian Tropical Fruit Research Institute (ITFRI), West Sumatra, Indonesia
Axel D. Poulsen - Royal Botanic Garden Edinburgh, Edinburgh, Scotland, UK
Andrew Powling - School of Biological Sciences, University of Portsmouth, Portsmouth, UK
Elham Sumarga - School of Life Sciences & Technology, Institut Teknologi Bandung, Bandung, Indonesia
Meekiong Kallu - University Malaysia Sarawak, Samarahan, Sarawak, Malaysia
Harry Wiriadinata - Herbarium Bogoriense, Indonesian Institute of Sciences, Bogor, Indonesia
Ulrich Meve - Lehrstuhl für Pflanzensystematik, Universität Bayreuth, Bayreuth, Germany
Mien A. Rifai - Akademi Ilmu Pengetahuan Indonesia (API), Jakarta, Indonesia

NEW SPECIES OF *CATANATHERA* AND *MEDINILLA* (MELASTOMATACEAE) FROM HALMAHERA, INDONESIA AND A NEW NAME FOR A *MEDINILLA* FROM MADAGASCAR

Received January 12, 2017; accepted May 04, 2017

JAN-FRITS VELDKAMP

Naturalis Biodiversity Center, Herbarium, PO Box 9517, 2300 RA Leiden, the Netherlands. Email: jef.veldkamp@naturalis.nl

ABDULROKHMAN KARTONEGORO

Naturalis Biodiversity Center, Herbarium, PO Box 9517, 2300 RA Leiden, the Netherlands. Email: abdulrokhman.kartonegoro@naturalis.nl
Research Center for Biology-LIPI, Jl. Raya Jakarta-Bogor-KM. 46, Cibinong, 16911, Indonesia

ABSTRACT

VELDKAMP, J. F. & KARTONEGORO, A. 2017. New species of *Catanthera* and *Medinilla* (Melastomataceae) from Halmahera, Indonesia and a new name for a *Medinilla* from Madagascar. *Reinwardtia* 16(1): 25 – 30. — One new species of *Catanthera* and four of *Medinilla* (Melastomataceae) from Halmahera, Moluccas, Indonesia, are described. One also occurs in Morotai. A new name was needed for *M. intermedia* H. Perrier (1932), non Blume (1831), from Madagascar.

Key words: Anisomorphous anthers, Moluccas, Morotai.

ABSTRAK

VELDKAMP, J. F. & KARTONEGORO, A. 2017. Jenis-jenis baru *Catanthera* dan *Medinilla* (Melastomataceae) dari Halmahera, Indonesia dan satu nama baru *Medinilla* dari Madagascar. *Reinwardtia* 16(1): 25 – 30. — Dipertelakan satu jenis baru *Catanthera* dan empat jenis *Medinilla* (Melastomataceae) dari Halmahera, Maluku, Indonesia serta satu jenis tersebar sampai ke Morotai. Satu nama baru diperlukan untuk menampung *M. intermedia* H. Perrier (1932), non Blume (1831), dari Madagascar.

Kata kunci: Anter anisomorfus, Maluku, Morotai.

INTRODUCTION

During a survey of the PT Weda Bay Nickel for their Flora Inventory Programme on Halmahera, Moluccas, Indonesia, many interesting species were collected. Here a summary of some Halmahera Melastomataceae is presented. *Catanthera* F. Muell. and *Medinilla* Gaudich. are new generic records for the island.

CATANATHERA F. Muell.

In his revision of *Catanthera* F. Muell. (Melastomataceae) Nayar (1982) gave a map that showed a disjunction in the distribution of the genus between western (Sumatra, Borneo) and eastern (New Guinea) Malesia, the genus apparently being absent in Celebes, the Moluccas and the Philippines. In the meanwhile collections have come in from Celebes, Morotai and Halmahera (Moluccas) and Palawan (Philippines).

In the Dissochaetae there are two whorls of stamens, an outer, alternipetalous one, where except for *Diplectria* (Blume) Rchb., the anthers are always fertile and in case of heteromorphy are the longest ones, and an inner, epipetalous one, which may have smaller anthers, be staminodial, or even

absent. In *Diplectria* the situation is reversed: the alternipetalous stamens are staminodial and the epipetalous are fertile. All species of *Catanthera* seen follow the usual mode, but the reverse one is present in the Halmahera collection (Bangun *et al.* 223). There the alternipetalous anthers are still the longest, but they are strap-shaped and staminodial, while the epipetalous although much more compact are fertile.

Catanthera keris Veldk., *spec. nov.* Fig. 1

Petioles 8 – 10 mm long, furfureous. *Leaf* blades elliptic, 5-plinerved, second pair of basal nerves branching off 2 – 3 mm above the basal ones, underneath between the nerves sparsely puberulous. *Inflorescence* cymose, furfureous. *Hypanthium* in flower 6 – 7 mm wide, shortly furfureous. Mature floral buds *ca.* 12 mm long. Petals ovate. Anthers very heteromorph. Alternipetalous anthers sterile, strap-shaped, *ca.* 11 mm long, plectrum *ca.* 0.7 mm long, appendages *ca.* 0.7 mm long. Epipetalous anthers fertile, straight, swollen, hemi-circular in side view, 6 – 6.5 mm long. — Type: *Bangun, Andriamahefarivo, Razakamalala & Mahroji* 223. (Holotype: L!, Isotype: BO!, MO).



Fig. 1. *Catanthera keris* Veldk. A. Inflorescences; B. Close up flower; C. Flower bud (Bangun 223, <http://www.tropicos.org/Specimen/100550447?projectid=55>)

Innovations furfuraceous. Leaf blades 4.5 – 4.7 by 2.3 – 3 cm wide, 1.5 – 2 times as long as wide, base rounded to truncate, apex acute or acuminate, second pair of basal nerves branching off 2–3 mm above the basal ones. Inflorescences fascicled, 2 – 10-flowered, 3.5 – 5 cm long, furfuraceous. Pedicel 10 – 20 mm long. Hypanthium campanulate to urceolate, in flower 5.5 – 6.5 by 6 – 7 mm wide, margin truncate, shortly furfuraceous. Ovary apex glabrous. Extra-ovarian chambers to the base of the ovary. Mature floral buds *ca.* 12 mm long. Petals ovate, *ca.* 12 mm long, pink. Anthers eight. Alternipetalous anthers stipitate, *ca.* 11 mm long. Epipetalous anthers filament glabrous, anthers cinnamon (*i.s.*); plectrum lanceolate, 1.5 – 2 mm long; lateral appendages auriculiform, *ca.* 0.6 mm long. Style glabrous.

Distribution. Indonesia, Moluccas, Halmahera, Weda Bay, Bukit Limber 00° 32' 41" N, 127° 58' 32" E. Only known from the type.

Habitat. Degraded primary forest, open place, 860 m alt.

Etymology. The epithet has been derived from the sterile anthers, which to a fertile imagination resemble this Malesian weapon.

Notes. Remarkable for the very anisomorphic stamens, the longer ones strap-shaped with patent plectrum and appendages. There is a collection from Morotai, G. Pare-Pare (Kostermans 1243, BO, L) which is very similar, but the L specimen lacks petals and stamens.

Phenology. Flowering in December.

Collector's notes. Liana, 10 m tall. Stem grey. Leaves glaucous golden beneath. Peduncle pink. Hypanthium pale green and glaucous golden.

Petals, stigma pink. Stamens [prob. staminodes] white, striped yellow. Style white.

MEDINILLA Gaudich.

Medinilla in Malesia has an estimated number of *ca.* 215 species, many of which, especially in Celebes, the Lesser Sunda Isles, the Moluccas and New Guinea, are still undescribed (Bodegom & Veldkamp, 2001). Most species have a limited distribution. For the Moluccas 12 species have been recorded, but none yet for Halmahera (Bakhuizen f., 1943).

The author of the generic name is Gaudichaud (1826), not De Candolle (1828) as is usually cited (Bodegom & Veldkamp, 2001: 534–537).

Remarkably, strongly anisomorphic anthers have not been described for the genus, perhaps because the authors have neglected their presence. A survey is thus much needed, which was beyond the scope of this paper.

1. *Medinilla pellita* Veldk. & Karton., *spec. nov.* – Fig. 2.

MEDINILLA VERRUCIFERA Ohwi, ined.

Shrub (terrestrial ?). Innovations glabrous (n.v.). Branchlets terete, brown, with verrucose lenticels, and many black, branched adventitious roots, forming a kind of pelt. Leaves opposite, equal (?). Leaf axils glabrous. Petioles flattened, 0.5 – 1 cm by *ca.* 10 mm, glabrous. Leaf blades elliptic, 13 (and longer) by 6 – 12 cm wide, *ca.* 1.6 times as long as wide, *ca.* 13 (or more) times as long as the petiole, base attenuate, margin entire, glabrous, apex acute (?), underneath glabrous, submarginal vein conspicuous. Primary venation 3- or 5-plinerved, all arising from the very base; secondary nerves inconspicuous, oblique; tertiary nerves absent. Hypanthium at anthesis urceolate, *ca.* 3.5




Fig. 3. *Medinilla squillula* Veldk. A. Habit; B. Branches; C. Fascicle of uniflorous inflorescences (Lowry & Phillipson 7287, <http://www.tropicos.org/Specimen/100560752>)



Fig. 4. *Medinilla uninervis* Veldk. A. Habit. Note 1-nerved leaves; B. Infructescences; C. Immature and mature fruits. (Mahroji, Fabanyo & Soleman 69, <http://www.tropicos.org/ImageSearch.aspx>)

Branchlets cinereous, with verrucose lenticels. Leaves equal. Leaf axils glabrous. Petioles flattened and broadened at base, 5 – 6 mm wide, glabrous. Leaf blades *ca.* 2.75 times as long as wide, 3.7 – 4.25 times as long as the petiole, base cuneate, margin entire, glabrous, apex acuminate, underneath glabrous, submarginal veins inconspicuous; secondary nerves inconspicuous, oblique; tertiary nerves absent. Inflorescences fascicled, simple cymes. Bracts absent. Bracteoles present. Pedicels glabrous. Hypanthium at anthesis short cylindrical, 6 – 9 by 4.5 – 5 mm, glabrous. Extra-ovarian chambers reaching to *ca.* half the length of the hypanthium. Epipetalous anthers fertile, shrimp-shaped, 4 – 5 mm long, ventrally bullate, superior in flower at anthesis; alternipetalous anthers staminodial, strap-shaped, 7 – 10 mm long, inferior at anthesis. Dorsal spur lanceolate, 0.6 – 1 mm long; lateral appendages connate into a Y-shaped body, not recurved, 1 – 1.2 mm long. Style glabrous. Fruits not seen.

Distribution. Indonesia, Moluccas, Halmahera, Weda Bay road to Bukit Limber, 00° 33' 00" N, 127° 59' 34" E. Only known from the type.

Habitat. Epiphytic on margin of humid forest, 935 m alt.

Phenology. Flowering in January.

Etymology. The epithet refers to fertile anthers that resemble small shrimps.

Collector's notes. Small [epiphytic] shrub, *ca.* 6 m above the ground. Leaves dark green above, paler below. Petiole tinged yellow-orange. Pedicel, ovary pink-red. Filaments, style cream. Anthers yellow.

3. *Medinilla uninervis* Veldk., *spec. nov.* – Fig. 4.

Shrub glabrous. Branches terete. *Leaf* axils glabrous. Petioles 1.5 – 3 cm by 2 – 2.7 mm. Blades elliptic, 12 – 17 by 6 – 8 cm, 1.9 – 2.1 as long as wide, 1-nerved, base attenuate, apex obtuse and abruptly acuminate to nearly aristate. *Inflorescences* in axillary to cauliflorous fascicles. Flowers solitary, *peduncles* 1.5 – 3 mm long, pedicels 5–6 mm long. Flowers 5-merous. — Type:



Fig. 5. *Medinilla zoster* Veldk., A. Habit; B. Inflorescences; C. Flower (Callmänder, *et al.* 1067. <http://www.tropicos.org/Image/100231129> and <http://www.tropicos.org/Specimen/100550913>)

Mahroji, Fabanyo & Soleman 69 (Holotype: L!, Isotype: BO!, MO).

Liana. Innovations probably glabrous (not seen). Branchlets cinereous, with verrucose lenticels. Leaves in whorls of 3 or 4, equal. Petioles flattened, glabrous. Leaf blades 5 – 8 times as long as the petiole, margin entire, glabrous, underneath glabrous, submarginal veins conspicuous, 1-nerved; secondary nerves conspicuous, oblique; tertiary nerves absent. Bracts present (scars only); bracteoles not seen (early caducous?). Pedicels glabrous. Hypanthium urceolate, 8 – 10 by 6 – 8 mm, glabrous, truncate. Petals *ca.* 11 by 6 mm. Anthers anisomorph, fertile, the shorter ones *ca.* 5 mm long, dorsal spur *ca.* 1 mm long, lateral appendages free, *ca.* 1 mm long; the longer ones *ca.* 9 mm long, dorsal spur 0.5 – 1 mm long, lateral appendages fused, T- or Y-shaped, 1 – 1.3 mm long. Extra-ovarian chambers reaching to the base of the hypanthium. Fruits *ca.* 10 by 8 mm, glabrous.

Distribution. Indonesia, Moluccas, Halmahera, km 6 to Bukit Limber, 00° 30' 38" N, 128° 00' 25" E (Type); Weda Bay, Bukit Limber 00° 29' 42" N, 128° 1' 20" E (*Bangun, Haris & Mahroji 860* (BO, L, MO); without locality (*Pleyte 348*, BO, L, flower!)).

Habitat. Disturbed forest, 260 – 450 m alt.

Phenology. Flowering in September, fruiting in October.

Collector's notes. Woody liana, climbing up to 12 m. Outer bark grey, inner bark brown. Bole cream coloured, brown. Fruit cream, pale red, pink turning maroon [*i.e.* chestnut-coloured, probably the hypanthium in flower is meant; in fact the ripe fruits are dark purplish blue, "blueberry" blue].

Notes. Most peculiar is the apparent pinnate venation with a single midrib, hence the epithet. Uninerved leaves have been seen in *M. membranacea* Merr. (Luzon, the Philippines), *M. radicans*

(Blume) Blume var. *radicans* (Java and the Lesser Sunda Isles) (see below) and *M. squillula* Veldk., described here from Halmahera.

4. *Medinilla zoster* Veldk., *spec. nov.* – Fig. 5.

Innovations glabrous, branches terete. Leaves paired or in whorls of four, petiolate, obovate, widely 3-plinerved. Inflorescences axillary, later cauliflorous, peduncle well-developed, 1- or 2-(or few?)-flowered. Flowers 5- or 6-merous. Anthers anisomorphous. — Type: *Callmänder, Mahroji & Soleman 1067* (Holotype: L!, Isotype: BO!, MO).

Branchlets terete, greyish, with verrucose lenticels. Leaves equal, petiolate, axils glabrous. Petioles terete, 1.7 – 2.5 cm by 1.5 – 2 mm, glabrous. Leaf blades obovate, 5.5 – 10 by 3.5 – 8.5 cm, 1.55 – 2.4 times as long as wide, 3.4 – 5.6 times as long as petiole, base attenuate, margin entire, glabrous, apex obtuse, abruptly apiculate, underneath glabrous, submarginal veins inconspicuous, 3-plinerved, the lateral nerves arising 9 – 10 mm from base. Secondary nerves inconspicuous, oblique; tertiary nerves absent. Inflorescences axillary to cauliflorous, solitary, simple cymes, 1 – 3-flowered. Peduncle 1 – 3 cm long. Bracts and bracteoles *n.v.* Pedicels at anthesis and in fruit 9 – 15 mm long, glabrous. Hypanthium at anthesis wine-glass-shaped, 10 – 11 by 6.5 – 7 mm, glabrous, truncate. Extra-ovarian chambers reaching to the base of the hypanthium. Stamens anisomorphous, the epipetalous ones fertile, the alternipetalous ones staminodial; epipetalous anthers more or less tubular, *ca.* 8 mm long, alternipetalous anthers more or less S-shaped, apex attenuate, 10 – 11 mm long. Dorsal spur oblong, *ca.* 1 mm long (inserted 1 – 1.2 mm above the filament in the staminodes); lateral appendages free, not recurved, 1.3 – 1.5 mm long. Style with microscopic glassy hairlets. Fruits urceolate, *ca.* 12 by 10 mm, rim truncate.

Distribution. Indonesia, Moluccas, Halmahera, Weda Bay, Doromesmesan, 00° 29' 23" N, 127°

54° 30" (Type); *Gushilman et al.* 213 (BO, L, MO; frt), 00° 29'56"N, 127° 54'22"E; Morotai, Sambiki River (*Kostermans* 898; BO, L; frt).

Habitat. Open logged area in tropical lowland forest on limestone, 60–110 m alt.

Phenology. Flowering in January; fruits in May, December.

Etymology. The epithet refers to the lianoid habit, "zoster" = belt, girdle.

Collector's notes. Woody liana. Bole red. *Peduncle* green. Pedicel and calyx pink. Petals white [with a tinge of pink]. Fruit white.

Notes. *Callmander et al.*, 1067 has pinkish pedicels and fruits (fig. 10, 11), while in *Gushilman et al.*, 213 they are purple (fig. 13).

Additional notes.

During searches to identify the current collections, it was noted that *M. intermedia* Blume (1831), an endemic from Java, has a later homonym: *H. intermedia* H. Perrier (1932; 1951) from Madagascar. The two have been much confused on the internet. We here rename the latter in honour of its discoverer and describer, Joseph Marie Henri Alfred ("Henri") Perrier de la Bâthie (1873–1958), outstanding explorer of Madagascar (Dorr, 1997: 338).

Medinilla perrieri Veldk. & Karton., *nom. nov.*

Medinilla intermedia H. Perrier, *Mém. Acad. Malgache* 12: 180. 1932; in Humbert, *Fl. Madagascar* 153: 250, t. XL, f. 1–6, non Blume (1831). — Lectotype: *Perrier de la Bâthie 14181* (Holotype: P00061493; Isotype: P00061494), designated here.

MEDINILLA RADICANS (Blume) Blume

Flora 14 (1831) 509; *Rumphia* 1 (1835) 15, t. 3. — *Melastoma radicans* Blume, *Bijdr.* 17 (1826) 1069. — Type: *Anon. s.n.* (L, holo), Java, Mt. Salak, first step designation by Bakhuizen v.d. Brink f. (1943: 161), L. 908.129-113, designated here. Note that Blume's drawing of the transverse sec-

tion of the hypanthium shows five cells while elsewhere (1826, 1831, 1835) he described the flowers as "octandris (rarissime decandris)".

ACKNOWLEDGEMENTS

Dr. P. Baas (L) kindly informed me on the nature of the appendages in *M. pellita*. The photographers of the illustrations, Ms. T. J. F. Bangun, and Messrs. I. Gushilman, R. Mahroji, and P. B. Philipson are cordially thanked for their permission to use them.

REFERENCES

- BAKHUIZEN V. D. BRINK, R. C. Jr. 1943. A contribution to the knowledge of the Melastomataceae occurring in the Malay Archipelago especially in the Netherlands East Indies. *Rec. Trav. Bot. Néerl.* 40: 147–198.
- BODEGOM, S. & VELDKAMP, J. F. 2001. A revision of the pseudo-stipular species of *Medinilla* Gaud. ex DC. (Melastomataceae-Melastomatoideae-Miconieae). *Blumea* 46: 527–567.
- DORR, L. J. 1997. *Plant collectors in Madagascar and the Comoro Islands*. Royal Botanic Gardens, Kew.
- MANSFELD, R. 1925. Die Melastomataceen von Papuasien. *Bot. Jahrb. Syst.* 60: 115–130.
- MERRILL, E. D. & PERRY, L. M. 1943. Plantae papuanae archboldianae, XIII. *J. Arnold Arbor.* 24: 422–434.
- NAYAR, M. P. 1982. Revision of the genus *Catanthera* F. v. Muell. (Melastomataceae). *Reinwardtia* 10: 35–61.
- OHWI, J. 1943. The Kanehira-Hatusima 1940 collection of New Guinea plants. XVI. *Melastomataceae. Bot. Mag. (Tokyo)* 57: 6–16.
- PERRIER DE LA BÂTHIE, H. 1932. Les Mélastomacées de Madagascar. *Mém. Acad. Malgache* 12: 180.
- PERRIER DE LA BÂTHIE, H. 1951. *Mélastomacées*. In: HUMBERT, H. *Flore de Madagascar* 153 Firmin-Didit & C^o., Paris. Pp. 250–252, t. XL, f. 1–6.
- REGALADO, J. C. Jr. 1990. Revision of *Medinilla* (Melastomataceae) of Borneo. *Blumea* 35: 5–70.
- REGALADO, J. C. Jr. 1995. Revision of Philippine *Medinilla* (Melastomataceae). *Blumea* 40: 113–193.

INSTRUCTION TO AUTHORS

Scope. *Reinwardtia* is a scientific irregular journal on plant taxonomy, plant ecology and ethnobotany published in June and December. Manuscript intended for a publication should be written in English.

Titles. Titles should be brief, informative and followed by author's name and mailing address in one-paragraphed.

Abstract. English abstract followed by Indonesian abstract of not more than 250 words. Keywords should be given below each abstract.

Manuscript. Manuscript is original paper and represent an article which has not been published in any other journal or proceedings. The manuscript of no more than 36 pages by using Times New Roman 11, MS Word for Windows of A4 with double spacing, submitted to the editor through <reinwardtia@mail.lipi.go.id>. New paragraph should be indented in by 5 characters. For the style of presentation, authors should follow the latest issue of *Reinwardtia* very closely. Author(s) should send the preferred running title of the article submitted. Every manuscript will be sent to two blind reviewers.

Identification key. Taxonomic identification key should be prepared using the aligned couplet type.

Nomenclature. Strict adherence to the International Code of Nomenclature is observed, so that taxonomic and nomenclatural novelties should be clearly shown. English description for new taxon proposed should be provided and the herbaria where the type specimens area deposited should be presented. Name of taxon in taxonomic treatment should be presented in the long form that is name of taxon, author's name, year of publication, abbreviated journal or book title, volume, number and page.

Map/line drawing illustration/photograph. Map, line drawing illustration, or photograph preferably should be prepared in landscape presentation to occupy two columns. Illustration must be submitted as original art accompanying, but separated from the manuscript. The illustration should be saved in JPG or GIF format at least 350 pixels. Legends or illustration must be submitted separately at the end of the manuscript.

References. Bibliography, list of literature cited or references follow the Harvard system as the following examples.

- Journal : KRAENZLIN, F. 1913. *Cyrtandraceae* novae Philippinenses I. *Philipp. J. Sci.* 8: 163–179.
MAYER, V., MOLLER, M., PERRET, M. & WEBER, A. 2003. Phylogenetic position and generic differentiation of *Epithemateae* (*Gesneriaceae*) inferred from plastid DNA sequence data. *American J. Bot.* 90: 321–329.
- Proceedings : TEMU, S. T. 1995. Peranan tumbuhan dan ternak dalam upacara adat “Djoka Dju” pada suku Lio, Ende, Flores, Nusa Tenggara Timur. In: NASUTION, E. (Ed.). *Prosiding Seminar dan Lokakarya Nasional Etnobotani II*. LIPI & Perpustakaan Nasional: 263–268. (In Indonesian).
SIMBOLON, H. & MIRMANTO, E. 2000. Checklist of plant species in the peat swamp forests of Central Kalimantan, Indonesia. In: IWAKUMA, T. *et al.* (Eds.) *Proceedings of the International Symposium on: Tropical Peatlands*. Pp.179-190.
- Book : RIDLEY, H. N. 1923. *Flora of the Malay Peninsula 2*. L. Reeve & Co. Ltd, London.
- Part of Book : BENTHAM, G. 1876. *Gesneriaceae*. In: BENTHAM, G. & HOOKER, J. D. *Genera plantarum 2*. Lovell Reeve & Co., London. Pp. 990–1025.
- Thesis : BAIRD, L. 2002. *A Grammar of Kéo: An Austronesian language of East Nusantara*. Australian National University, Canberra. [PhD. Thesis].
- Website : <http://www.nationaalherbarium.nl/fmcollectors/k/KostermansAJGH.htm>. Accessed 15 February 2012.



Reinwardtia

Published by Herbarium Bogoriense, Botany Division, Research Center for Biology,
Indonesian Institute of Sciences

Address: Jln. Raya Jakarta-Bogor Km. 46 Cibinong 16911, P.O. Box 25 Cibinong

Telp. (+ 62) 21 8765066; Fax (+62) 21 8765062

E-mail: reinwardtia@mail.lipi.go.id

***REINWARDTIA* Author Agreement Form**

Title of article :

Name of Author(s) :

I/We hereby declare that:

- My/Our manuscript was based on my/our original work.
- It was not published or submitted to other journal for publication.
- I/we agree to publish my/our manuscript and the copyright of this article is owned by Reinwardtia.
- We have obtained written permission from copyright owners for any excerpts from copyrighted works that are included and have credited the sources in our article.

Author signature (s)

Date

Name

REINWARDTIA

Vol. 16. No. 1. 2017

CONTENTS

Page

DINI PUSPITANINGRUM, WENDY A. MUSTAQIM & MARLINA ARDIYANI. A new record of <i>Etingera pauciflora</i> (Zingiberaceae) in Java, Indonesia	1
SRI RAHAYU & MICHELE RODDA. <i>Hoya narcissiflora</i> (Apocynaceae, Asclepiadoideae), a new species from Borneo	5
IYAN ROBIANSYAH. Predicting habitat distribution of endemic and critically endangered <i>Dipterocarpus littoralis</i> in Nusakambangan, Indonesia	11
LULUT DWI SULISTYANINGSIH. A newly described and recorded infraspecific taxa of <i>Musa borneensis</i> Becc. (Musaceae) from Sulawesi, Indonesia	19
JAN-FIRTS VELDKAMP & ABDULROKHMAN KARTONEGORO. New species of <i>Catanthera</i> and <i>Medinilla</i> (Melastomataceae) from Halmahera, Indonesia and a new name for a <i>Medinilla</i> from Madagascar.....	25
PURWANINGSIH, RUDDY POLOSAKAN, RAZALI YUSUF & KUSWATA KARTAWINATA. Phytosociological study of the montane forest on the south slope of Mt. Wilis, East Java, Indonesia.....	31
IAN M. TURNER. A new combination for subspecies of <i>Radermachera quadripinnata</i> (Bignoniaceae)	47

Reinwardtia is a LIPI accredited Journal (792/AU3/P2MI-LIPI/04/2016)

<http://e-journal.biologi.lipi.go.id/index.php/reinwardtia>

Herbarium Bogoriense
Botany Division
Research Center for Biology – Indonesian Institute of Sciences
Cibinong Science Center
Jln. Raya Jakarta – Bogor, Km 46
Cibinong 16911, P.O. Box 25 Cibinong
Indonesia

