



In Silico Study of Secondary Metabolites in *Dendrobium* spp. as SARS-CoV-2 Antivirus on Main Protease (Mpro)

Anggiresti Kinasih^{1,2}, Alim El Hakim^{1,2}, Dyah Ayu Puspita Arum^{1,2}, Aulia Noor Ramadhani^{1,2}, Endang Semiarti^{1,3*}

¹Faculty of Biology, Universitas Gadjah Mada, Sleman, Jln. Teknik Selatan, Sekip Utara, Bulaksumur Yogyakarta

²Biology Orchid Study Club, Faculty of Biology Universitas Gadjah Mada, Special Region of Yogyakarta, Jln. Teknik Selatan, Sekip Utara, Bulaksumur Yogyakarta

³Laboratory of Biotechnology, Faculty of Biology, Universitas Gadjah Mada, Special Region of Yogyakarta, Jln. Teknik Selatan, Sekip Utara, Bulaksumur Yogyakarta

*Corresponding Author:

e-mail: endsemi@ugm.ac.id

Article History

Received : 25 February 2022

Revised : 14 March 2022

Approved : 25 March 2022

Published : 31 March 2022

Keywords

antiviral, docking, *Dendrobium*, Mpro, SARS-CoV-2s

ABSTRACT

Infection and deaths cases by SARS-CoV-2 still increase and have not decreased significantly. Main protease (Mpro) is playing an important role in the replication of SARS-CoV-2 life cycle and causes of rapid transmission. Natural compounds are potential to be antiviral candidates with high bioavailability and low cytotoxicity. Orchids of *Dendrobium* genus have high diversity in Indonesia. *Dendrobium* has been used as traditional Chinese medicine and contains a group of secondary metabolites with antiviral activity. This study aimed to determine the potential of secondary metabolites of *Dendrobium* orchids as antiviral candidates against Mpro SARS-CoV-2 with in silico molecular docking. Secondary metabolites obtained from the KNApSack and PubChem act as ligands. N3 inhibitors as native ligands were obtained from the RCSB. Mpro SARS-CoV-2 (6LU7) as a target macromolecule. Molecular docking was carried out using the online Covid-19 Docking Server using AutoDock Vina device. The most negative binding affinity value for each ligand compared to the native ligand binding affinity. Visualization with Discovery Studio software has been used to observe the protein amino acid residues contact for each ligand. The binding affinity of the native ligand inhibitor N3 is -7.5 kcal/mol. Based on the results of Mpro docking, three phytochemicals from *Dendrobium* spp., i.e., dendrocandin B, denthysinone, and denthysinol compounds have binding affinities of -7.7 kcal/mol, -7.9 kcal/mol, and -8.1 kcal/mol, respectively. It can be concluded that in *Dendrobium* orchid, denthysinol has the highest chance of binding so it has the potential to inhibit the Mpro SARS-CoV-2 activity.

How to cite: Kinasih, A., Hakim, A.L., Arum, D.A.P., Ramadhani, A.N., Putra, A.C.D.S & Semiarti, E. (2022). In Silico Study of Secondary Metabolites in *Dendrobium* spp. as SARS-CoV-2 Antivirus on Main Protease (Mpro). *Jurnal Riset Biologi dan Aplikasinya*, 4(1):19-25. DOI: 10.26740/jrba.v4n1.p19-25.

INTRODUCTION

The pandemic of COVID-19 caused by positive-sense RNA of β -coronaviridae known as Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2). The first case started to spread in Wuhan, China, in December 2019 (Coronaviridae Study Group of the International Committee on Taxonomy of Viruses, 2020; Holshue *et al.* 2020).

There were 2-3% infection and death cases caused by SARS-CoV-2 and continue to increase without any significant decreasing number (Hamid *et al.* 2020). World Health Organization (WHO) reported that for now on, effective treatment using antiviral agents against SARS-CoV-2 is still not available (Pandey *et al.* 2020). Several symptoms of COVID-19 disease are sputum production (28%), headache

(8%), hemoptysis (5%), and diarrhea (3%) (Huang *et al.* 2020). According to several studies, other symptoms such as anosmia, fever, cough, alveolar destruction, and acute pneumonia was detected after ten days of infection with more severe symptoms on comorbid patients and old patients >70 years old (Chen *et al.* 2020; Li *et al.* 2020; Tian *et al.* 2020; Wang W *et al.* 2020; Zhu *et al.* 2020).

Indonesia is known as a country with high biodiversity that has several plants used as traditional medicine, one of them is orchids (Darmawati *et al.* 2018; Fandani *et al.* 2018). Orchids include in the Orchidaceae family with high distribution over the world and very abundance in species richness. There are almost 5000 of 30.000 species of orchids can be found in the Indonesia archipelago (Sapto, 2009). The genus of *Dendrobium* become the second largest genus in the Orchidaceae family (Moudi *et al.*, 2013). *Dendrobium* has been used as natural traditional medicine in China and the studies show that some of the species contain secondary metabolites group with antiviral activity (Fan *et al.* 2001). Particular of secondary metabolites group as flavonoids, alkaloids, derivatives of bibenzyl have potential in the medication field (Gutierrez, 2010).

Herbal medication is a treatment using complex mixture of plant organic compounds (Ren *et al.* 2020). A secondary metabolite is an intermediate substance which do not have a role in the main metabolic processes, specifically and has a certain function in each plant family (Semiarti *et al.*, 2020). Plant secondary metabolite useful for inhibitor of specific disease with lower side effect also beneficial as alternative of an antiviral candidate with high bioavailability and low cytotoxic effect (Umadevi *et al.* 2020). Catechins, zingerols, gingerols, alicins, quercetins, and epitacetin reported have inhibition activity towards SARS-CoV-2 by *in silico* study (Khaerunnisa *et al.* 2020). The quest of antiviral candidates from the natural substance as anti-SARS-CoV-2 facing challenges due to differences in genome sequence with SARS-CoV-1 or MERS-CoV (Septiana, 2020).

The antiviral candidate analysis in botany sectors should be started with *in silico* study (Trivedi *et al.* 2020). *In silico* study expected to optimize research process and prevent loss in *in vitro* and *in vivo* steps, appropriately, accurately, rapidly, and effectively (Sumon *et al.* 2021). Mostly researches that discusses the effects of the secondary metabolite is still in progress of *in vitro* testing and the *in vivo* testing effectiveness is not

yet guaranteed (Septiana, 2020). Experimental research to find drug candidates from natural substance is difficult in the pandemic condition. *In silico* method approach based on molecular docking need to be done due to it is relatively fast and safe. The aim of this study was to determine the potential of secondary metabolite on *Dendrobium* orchid species as an antiviral candidate on main protease (M^{pro}) SARS-CoV-2 using molecular docking by *in silico* method.

MATERIALS AND METHODS

Molecular Docking

Secondary metabolites of *Dendrobium* listed from KNApSack: Metabolite Ecology. 3-dimension structure of secondary metabolites downloaded from PubChem databases with .sdf format. Molecular docking conducted at online platform small molecule Covid-19 Docking Server parameter <https://ncov.schanglab.org.cn/index.php>. (Kong, 2020). Structure of secondary metabolites act as ligand in docking. Sars-CoV2 main Protease (M^{pro}) with PDB ID: 6LU7 selected as target (Jin *et al.* 2020). Centre Coordinate of Grid Box set in x = -10.85 Å; y = 12.58 Å; z = 68.72 Å and the size of grid box 30 Å (r=12) (Vidomini *et al.* 2021). Rate of exhaustiveness in 12. The higher rate of exhaustiveness would get a more constant result (Forli, 2016). The result downloaded in .zip format

Ligand Profiling

Binding affinity score between target and ligand which lower than inhibitor N3 (Table 1) as native ligand would analyzed furthermore by ligand profiling. Manifesting ligand profile by Lipinski's rule of five that is hydrogen bond donors (the total number of nitrogens-hydrogen and oxygen-hydrogen bonds), hydrogen bond acceptors (all nitrogen or oxygen atoms), molecular mass less, and lipophilicity (log P) conducted by <http://www.swissadme.ch/index.php>

Data Visualization

Ligand and target 3-dimension structure conformation visualized in Software Discovery Studio 2020. The three strongest ligand and target interaction analyzed the amino acid residue contact and interaction in binding pocket.

RESULTS AND DISCUSSION

Dendrobium is generally found growing well in high areas or mountainous areas 1400-1600 m above sea level, in a fairly humid and foggy environment, and with a mild average temperature (Lam *et al.*, 2015). In this study, secondary metabolites of the genus *Dendrobium* acted as

ligand-independent variables. Secondary metabolites composed of *Dendrobium* polysaccharide compounds have antiviral activity (Hwang et al., 2012). Binding of secondary metabolites to viral nucleoprotein regions can inhibit oligomerization and export of viral nucleoproteins (Ti et al., 2021). Medioresinol is capable of causing the accumulation of intracellular ROS and apoptosis of microbial cells (Hwang et al., 2012). Terpenoids in *Dendrobium nobile* play a role in the proliferative activity of murine T and B lymphocytes as antiviral immunomodulators (Singh et al., 2012). Naringenin is believed to have potential as a treatment compound that can inhibit SARS-CoV in Mpro and ACE2 (Tutunchi et al., 2020).

The main protease (M^{pro}) is known to play an important role in the live replication process of SARS-CoV-2 and is one of the causes of rapid transmission (Purwaniati and Asnawi, 2020). The M^{pro} structure of SARS-CoV-2 consists of A and B Chains (Zhang et al. 2020). M^{pro}, also called the 3CL protease, is a 33.8-kDa cysteine protease that

mediates functional polypeptide maturation in the assembly of viral replication machinery (Cui et al. 2020). The role of M^{pro} in the life cycle of SARS-CoV-2 uses one of the important targets in the design of COVID-19 antivirals (Purwaniati and Asnawi, 2020). In this study, M^{pro} SARS-CoV-2 (6LU7) was used as a macromolecular target.

N₃ inhibitors or known as Michael Acceptor Inhibitors have been investigated to be able to inhibit the virus and specifically inhibit M^{pro} SARS-CoV (Yang et al. 2005). In this study, N₃ inhibitors were used as native ligands obtained from the RCSB website (Jain and Mujwar, 2020). The N₃ inhibitor covalently binds to SARS-CoV-2 M^{pro} via the Michael reaction and blocks its active site (Griffin, 2020; Xue et al. 2008). N₃ cytotoxicity 50% concentration of 133µM, ebselen and N₃ antiviral activity had the strongest effect at 10 M concentration for Vero cells infected by SARS-CoV-2, N inhibitor compounds were able to penetrate cell membranes to access more targeted targets (Jin et al., 2020).

Table 1. Ligand Profiling secondary metabolite of *Dendrobium*

Secondary metabolites	Binding affinity Bioavailability		Lipinski's rule				
	Kcal/Mol	Score	Log ^p	Violation	H-Bond		MR (g/mol)
					Donor	Acceptor	
Medioresinol	-8	0.55	3.29	0	2	7	388.41
Denthyrsinol	-8.1	0.55	3.98	0	4	6	478.49
Dendroside D	-8.2	0.17	2.21	3	9	14	592.63
Nobilin E	-8	0.55	3.38	1	3	8	544.59
(+)-Lirioresinol B	-7.9	0.55	3.52	0	2	8	418.44
Denthyrsinone	-7.9	0.56	2.41	1	3	8	522.50
Naringenin	-7.8	0.55	1.75	0	3	5	272.25
Dendrocandin B	-7.7	0.55	4.24	0	2	8	482.52
4,4'-Dihydroxy-3,5 dimethoxybibenzyl	-7.6	0.55	2.67	0	2	4	274.31
Acanthoside B	-7.6	0.17	3.5	2	5	13	580.58
Inhibitor N3	-7.5						

Table 2. Binding affinity and Binding Pocket secondary metabolites of *Dendrobium* with M^{pro}

Species	Secondary metabolites	Binding affinity (kcal/mol)	Bio-availability Score	Binding Pocket
<i>Dendrobium thysiflorum</i>	Denthyrsinol	-8.1	0.55	ASN142, GLY143, THR26, CYS145, LEU27, GLU166, SER144,
<i>Dendrobium thysiflorum</i>	Denthyrsinone	-7.9	0.56	CYS145, HIS163, HIS41, GLU166, PHE140, GLY143,
<i>Dendrobium candidum</i>	Dendrocandin B	-7.7	0.55	CYS145, HIS172, MET49, THR25, GLU166, HIS41, MET165, PHE140, THR190,
	Inhibitor N3	-7.5	0.17	ASN142, GLU166, LEU27, CYS145, HIS163, MET49, GLN189, HIS41,




Figure 1. Visualization of strongest conformation secondary metabolites of *Dendrobium* spp., with M^{pro}
A. Dendrocandin, B. Denthyrsinone, C. Denthyrsinol, D. Inhibitor N3

In this research, an *insilico* study was conducted with the mechanism of the stages (1) ligand preparation, (2) target preparation, (3) target ligand molecule docking, (4) ligand-protein interaction visualization, and (5) data analysis (Sari et al. 2020). Molecular docking is a popular method in *in silico*

studies, especially to initiate drug design (de Ruyck et al. 2016). The docking process is divided into two types, namely blind docking and oriented docking (Pradani et al., 2021). Blind docking is a docking process without knowing the exact location of the active site of the receptor, while oriented docking is done after knowing the exact location of the active site of the receptor (Syahputra et al., 2014). Molecular docking can be done with the PyRx software using the Autodock tool.

In this study, molecular docking was carried out with the Online Covid-19 Docking Server using the AutoDock Vina device (Vicidomini et al. 2021). Autodock Vina and CoDock PP were used as docking engines to predict the binding mode between COVID-19 targets and potential ligands. Covid-19 Docking Server is economically and operationally easy to use (Kong et al. 2020). One of the studies using the Online Covid-19 Docking Server is Pendyala and Patras (2020), which examines several bioactive compounds in food ingredients to inhibit M^{pro} COVID-19. Research conducted by Barik et al. (2020) also used the Online Covid-19 Docking Server as a method to analyze the types of bonds in chloroquine, hydroxychloroquine, ramdesivir and arbidol compounds. The web-based docking platform has been validated by its developers with the evaluation of re-docking trials for targets, including M^{pro} with N₃ inhibitors (Vicidomini et al. 2021).

The most negative binding affinity value for each ligand compared to the native ligand binding affinity (Chen et al. 2016; Xue et al. 2008). Based on the results in Table 1., there are 10 secondary metabolites of *Dendrobium* with a lower binding

affinity value than the native ligand Inhibitor N3. The metabolites dendroside D, denthyrsinol, medioresinol, and nobilin E respectively had the lowest binding affinity, namely -8.2 kcal/mol, -8.1 kcal/mol, -8 kcal/mol, and -8 kcal/mol. Binding affinity indicates the strength of the drug to bind to the receptor (Arwansyah et al. 2014). The lower the binding affinity value, the higher the affinity between the receptor and the ligand and vice versa (Prabowo and Santoso, 2018). The smaller the binding affinity value, the more stable the interaction between the ligand and the receptor (Arwansyah et al. 2014).

The ligand profile in Table 1 shows that of the 10 secondary metabolites of *Dendrobium*, only Dendroside D and Acanthoside B are not potential drugs because they have Violation >2 (Wells and McGee, 2008). The profile of ligands according to Lipinski's five rules includes molecular weight, logP (lipophilicity), number of hydrogen bond donors, number of hydrogen bond acceptors, and Violation (Rachmania et al. 2015). Lipinski's rule describes the solubility of certain compounds to penetrate cell membranes by passive diffusion (Syahputra et al. 2014). Compounds are more permeable and active as ligands if they have <5 hydrogen bond donors, <10 hydrogen bond acceptors, molecular mass <500 Da, and logP value <5 (Afriza et al. 2018). The value of Mr (molecular weight), H-Donor and H-acceptor is a measure of the permeability of a drug to be able to pass through the lipid bilayer of a cell (Suhadi et al. 2019). Lipophilicity is defined as the logarithm of the ratio of the hydrophobicity of the drug partitioning into the organic phase to the aqueous phase and is referred to as logP (Ivanovic et al. 2019; Utomo et al. 2017). Violation is related to the potential for dependence and the amounts of illegal drugs used (Wells and McGee, 2008). Hydrogen bonds generally act as a facilitator to increase the binding affinity of ligands by moving

protein-bound water molecules into a large volume of solvent (Chen et al., 2016).

Visualization of ligand-protein interactions of 10 secondary metabolites showed that there were only 3 strongest metabolite modes. Secondary metabolites with the strongest mode against M^{pro} were found in *Dendrobium thysiflorum* and *Dendrobium candidum* orchids. The strongest metabolite mode in terms of the lowest binding affinity and interactions with the most hydrogen bonds, especially on key amino acids. Based on the results of Table 2, the metabolites that have the highest probability of binding to M^{pro} are dendrocandin B, denthysrinone, and denthysrinol. Denthysrinol has the lowest binding affinity of -8.1 kcal/mol which is predicted to be more stable when bound to M^{pro}. The interactions formed by Denthysrinol and M^{pro} are hydrogen bonds, electrostatic interactions, and hydrophobicity with the amino acids GLY143, LEU27, SER144, THR26 and ASN142, CYS145, GLU166 as key amino acids. The interaction between the ligand and the protein receptor is expected to inhibit the performance of M^{pro} SARS-CoV-2 (Trott and Olson, 2010).

CONCLUSION

Based on molecular docking using *in silico* method, it can be concluded that Denthysrinol is a the most potential secondary metabolite in orchid *Dendrobium thysiflorum* as an antiviral candidate on main protease (M^{pro}) SARS-CoV-2.

REFERENCES

- Afriza, D., Suriyah, W. H., & Ichwan, S. J. A. (2018). In silico analysis of molecular interactions between the anti-apoptotic protein survivin and dentatin, nordentatin, and quercetin. *Journal of Physics: Conference Series*, 1073(3), 0–7. <https://doi.org/10.1088/1742-6596/1073/3/032001>.
- Arwansyah, A., Ambarsari, L. and Sumaryada, T.I. (2014). Simulasi docking senyawa kurkumin dan analognya sebagai inhibitor reseptor androgen pada kanker prostat [English: “Docking simulation of curcumin and its analogues as androgen receptor inhibitors in prostate cancer”]. *Current Biochemistry*, 1(1): 11–19.
- Barik, A., Rai, G., & Modi, G. (2020). *Molecular docking and binding mode analysis of selected FDA approved drugs against COVID-19 selected key protein targets: An effort towards drug repurposing to identify the combination therapy to combat COVID-19*. <http://arxiv.org/abs/2004.06447>.
- Chen, D., Oezguen, N., Urvil, P., Ferguson, C., Dann, S. M., & Savidge, T. C. (2016). Regulation of protein-ligand binding affinity by hydrogen bond pairing. *Science Advances*, 2(3). <https://doi.org/10.1126/sciadv.1501240>.
- Chen, N., Zhou, M., Dong, X., Qu, J., Gong, F., Han, Y., Qiu, Y., Wang, J., Liu, Y., Wei, Y., Xia, J., Yu, T., Zhang, X., & Zhang, L. (2020). Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *The Lancet*, 395(10223), 507–513. [https://doi.org/10.1016/S0140-6736\(20\)30211-7](https://doi.org/10.1016/S0140-6736(20)30211-7).
- Coronaviridae Study Group of the International Committee on Taxonomy of Viruses. (2020). The species severe acute respiratory syndrome-related coronavirus: classifying 2019-nCoV and naming it SARS-CoV-2. *Nature microbiology*, 5(4): 536–544.
- Cui, W., Y. Kailin, and Y. Haitao. (2020). Recent Progress in the Drug Development Targeting SARS-CoV-2 Main Protease as Treatment for COVID-19. *Frontiers in Molecular Biosciences*, 7(398): 1–10. doi: 10.3389/fmolb.2020.616341
- Darmawati, I. A. P., Rai, I. N., Dwiyani, R., & Astarini, I. A. (2018). Short communication: The diversity of wild Dendrobium (Orchidaceae) in Central Bali, Indonesia. *Biodiversitas*, 19(3), 1110–1116. <https://doi.org/10.13057/biodiv/d190345>.
- de Ruyck, J., Brysbaert, G., Blosssey, R., & Lensink, M. F. (2016). Molecular docking as a popular tool in drug design, an in silico travel. *Advances and Applications in Bioinformatics and Chemistry*, 9(1), 1–11. <https://doi.org/10.2147/AABC.S105289>.
- Fan, C., Wang, W., Wang, Y., Qin, G., & Zhao, W. (2001). Chemical constituents from *Dendrobium densiflorum*. *Phytochemistry*, 57(8), 1255–1258. [https://doi.org/10.1016/S0031-9422\(01\)00168-6](https://doi.org/10.1016/S0031-9422(01)00168-6).
- Fandani, H. S., Mallomasang, S. N., & Korja, I. N. (2018). Keanekaragaman Jenis Anggrek pada beberapa Penangkaran di Desa Ampera dan Desa Karunia Kecamatan Palolo Kabupaten Sigi. *Jurnal Warta Rimba*, 6(9), 14–20.
- Forli, S., Huey, R., Pique, M. E., Sanner, M., Goodsell, D. S., & Arthur, J. (2016). 00006565-201002000-00017. 11(5), 905–919. <https://doi.org/10.1038/nprot.2016.051>. Computation al
- Griffin, J. W. D. (2020). SARS-CoV and SARS-CoV-2 main protease residue interaction networks change when bound to inhibitor N3. *Journal of Structural Biology*, 211(3), 107575. <https://doi.org/10.1016/j.jsb.2020.107575>.
- Gutiérrez, R. M. P. (2010). Orchids: A review of uses in traditional medicine, its phytochemistry and pharmacology. *Journal of Medicinal Plants Research*, 4(8), 592–638. <https://doi.org/10.5897/JMPR10.012>.
- Hamid, S., Mir, M. Y., & Rohela, G. K. (2020). Novel coronavirus disease (COVID-19): a pandemic (epidemiology, pathogenesis and potential therapeutics). *New Microbes and New Infections*, 35, 100679. <https://doi.org/10.1016/j.nmni.2020.100679>.
- Holshue, M. L., DeBolt, C., Lindquist, S., Lofy, K. H., Wiesman, J., Bruce, H., Spitters, C., Ericson, K., Wilkerson, S., Tural, A., Diaz, G., Cohn, A., Fox, L., Patel, A., Gerber, S. I., Kim, L., Tong, S., Lu, X., Lindstrom, S., Pillai, S. K. (2020). First Case of 2019 Novel Coronavirus in the United States. *New England Journal of Medicine*, 382(10), 929–936. <https://doi.org/10.1056/nejmoa2001191>.
- Huang, M., Tang, T., Pang, P., Li, M., Ma, R., Lu, J., Shu, J., You, Y., Chen, B., Liang, J., Hong, Z., Chen, H., Kong, L., Qin, D., Pei, D., Xia, J., Jiang, S., & Shan, H. (2020). Treating COVID-19 with Chloroquine. *Journal of Molecular Cell Biology*, 12(4), 322–325. <https://doi.org/10.1093/jmcb/mjaa014>.
- Hwang, J. H., Hwang, I. S., Liu, Q. H., Woo, E. R., & Lee, D. G. (2012). Medioresinol leads to intracellular ROS accumulation and mitochondria-mediated apoptotic cell death in *Candida albicans*. *Biochimie*, 94(8), 1784–

1793. <https://doi.org/10.1016/j.biochi.2012.04.010>.
- Ivanović, V., Rančić, M., Arsić, B., & Pavlović, A. (2020). Lipinski's rule of five, famous extensions and famous exceptions. *Popular Scientific Article*, 3(1), 171-177.
- Jain, R., & Mujwar, S. (2020). Repurposing metocurine as main protease inhibitor to develop novel antiviral therapy for COVID-19. *Structural Chemistry*, 31(6), 2487-2499. <https://doi.org/10.1007/s11224-020-01605-w>.
- Jin, Z., Du, X., Xu, Y., Deng, Y., Liu, M., Zhao, Y., Zhang, B., Li, X., Zhang, L., Peng, C., Duan, Y., Yu, J., Wang, L., Yang, K., Liu, F., Jiang, R., Yang, X., You, T., Liu, X., Yang, H. (2020). Structure of Mpro from SARS-CoV-2 and discovery of its inhibitors. *Nature*, 582(7811), 289-293. <https://doi.org/10.1038/s41586-020-2223-y>.
- Khaerunnisa, S., Kurniawan, H., Awaluddin, R., Suhartati, S., & Soetjipto, S. (2020). Potential Inhibitor of COVID-19 Main Protease (Mpro) From Several Medicinal Plant Compounds by Molecular Docking Study Molecular Docking, ADME-Toxicity Prediction, and Evaluation of Curcumin Derivative Compound as Inhibitor Inflammation on Rheumatoid Arth. *Preprints, March*. <https://doi.org/10.20944/preprints202003.0226.v1>.
- Kong, R., Yang, G., Xue, R., Liu, M., Wang, F., Hu, J., Guo, X., & Chang, S. (2020). COVID-19 Docking Server: A meta server for docking small molecules, peptides and antibodies against potential targets of COVID-19. *Bioinformatics*, 36(20), 5109-5111. <https://doi.org/10.1093/bioinformatics/btaa645>.
- Lam, Y., Ng, T. B., Yao, R. M., Shi, J., Xu, K., Sze, S. C. W., & Zhang, K. Y. (2015). Evaluation of chemical constituents and important mechanism of pharmacological biology in *Dendrobium* plants. *Evidence-Based Complementary and Alternative Medicine*, 2015. <https://doi.org/10.1155/2015/841752>.
- Li, X., Geng, M., Peng, Y., Meng, L., & Lu, S. (2020). Molecular immune pathogenesis and diagnosis of COVID-19. *Journal of Pharmaceutical Analysis*, 10(2), 102-108. <https://doi.org/10.1016/j.jpaha.2020.03.001>
- Moudi, M., Go, R., Yong, C., Yien, S., & Nazre, M. (2013). *A Review on Molecular Systematic of the Genus Dendrobium Sw.* 2 (April), 71-78.
- Pandey, P., Rane, J. S., Chatterjee, A., Kumar, A., & Ray, S. (2020). Targeting SARS-CoV-2 Spike Protein of COVID-19 with Naturally Occurring Phytochemicals: An in Silico Study for Drug Development. *ChemRxiv*. <https://doi.org/10.26434/chemrxiv.12094203.v1>
- Pendyala, B., & Patrasa, A. (2020). In silico Screening of Food Bioactive Compounds to Predict Potential Inhibitors of COVID-19 Main protease (M. *ChemRxiv*, 11-44).
- Prabowo, S.A.A.E. and Santoso, B., 2018. Profil in silico Interaksi Senyawa Alam ketumbar dan Adas Bintang sebagai Inhibitor Peptida Deformilase *Mycobacterium tuberculosis* (3SVJ dan 1WS1) menggunakan Bantuan PyRx-Vina [English: "In silico Profile of Interaction of Coriander and Star Anise Natural Compounds as Peptide Deformylase Inhibitors of *Mycobacterium tuberculosis* (3SVJ and 1WS1) using PyRx-Vina Assistance"]. *Proceeding of the URECOL*, pp.402-408.
- Pradani, T.C., Manampiring, A.E., Kepel, B.J., Budiarmo, F.D. and Bodhi, W., 2021. Molecular Docking Terhadap Senyawa Kurkumin dan Arturmeron pada Tumbuhan Kunyit (*Curcuma Longa* Linn.) yang Berpotensi Menghambat Virus Corona [English: "Molecular Docking of Curcumin and Arturmeron Compounds in Turmeric Plants (*Curcuma Longa* Linn.) Potentially Inhibiting Corona Virus"]. *eBiomedik* 9(2): 208-214.
- Purwaniati, and A. Asnawi. (2020). Target Kerja Obat Antivirus Covid-19: Review Drug Target of Antivirus Covid-19: Review. *Jurnal Farmagazine*, 7(2): 30-42
- Rachmania, R. A., Supardi, and O. A. Larasati. (2015). In Silico Analysis of Diterpenoid Lactone Compounds of Bitter Herbs (*Andrographis paniculata* Nees) on Alpha-Glucosidase Receptor as Antidiabetic Type II Agents. *PHARMACY*, 12(2): 210-222.
- Rane, J. S., Chatterjee, A., Kumar, A., & Ray, S. (2020). Targeting SARS-CoV-2 Spike Protein of COVID-19 with Naturally Occurring Phytochemicals: An in Silico Study for Drug Development. *ChemRxiv*, <https://doi.org/10.26434/chemrxiv.12094203.v1>.
- Sari, I.W., Junaidin, J. and Pratiwi, D., 2020. Studi Molecular Docking Senyawa Flavonoid Herba Kumis Kucing (*Orthosiphon stamineus* B.) Pada Reseptor A-Glukosidase Sebagai Antidiabetes Tipe 2 [English: "Molecular Docking Study of Flavonoid Compounds of Cat's Whisker Herb (*Orthosiphon stamineus* B.) on -Glucosidase Receptors as Type 2 Antidiabetic"]. *Jurnal Farmagazine*, 7(2), pp.54-60.
- Semiarti, E., Purwantoro, A., & Puspita Sari, I. (2020). *Biotechnology Approaches on Characterization, Mass Propagation, and Breeding of Indonesian Orchids *Dendrobium lineale* (Rolle) and *Vanda tricolor* (Lindl.) with Its Phytochemistry*, 1-14. https://doi.org/10.1007/978-3-030-11257-8_12-1.
- Septiana, E., 2020. Prospek Senyawa Bahan Alam Sebagai Antivirus Dalam Menghambat SARS-CoV-2 [English: "Prospects of Natural Compounds As Antivirals In Inhibiting SARS-CoV-2"]. *Biotrends* 11(1): 30-38.
- Singh, Siddhartha Singh, Amit Kumar, Sunil Kumar, Mukul Pandey, Pramod Singh, & Mayanglambam. (2012). Medicinal Properties and Uses of Orchids: a Concise Review. *Elixir Appl. Botany*, 2012 (November), 11627-11634. https://www.researchgate.net/publication/292131192_Medicinal_properties_and_uses_of_orchids_a_concise_review
- Suhadi, A., Rizarullah, R., & Feriyani, F. (2019). Simulasi Docking Senyawa Aktif Daun Binahong Sebagai Inhibitor Enzyme Aldose Reductase. *Sel Jurnal Penelitian Kesehatan*, 6(2), 55-65. <https://doi.org/10.22435/sel.v6i2.1651>
- Sumon, T. A., Hussain, M. A., Hasan, M. T., Hasan, M., Jang, W. J., Bhuiya, E. H., Chowdhury, A. A. M., Sharifuzzaman, S. M., Brown, C. L., Kwon, H. J., & Lee, E. W. (2021). A Revisit to the Research Updates of Drugs, Vaccines, and Bioinformatics Approaches in Combating COVID-19 Pandemic. *Frontiers in Molecular Biosciences*, 7. <https://doi.org/10.3389/fmolb.2020.585899>.
- Suryadi Budi Utomo, Fajar Sanubari, Budi Utami, dan N. D. N., & Program. (2017). Aktivitas Analgesik Senyawa Turunan Meperidin Menggunakan Metode Semiempiris Am1 Analysis of a Quantitative Relationship Between the Structure and Analgesic Activity of Meperidin Derivatives Using. (*Jurnal Kimia Dan Pendidikan Kimia*, 2(3), 158-168.
- Syahputra, G., L. Ambarsari, and T. Sumaryada. (2014). Simulasi Docking Kurkumin Enol, Bisdemetoksikurkumin dan Analognya sebagai Inhibitor Enzim 12-Lipoksigenase [English: "Docking Simulation of Enol Curcumin,

- Bisdemethoxycurcumin and Their Analogues as Enzyme 12-Lipoxygenase Inhibitors”]. *Jurnal Biofisika*, 10 (1): 55-67
- Ti, H., Zhuang, Z., Yu, Q., & Wang, S. (2021). Progress of plant medicine derived extracts and alkaloids on modulating viral infections and inflammation. *Drug Design, Development and Therapy*, 15, 1385–1408. <https://doi.org/10.2147/DDDT.S299120>
- Tian, S., Hu, N., Lou, J., Chen, K., Kang, X., Xiang, Z., Chen, H., Wang, D., Liu, N., Liu, D., Chen, G., Zhang, Y., Li, D., Li, J., Lian, H., Niu, S., Zhang, L., & Zhang, J. (2020). Characteristics of COVID-19 infection in Beijing. *Journal of Infection*, 80 (4), 401–406. <https://doi.org/10.1016/j.jinf.2020.02.018>
- Trivedi, G. N., Karlekar, J. T., Dhameliya, H. A., & Panchal, H. (2020). A review on the novel coronavirus disease based on in-silico analysis of various drugs and target proteins. *Journal of Pure and Applied Microbiology*, 14 (1), 849–860. <https://doi.org/10.22207/JPAM.14.SPL1.22>
- Trott, O., & Olson, A. J. (2009). AutoDock Vina: Improving the speed and accuracy of docking with a new scoring function, efficient optimization, and multithreading. *Journal of Computational Chemistry*, 31(2), NA-NA. <https://doi.org/10.1002/jcc.21334>
- Tutunchi, H., Naeini, F., Ostadrahimi, A., & Hosseinzadeh-Attar, M. J. (2020). Naringenin, a flavanone with antiviral and anti-inflammatory effects: A promising treatment strategy against COVID-19. *Phytotherapy Research*, 34 (12), 3137–3147. <https://doi.org/10.1002/ptr.6781>
- Umadevi, P., Manivannan, S., Fayad, A. M., & Shelvy, S. (2020)p. In silico analysis of phytochemicals as potential inhibitors of proteases involved in SARS-CoV-2 infection. *Journal of Biomolecular Structure and Dynamics*, 0(0), 1–9. <https://doi.org/10.1080/07391102.2020.1866669>
- Utomo, S. B., F. Sanubari, B. Utami, and N. D. Nurhayati. (2017). Analisis Hubungan Kuantitatif Struktur dan Aktivitas Analgesik Senyawa Turunan Meperidin Menggunakan Metode Semiempiris AM1 [English: “Analysis of the Quantitative Relationship of the Structure and Analgesic Activity of Meperidine Derivative Compounds Using the AM1. Semiempirical Method”]. *Jurnal Kimia dan Pendidikan Kimia*, 2(3): 158-168
- Vicidomini, C., Roviello, V., & Roviello, G. N. (2021). In silico investigation on the interaction of chiral phytochemicals from opuntia ficus-indica with sars-cov-2 mpro. *Symmetry*, 13(6). <https://doi.org/10.3390/sym13061041>.
- Wang, C., Horby, P.W., Hayden, F.G. and Gao, G.F. (2020). A novel coronavirus outbreak of global health concern. *The lancet*, 395 (10223): 470-473.
- Wang, W., Tang, J., & Wei, F. 2020. Updated understanding of the outbreak of 2019 novel coronavirus (2019-nCoV) in Wuhan, China. *Journal of Medical Virology*, 92 (4), 441–447. <https://doi.org/10.1002/jmv.25689>.
- Wang, Z., Zhao, M., Cui, H., Li, J., & Wang, M. (2020). Transcriptomic Landscape of Medicinal Dendrobium Reveals Genes Associated With the Biosynthesis of Bioactive Components. *Frontiers in Plant Science*, 11 (April), 1–11. <https://doi.org/10.3389/fpls.2020.00391>
- Wells, J. E., & McGee, M. A. (2008). Violations of the usual sequence of drug initiation: Prevalence and associations with the development of dependence in the New Zealand Mental Health Survey. *Journal of Studies on Alcohol and Drugs*, 69 (6), 789–795. <https://doi.org/10.15288/jsad.2008.69.789>.
- Xue, X., Yu, H., Yang, H., Xue, F., Wu, Z., Shen, W., Li, J., Zhou, Z., Ding, Y., Zhao, Q., Zhang, X. C., Liao, M., Bartlam, M., & Rao, Z. (2008). Structures of Two Coronavirus Main Proteases: Implications for Substrate Binding and Antiviral Drug Design. *Journal of Virology*, 82 (5), 2515–2527. <https://doi.org/10.1128/jvi.02114-07>
- Zhu, N., Zhang, D., Wang, W., Li, X., Yang, B., Song, J., Zhao, X., Huang, B., Shi, W., Lu, R., Niu, P., Zhan, F., Ma, X., Wang, D., Xu, W., Wu, G., Gao, G. F., & Tan, W. (2020). A Novel Coronavirus from Patients with Pneumonia in China, 2019. *New England Journal of Medicine*, 382(8), 727–733. <https://doi.org/10.1056/nejmoa2001017>.