

## WORLD CONFEDERATION FOR PHYSICAL THERAPY

# *Ethical Principles for Physical Therapists*

**T**he need for physical therapy is universal. The service of the physical therapist is, therefore, unrestricted by considerations of nationality, race, creed, colour, politics or social status.

The activities and relationships of physical therapists must vary widely in the many countries and spheres in which they are pursued and each national organisation must be free to make its own detailed code of ethics to which it may require its members to conform.

**Heat and Exercise—Continued from Page 7.**

### PART II.

The second method of treatment favoured for gynaecological conditions is that of Exercise—here again divided into breathing exercises, assisted, free and resisted exercises.

#### 1. Breathing Exercises.

Breathing exercises are given as a prophylactic measure against chest complaints following abdominal operations. In the Bantu respiratory troubles, particularly after Caesarean Section, seem to arise very quickly, and pre- and post-operative breathing exercises, together with leg exercises for the prevention of thrombosis, are of definite value.

If, in spite of these exercises, atelectasis occurs the use of a vibrator and back clapping will loosen secretions and promote healthier action of the lungs.

#### 2. Assisted exercises.

In the treatment of post-natal dropped foot, which is known by various names, from "obstetrical palsy" to "traumatic neuritis of the puerperium," assisted exercises are used, following electrical stimulation by the current most suited to the condition. (In a very slight case, surging faradism will be tolerated, but in a bad case one of the progressive currents will be required.)

The Bantu patient on the whole has a smaller pelvis, from the obstetric capacity point of view, than the European. Consequently the pressure effect of the engaging head is more severe on the large lumbo-sacral nerve trunks which cross the alae of the sacrum. Damage to these nerve trunks, which form part of the great sciatic nerve, account for the comparatively large number of dropped foot cases seen in Non-European hospital practise.

#### 3. Free exercises.

These include the well known ante- and post-natal exercises about which so much has already been written.

Dysmenorrhoea is greatly improved by free exercises. In similar cases among European women ballet lessons have proved helpful. Women suffering from this disability enjoy "free exercises," served in this fashion.

#### 4. Resisted exercises.

The use of Kegels perineometer can be described as a "resisted exercise." Inserted vaginally it is used at Edenvale with marked success in causes of cystocele, rectocele and prolapse; and, in conjunction with faradism, in cases of stress incontinence, a lax pelvic floor, or for general toning of the levator ani in elderly hysterectomy cases.

Inserted rectally it can be used to improve ano-rectal tone and continence after cancer operations, where the removal of the bladder, uterus and vagina has become necessary.

In conclusion it can be truthfully said that with accurate diagnosis and well controlled therapy, great improvement in all types of gynaecological cases is possible by the use of heat and exercises.

The following general principles, to which all Member-Organisations of the World Confederation for Physical Therapy have declared their adherence, will serve as a guide to national organisations in building up their individual Code of Ethics:

1. Physical Therapists recognise not only the responsibilities but also the limitations of their professional functions. They therefore only treat those patients referred to them by a registered medical practitioner.
2. Physical Therapists carry out with loyalty and the utmost skill any physical treatment prescribed by a registered medical practitioner which is aimed at restoring, improving or maintaining the mental and physical fitness of the patient.
3. Physical Therapists maintain at all times the highest standards of knowledge and skill.
4. Physical Therapists respect the culture and religious beliefs of their patients and associates.
5. Physical Therapists co-operate loyally with their physical therapy colleagues and with members of allied health professions and they sustain the confidence of the patient in all members of the health team.
6. Physical Therapists hold in confidence all personal information entrusted to them and will not discuss a patient's affairs with others than those responsible for the patient's care.
7. Physical Therapists recognise that failure of a colleague to conform to the ethical principles of the profession is detrimental to the patient and therefore such conduct should be reported to the proper authority. Unethical conduct should, however, never be reported in the presence of patients or associates.
8. Physical Therapists share responsibility with other citizens and members of other professions in meeting the needs of the public in matters of health, and should participate in all governmental and voluntary efforts to meet those needs.
9. Physical Therapists do not permit their names to be used in connection with the advertisement of products or with any other form of self-advertisement except as expressly authorised by the Code of Ethics of their national association.
10. Physical Therapists believe that just remuneration for their services should be received, but they may accept only such compensation or professional fee as the contract, actual or implied, provides.
11. Physical Therapists adhere at all times to standards of professional and personal ethics which reflect credit upon the profession.