

Efficacy and safety of tralokinumab treatment in adults of different racial subgroups with moderate-to-severe atopic dermatitis in three randomized, placebo-controlled phase 3 trials

Tiffany Mayo¹, April Armstrong², Leon Kircik³, Jonathan I. Silverberg⁴, Andrew Blauvelt⁵, Ben Esdaile⁶, Shannon Schneider⁷, Thomas Mark⁸, Melinda Gooderham⁹, Andrew F. Alexis¹⁰

¹University of Alabama at Birmingham, Birmingham, AL, USA; ²Keck School of Medicine of University of Southern California, Los Angeles, CA, USA; ³Icahn School of Medicine at Mount Sinai, New York, NY, USA; ⁴The George Washington School of Medicine and Health Sciences, Washington, DC, USA; ⁵Oregon Medical Research Center, Portland, OR, USA; ⁶Whittington Health National Health Service Foundation Trust, London, UK; ⁷LEO Pharma Inc., Madison, NJ, USA; ⁸LEO Pharma A/S, Ballerup, Denmark; ⁹SKIN Centre for Dermatology, Peterborough, ON, Canada; ¹⁰Weill Cornell Medicine, New York, NY, USA

Introduction

- Atopic dermatitis (AD) is a chronic inflammatory skin disease associated with significant disease burden
- Although AD is highly prevalent in patients with skin of color, data on the efficacy and safety of AD therapies in these patients is limited since most clinical trials enroll predominantly White patients¹
 - Several standard measures, including EASI, can underestimate AD severity in dark skin¹
- Tralokinumab, a specific, high-affinity interleukin-13 inhibitor, is approved in Europe, Canada, and the United States for the treatment of adults with moderate-to-severe AD
- ECZTRA 1 (NCT03131648), ECZTRA 2 (NCT03160885), and ECZTRA 3 (NCT03363854) were randomized phase 3 trials assessing the safety and efficacy of tralokinumab or tralokinumab + TCS, as needed.
 - ECZTRA 1 and 2 were placebo-controlled trials and ECZTRA 3 was placebo + TCS controlled

Objective


To evaluate the efficacy and safety of tralokinumab +/- TCS versus placebo +/- TCS, by self-identified racial subgroup (Asian, Black, White) in adults with moderate-to-severe AD across three phase 3 trials

Materials and Methods

Patients and treatment

- ECZTRA 1 and 2 were two identically designed, multinational, double-blind, randomized, placebo-controlled, 52-week trials (Figure 1)
 - Patients were randomized 3:1 to subcutaneous tralokinumab 300 mg or placebo every 2 weeks (Q2W) for an initial 16 weeks following a 600 mg loading dose
 - Patients who achieved IGA 0/1 or EASI-75 at Week 16 with tralokinumab were rerandomized to tralokinumab Q2W or every 4 weeks or placebo, for an additional 36 weeks
 - Rescue treatment could be used at the discretion of the investigator to control intolerable symptoms
- In ECZTRA 3, patients were randomized 2:1 to subcutaneous tralokinumab 300 mg + TCS as needed or placebo + TCS as needed Q2W for an initial treatment period of 16 weeks following a 600 mg loading dose
 - Patients who achieved IGA 0/1 or EASI-75 at Week 16 with tralokinumab were rerandomized to tralokinumab Q2W or every 4 weeks, with TCS as need, for an additional 16 weeks
- Patients self-reported their racial subgroup

Figure 1. Summary of tralokinumab ECZTRA trials included in these analyses



Analyses

- As shown in Figure 1, data are presented as:
 - pooled from ECZTRA 1/2/3
 - pooled from ECZTRA 1/2
 - ECZTRA 3
- Efficacy outcomes assessed were:
 - Proportion of patients achieving EASI-75, IGA 0/1
 - Change from baseline in EASI, Peak Pruritus NRS, DLQI, and POEM
- Data are presented at Week 16 (ECZTRA 1/2/3) and Week 52 (ECZTRA 1/2)
- Data are presented as observed regardless of rescue medication use. Multiple imputation was used for missing data
- Data were used as per Food and Drug Administration (FDA) label and United States Prescribing Information (USPI; i.e., data from 2 sites were excluded as per FDA guidance)

Results

Patients, Demographics, and Clinical Characteristics

- This post hoc analysis included 1876 patients (USPI population) across ECZTRA 1, 2, and 3 who self-reported their race as Asian, Black, or White (Figure 1, Table 1)
- Baseline demographic and disease characteristics were largely balanced between treatment groups and across racial subgroups (Table 1)
 - AD severity was more moderate in the Black subgroup

Table 1. Baseline demographic and disease characteristics of patients by racial subgroup in pooled E1/2/3

	Asian N=407		Black N=134		White N=1335	
	Tralokinumab (n=291)	Placebo (n=116)	Tralokinumab (n=95)	Placebo (n=39)	Tralokinumab (n=992)	Placebo (n=343)
Mean age, y (SD)	35.2 (12.3)	32.4 (13.2)	39.3 (14.1)	38.6 (16.8)	39.0 (14.8)	38.7 (14.6)
Male, n (%)	176 (60.5)	76 (65.5)	38 (40.0)	20 (51.3)	578 (58.3)	209 (60.9)
Ethnicity, n (%)						
Hispanic or Latino	2 (0.7)	3 (2.6)	3 (3.2)	0	68 (6.9)	26 (7.6)
Region, n (%)						
North America	100 (34.4)	48 (41.4)	86 (90.5)	36 (92.3)	284 (28.6)	89 (25.9)
Europe	20 (6.9)	8 (8.6)	8 (8.4)	3 (7.7)	640 (64.5)	233 (67.9)
Australia	17 (5.8)	7 (6.0)	1 (1.1)	-	68 (6.9)	21 (6.1)
Japan	96 (33.0)	31 (26.7)	-	-	-	-
Asia	58 (19.9)	20 (17.2)	-	-	-	-
Country, n (%)						
United States	43 (14.8)	23 (19.8)	82 (86.3)	35 (89.7)	178 (17.9)	52 (15.2)
Canada	57 (19.6)	25 (21.6)	4 (4.2)	1 (2.6)	106 (10.7)	37 (10.8)
Japan	96 (33.0)	31 (26.7)	-	-	-	-
Patients with IGA 3, n (%)	138 (47.4)	53 (45.7)	61 (64.2)	23 (59.0)	507/1001 (50.6)	169/346 (48.8)
Patients with IGA 4, n (%)	153 (52.6)	63 (54.3)	33 (34.7)	15 (38.5)	490/1001 (49.0)	175/346 (50.6)
Mean EASI (SD), n	33.2 (15.1)	34.1 (14.2)	29.2 (13.0), 94	33.1 (15.8), 38	31.5 (13.4), 997	31.9 (13.3), 344
Mean SCORAD score (SD)	70.7 (14.7)	71.4 (12.4)	66.4 (12.4), 94	67.6 (12.7), 38	69.7 (12.8), 997	70.8 (12.8), 344
Mean DLQI (SD), n	17.5 (7.1), 289	18.0 (6.6), 114	16.9 (7.1), 94	17.2 (9.7), 37	17.3 (7.0), 984	17.4 (6.9), 342
Mean Worst Pruritus NRS (SD), n	7.8 (1.4), 288	7.8 (1.4), 116	8.1 (1.6), 94	7.7 (1.7), 37	7.7 (1.5), 991	7.9 (1.4), 342

Tralokinumab improved signs and symptoms of AD across racial subgroups at Week 16

- Across three pooled trials at Week 16, tralokinumab significantly improved efficacy outcomes in the Asian and White subgroups relative to placebo. Similar results were found in the smaller Black subgroup, although statistical significance was not reached for all endpoints vs placebo (Figures 2-3)
- Lower efficacy was observed for the Black subgroup relative to Asian and White subgroups when the monotherapy trials were pooled, driven by higher placebo response rates. In contrast, higher efficacy was observed for the Black subgroup relative to Asian and White subgroups in the TCS combination trial

Figure 2. Percentage of patients achieving EASI-75 and IGA 0/1 by racial subgroup at Week 16





Figure 3. Change from baseline in EASI, ITCH, POEM, and DLQI by racial subgroup at Week 16




Error bars show standard error.

Improvements in efficacy outcomes beyond Week 16 were consistent across racial subgroups

- At Week 52 (pooled monotherapy trials), EASI-75 was achieved in 58% of Asian patients, 66% of Black patients, and 63% of White patients (Figure 4)
 - IGA 0/1 was achieved in 32% of Asian patients, 38% of Black patients, and 38% of White patients (Figure 4)
- Improvements in efficacy outcomes after Week 16 were also observed across racial subgroups in ECZTRA 3 (TCS combination trial)

Figure 4. Percentage of patients achieving EASI-75 and IGA 0/1 by racial subgroup at Week 52 (pooled monotherapy trials)



The safety profile of tralokinumab treatment was consistent across racial subgroups

- Tralokinumab was generally well-tolerated, with a safety profile comparable to placebo and largely consistent across racial subgroups (Table 2)
- Rates of adverse events (AEs), serious AEs, and AEs leading to drug withdrawal were low in all treatment groups
- Conjunctivitis rates were lower in the Asian and Black relative to White subgroup (Table 3)

Table 2. Summary of AEs through Week 16 in ECZTRA 1/2/3 by racial subgroup

PVE	Asian (n=292)		Black (n=115)		Tralokinumab (n=94)		Placebo (n=38)		Tralokinumab (n=989)		Placebo (n=340)	
	n (%)	Rate (nE/100 PVE)	N (%)	Rate (nE/100 PVE)	N (%)	Rate (nE/100 PVE)	N (%)	Rate (nE/100 PVE)	N (%)	Rate (nE/100 PVE)	N (%)	Rate (nE/100 PVE)
All AEs	197 (67.5)	570.59	79 (68.7)	633.90	52 (55.3)	542.36	25 (65.8)	512.12	713 (72.1)	753.21	243 (71.5)	772.59
Serious AEs	6 (2.1)	6.9	1 (0.9)	2.9	2 (2.1)	7.4	2 (5.3)	27.9	25 (2.5)	8.8	13 (3.8)	17.0
Severity												
Mild	161 (55.1)	424.4	61 (53.0)	455.3	45 (47.9)	408.6	17 (44.7)	316.5	590 (59.7)	500.9	181 (53.2)	412.9
Moderate	76 (26.0)	132.2	42 (36.5)	157.7	19 (20.2)	122.5	13 (34.2)	167.6	362 (36.6)	224.4	149 (43.8)	311.4
Severe	11 (3.8)	13.8	5 (4.3)	20.8	3 (3.2)	11.1	2 (5.3)	27.9	56 (5.7)	27.8	31 (9.1)	48.2
Leading to withdrawal from trial	7 (2.4)	8.0	5 (4.3)	17.8	5 (5.3)	26.0	1 (2.6)	9.3	20 (2.0)	8.1	8 (2.4)	11.0
Outcome												
Not Recovered/Not Resolved	39 (13.4)	65.5	18 (15.7)	65.4	14 (14.9)	55.7	7 (18.4)	74.4	156 (15.8)	70.3	43 (12.6)	60.2
Recovering/Resolving	16 (5.5)	18.4	8 (7.0)	23.8	7 (7.4)	26.0	-	-	41 (4.1)	15.6	21 (6.2)	26.1
Recovered/Resolved	179 (61.3)	483.1	68 (59.1)	544.6	49 (52.1)	445.7	22 (57.9)	428.3	668 (67.5)	655.4	233 (68.5)	680.1
Recovered/Resolved with Sequelae	3 (1.0)	3.4	-	-	1 (1.1)	3.7	-	-	14 (1.4)	4.7	2 (0.6)	3.0
Unknown	-	-	-	-	2 (2.1)	11.1	1 (2.6)	9.3	20 (2.0)	7.1	3 (0.9)	3.0

Table 3. Summary of selected AEs by SOC and preferred term through Week 16 in ECZTRA 1/2/3 by racial subgroup

All AEs	Asian (n=292)		Black (n=115)		Tralokinumab Q2W (n=94)		Placebo (n=38)		Tralokinumab Q2W (n=989)		Placebo (n=340)	
	n (%)	Rate (nE/100 PVE)	N (%)	Rate (nE/100 PVE)	N (%)	Rate (nE/100 PVE)	N (%)	Rate (nE/100 PVE)	N (%)	Rate (nE/100 PVE)	N (%)	Rate (nE/100 PVE)
Infections and infestations	87 (29.8)	141.50	40 (34.8)	196.42	27 (28.7)	133.73	12 (31.6)	139.67	446 (45.1)	237.71	145 (42.6)	232.08
Viral upper respiratory tract infection	28 (9.6)	35.66	9 (7.8)	29.76	8 (8.5)	37.15	4 (10.5)	46.56	191 (19.3)	80.14	55 (16.2)	72.34
Conjunctivitis	5 (1.7)	5.75	1 (0.9)	2.98	3 (3.2)	14.86	1 (2.6)	9.31	79 (8.0)	31.24	9 (2.6)	9.04
Upper respiratory tract infection	17 (5.8)	21.86	6 (5.2)	20.83	3 (3.2)	11.14	2 (5.3)	18.62	60 (6.1)	22.07	15 (4.4)	15.07
Skin infection	2 (0.7)	3.45	2 (1.7)	5.95	1 (1.1)	3.71	1 (2.6)	9.31	14 (1.4)	5.09	10 (2.9)	10.05
Herpes simplex	5 (1.7)	8.05	1 (0.9)	2.98	1 (1.1)	3.71	-	-	17 (1.7)	6.45	4 (1.2)	5.02

Conclusions

- In this post hoc analysis, tralokinumab was well-tolerated and improved the signs and symptoms of moderate-to-severe AD, regardless of race, with further improvements up to 52 weeks of treatment
- Limitations of this analysis include disparate sample sizes across racial subgroups

Abbreviations

adj., adjusted; AE, adverse event; AD, atopic dermatitis; DLQI, dermatology life quality index; E, number of adverse events; EASI, Eczema Area and Severity Index; IGA, Investigator's Global Assessment; n, number of patients achieving the indicated metric, or with 1 event; nE, number of events; nP, number of patients; N, number of patients with recorded observation; NRS, numerical rating scale; PVE, patient-years of exposure; Q2W, every 2 weeks; SD, standard deviation; TCS, topical corticosteroids

References

1. Kaufman B, et al. *Exp Dermatol* 2018; 27: 340-357. 2. Wollenberg A, et al. *Br J Dermatol*. 2021; 184(3):437-449. 3. Silverberg J, et al. *Br J Dermatol*. 2021; 184(3):450-463.

Acknowledgements

This analysis was sponsored by LEO Pharma A/S. Medical writing according and editorial support from Alphabet Health by Clair Geary, PhD was funded LEO Pharma A/S, Ballerup, Denmark, to Good Publication Practice guidelines (<https://www.ismp.org/app3>). This work was previously presented at Fall Clinical 2022.

Disclosures

Tiffany Mayo has served as an investigator or consultant for Eli Lilly, ChemoCentryx, Pfizer, Janssen, Galderma, Bristol Myers Squibb, Acelyin, Novartis, Leo Pharma, Arcutis, and Procter and Gamble. April Armstrong has served as a research investigator and/or scientific advisor to AbbVie, Almirall, Arcutis, ASLAN, Beiersdorf, BLBMS, EPI, Incyte, Leo, UCB, Janssen, Lilly, Nimbus, Novartis, Ortho Dermatologics, Sun, Dermavant, Dermira, Sanofi, Regeneron, Pfizer, and Madmed. Leon Kircik has served either as an investigator, consultant, or speaker for AbbVie, Almirall, Amgen, Arcutis, Bausch Health Canada, Bristol Myers Squibb, Boehringer Ingelheim, Celgene, Cohesus, Dermavant, Dermira, Eli Lilly, Leo, MC2, Maruho, Novartis, Ortho Dermatologics, Pfizer, Dr. Reddy's Laboratories, Sun Pharma, UCB, Tori, and Xenoport. Jonathan I. Silverberg reports honoraria as a consultant/advisory board member from LEO Pharma and has acted as a consultant for and/or received grants/honoraria from AbbVie, AnaptysBio, Asana Biosciences, Galderma Research and Development, GSK, Glenmark, KiriKiri, LEO Pharma, Lilly, MedImmune, Merlo Therapeutics, Pfizer, PurixCore, Regeneron, and Sanofi. Andrew Blauvelt has served as a speaker (received honoraria) for AbbVie, Arcutis, Bristol-Myers Squibb, Eli Lilly and Company, Pfizer, Regeneron, Sanofi, and UCB, served as a scientific adviser (received honoraria) for AbbVie, Abcentra, Alligis, Almirall, Alumis, Amgen, AnaptysBio, Arcutis, Arena, Asana, Athenex, Bluebird Bio, Boehringer Ingelheim, Bristol-Myers Squibb, Cara Therapeutics, Dermavant, EcorR, Eli Lilly and Company, Escient, Evelo, Evmmune, Forto, Galderma, Highlight Pharma, Incyte, Inovance, Janssen, Lanxos, Leo, Merck, Novartis, Pfizer, Rapt, Regeneron, Sanofi Genzyme, Shirex Global Insights, Sun Pharma, TLL Pharmaceutical, TrialSpark, UCB Pharma, Vibromed, and Xenor, and has acted as a clinical study investigator (institution has received clinical study funds) for AbbVie, Almirall, Almirall, Amgen, Arcutis, Athenex, Boehringer Ingelheim, Bristol-Myers Squibb, Concert, Dermavant, Eli Lilly and Company, Evelo, Evmmune, Galderma, Incyte, Janssen, Leo, Merck, Novartis, Pfizer, Regeneron, Sun Pharma, and UCB Pharma. Ben Esdaile has served either as an investigator, advisor or speaker for LEO Pharma, L'Oréal, Thornton & Ross, Biodesma, Skin + Me and AbbVie. Shannon Schneider and Thomas Mark are employees of LEO Pharma. Thomas Mark owns LEO Pharma stock. Melinda Gooderham has been an investigator, speaker and/or advisor for AbbVie, Amgen, Akros, AnaptysBio, Asana, Arcutis, Arista, Bausch Health, BMS, Boehringer Ingelheim, Celgene, Dermira, Dermavant, Eli Lilly, Galderma, GSK, Incyte, Janssen, Kyowa Kirin, LEO Pharma, MedImmune, Merck, Meiji, Moonlake, Nimbus, Novartis, Pfizer, Reistone, Regeneron, Roche, Sanofi Genzyme, Sun Pharma, and UCB. Andrew F. Alexis has received grants (funds to institution) from LEO Pharma, Novartis, Almirall, Bristol Myers Squibb, Amgen, Vyne, Galderma, Valiant (Bausch Health), Cara, Arcutis, Dermavant, AbbVie, and Castle; has served as an advisory board member or consultant for LEO Pharma, Galderma, Pfizer, Sanofi-Regeneron, Dermavant, Beiersdorf, Ortho, L'Oréal, BMS, Bausch Health, UCB, Vyne, Arcutis, Janssen, Allergan, Almirall, AbbVie, Sol-Gel, Amgen, VisiQ, Eli Lilly, Swiss American, and Cutera; and a speaker for Regeneron, Sanofi-Genzyme, Pfizer, and Bristol Myers Squibb.

Scan to download a copy of this poster
Copies of this poster and its content, obtained through this QR code, are for personal use only and may not be reproduced without written permission from the authors

